

REMARKS

Claims 1-66 are pending. Claims 1-7, 10, 15-22, 31-42, 45, 50-55 and 64-66 are withdrawn from consideration as being drawn to non-elected inventions. Claims 8, 9, 23, 24, 25, 26, 28, 43, 44, 47, 56, 57, 58, 59, and 61 have been amended to better clarify aspects of the invention. Claims 11, 12, 13, 14, 27, 29, 30, 46, 48, 49, 60, 62 and 63 have been canceled without prejudice or disclaimer. New claims 67-70 have been added for consideration. Support for the new claims can be found throughout the specification and in the original claims as filed. Support for the amendments can be found throughout the specification, but in particular, on page 26, paragraphs [0070] and [0071]. No new matter has been entered by way of this amendment. Accordingly, if the new claims are entered for consideration, claims 1-10, 12, 15-26, 28, 31-45, 47, 50-59, 61, 64-70 are pending. Reconsideration of this application is respectfully requested.

Requirement for Restriction/Election

Due to the complexity of the supplemental requirement for restriction, for which Applicant provided a response on June 17, 2005, the Examiner has provided a summary of the supplemental requirement for restriction and has requested confirmation of Applicant's election of groups and the combination of agents on which Applicant wishes to pursue examination. Pursuant to the Examiner's request, Applicant confirms that Applicant wishes to elect the invention of:

Section A (the at least one first agent): Group I, claims 1, 23, 36 and 56, wherein the at least one first agent is minocycline or any tetracycline family derivative capable of crossing the blood brain barrier,

Section B (the at least one second agent, which is an anti-inflammatory): Group XVIII, claims 23, 36 and 56, wherein the at least one second agent is salicylates.

Section C (the at least one second agent, (now noted as the third agent) which is an inhibitor of glutamate induced excitotoxicity): Group XXII, claims 28 and 61, wherein the at least one second agent is memantine.

With respect to the requirement for choosing one of the four possible combinations of agents, Applicant elects the combination noted in 4) one agent of Section

A in combination with one agent of Section B in combination with one agent of Section C.

In addition, Applicant requests rejoinder of claims 15, 16, 17, 18, 31, 32, 33, 34, 50, 51, 64 and 65, with the claims currently under prosecution, since they read on the subject matter (anti-inflammatory agents) of the claims currently elected for prosecution. Applicant believes that they were inadvertently left out of the list of claims noted by the Examiner. Accordingly, rejoinder of these claims is believed to be in order and is courteously requested.

Objection to the Specification

The Examiner has objected to the specification because of the following informalities:

The word “functional” is misspelled in the title of Table 1 at the top of page 8 of the disclosure. The word “tetracycline” is misspelled on page 19, paragraph [0048], line 3 of the disclosure.

Applicant has provided a substitute Table 1 with an amended Title to reflect the correct spelling of the word “functional”. Applicant has also corrected the spelling of the word tetracycline in paragraph [0048] on page 19.

Based on the foregoing amendments, Applicant requests withdrawal of the objection.

Rejection under 35 U.S.C. §112, second paragraph

A. Regarding the term “derivative”

The Examiner has rejected claims 9, 26, 44 and 59 under 35 U.S.C. §112, second paragraph as being indefinite. In particular, the Examiner alleges that the term “derivative” of present claims 9, 26, 44 and 59 is a relative term that renders the claims indefinite. More particularly, the term “derivative” does not particularly point out the degree or type of derivation that a given compound may have in relation to the parent compound and still be considered a derivative as intended by Applicant. The Examiner alleges that Applicant has provided no specific definition for this term in the present specification. Lacking a clear meaning of the term “derivative”, the Examiner alleges that

the skilled artisan would not be reasonably apprised of the metes and bounds of the subject matter for which Applicant seeks patent protection.

Applicant respectfully traverses the Examiner's rejection and has amended the claims to delete reference to the term "derivative". The claims now recite "any tetracycline that crosses the blood brain barrier". Applicant asserts that one of skill in the art would be able to understand and ascertain the limitations of the claims, as currently amended. Withdrawal of the rejection is respectfully requested.

B. Regarding the phrase "related retrogenic degenerative neurological conditions"

The Examiner has rejected claims 8-9, 11-14, 43-44 and 46-49 as being indefinite. In particular, the Examiner alleges that the phrase "related retrogenic degenerative neurological conditions" is a phrase that renders the claims indefinite. The Examiner alleges that the corresponding disclosure does not point out the degree or type of relationship that a given retrogenic degenerative condition may have to those that are presently claimed (AAMI, MCI, CVD or AD) and still be considered a "related retrogenic degenerative neurological condition". Furthermore, Applicant has not provided any specific definition for this phrase in the present specification. Accordingly, the skilled artisan would not be reasonably apprised of the metes and bounds of the subject matter for which patent protection is desired. Moreover, the Examiner notes that the term "related retrogenic diseases" is defined as neurological conditions that present some degree of degeneration from the normal adult neurological condition. This includes measurable decline of normal cognitive, neurologic, or functional capacity.

As stated in the present application, in paragraph [0033], "The relationship between normal human development and AD has been termed "retrogenesis", and is defined as a process by which degenerative biological mechanisms reverse the order of acquisition of the same mechanisms in normal development." The application lists in table 1 the "levels of functional loss in Alzheimer's disease (AD) with corresponding developmental ages (DA) of functional acquisition" (page 8 of application). Accordingly, a retrogenic dementia can be defined as a dementia which follows this characteristic retrogenic functional clinical course in terms of sequence of loss of functional capacity.

Further in paragraph [0033], the retrogenic process “is also related to recent findings regarding the molecular biology of normal cellular development and the changes in these normal molecular processes in AD, CVD and other retrogenic dementias” (Reisberg et al., 2002, American Journal of Alzheimer’s Disease, 17:202-212, EXHIBIT A), is cited at this point in the application. This reference teaches that, “The major apparent mechanism for the retrogenic phenomenon in AD and related dementing disorders has recently been elucidated.” Activation of “mitogenic molecular markers including the mitogen-activated protein kinase (MAPK) cascade, cyclins, and cyclin dependent kinases” has been “related to the phosphorylation and hyperphosphorylation of tau, and, consequently, the development of neurofibrillary changes in AD” (Reisberg et al., 2002; page 205). Therefore, the retrogenic dementias also include the category of dementias known as the tauopathies.

Based on the foregoing, Applicant respectfully traverses the Examiner’s rejection and have amended the claims to recite “related retrogenic diseases” and asserts that given the definition for this term in the present application, as noted by the Examiner and as summarized above, one of skill in the art is capable of ascertaining what cognitive or neurologic diseases fall within the meaning of this term, and moreover, which diseases would be considered a “related retrogenic disease” as currently defined in the present specification. It is to be understood that as defined, a retrogenic disease is one that includes a measurable decline of normal cognitive, neurologic or functional capacity. Further support for the definition of this term may be found in the reference supplied as Exhibit A, which is an article published in the American Journal of Alzheimer’s Disease and Other Dementias, (Vol. 17, Number 4, July/August 2002) by the inventor of the present application, entitled: “Evidence and Mechanisms of Retrogenesis in Alzheimer’s and Other Dementias: Management and Treatment Import”. Based on the amendment to the claims, withdrawal of the rejection is respectfully requested.

Rejection under 35 U.S.C. §102(b)

The Examiner has rejected claims 8-9, 11, 13, 43-44, 46, 48, 56-60 and 62 under 35 U.S.C. §102(b) as being anticipated by Duncan (WO 02/020022 A1; March, 2002).

The Examiner alleges that Duncan teaches the administration of a tetracycline compound, such as minocycline, doxycycline or any tetracycline compound of the

formula (I) in an effective amount for the treatment of neurologic disorders, such as Alzheimer's disease, in an effective amount sufficient to down-regulate microglia expression, activation and production. The Examiner further alleges that Duncan teaches that the tetracycline compound may be administered with an agent capable of inhibiting inflammation in tissue, e.g. steroidal or non-steroidal anti-inflammatory drugs.

The Examiner has noted that present claims 8 and 43 require at least one first agent capable of inhibiting neuronal cell cycle progression at or before an early phase, at least one second agent capable of inhibiting neuronal cell cycle progression generally and optionally at least one third agent capable of inhibiting mitogenic stimulation. However, the Examiner notes that absent express disclosure by Applicant stating that the at least one first agent cannot be the same as the at least one second agent, the Examiner considers the use of minocycline or a tetracycline compound as taught by Duncan to meet both the limitation of (i) at least one first agent capable of inhibiting neuronal cell cycle progression at or before an early phase and (ii) at least one second agent capable of inhibiting neuronal cell cycle progression.

Furthermore, the Examiner alleges that regardless of whether minocycline or a tetracycline compound inhibits neuronal cell cycle progression selectively at or before an early phase or simply "generally" as Applicant has stated in claims 8 and 43, the Examiner alleges that the therapeutic endpoint is still the same, i.e., the inhibition of neuronal cell cycle progression, irrespective of what point in the cell cycle the compound actually inhibits progression. Such is further support that the at least one first agent and the at least one second agent are not required to be distinct from one another. Thus, for the purposes of examination, the Examiner has interpreted the limitations to the at least one first agent and the at least one second agent to be met by Duncan, who discloses the use of minocycline or a tetracycline compound.

In addition, while the Examiner had noted the recitation of the limitation "wherein the at least one first agent inhibits cell cycle progression...and/or reducing activated microglia-induced mitogenic stimulation" in present claim 57 (see lines 1-4) and the limitation "wherein the at least one first agent inhibits cell cycle progression at or prior to entry of a neuronal cell...and/or reducing activated microglia-induced mitogenic stimulation" in present claim 58 (see lines 1-4), the Examiner notes that such limitations

are considered a functional limitation of the agents, and, thus, fail to further limit the claimed method. Nevertheless, the Examiner alleges that because the claimed active agents are also taught by the prior art of Duncan for the same therapeutic purposes, there is no reason to doubt that the agents of the patent publication do not selectively act on cells in the S or G1 phase or that they are capable of inhibiting glutamate-induced excitotoxicity and/or reducing activated microglial-induced mitogenic stimulation, absent factual evidence to the contrary.

Lastly, the Examiner notes that while Alzheimer's disease is not expressly recited in present claim 8, absent factual evidence to the contrary, and absent any limiting definition provided by Applicant, the Examiner considers such a condition to meet Applicant's limitation of a "related retrogenic degenerative neurological condition".

Applicant's Position Regarding Duncan

Applicant respectfully traverses the Examiner's rejection and asserts that in order for a rejection under 35 U.S.C. §102(b) to be proper, the reference(s) must recite each and every element of the invention as claimed.

Applicant asserts that Duncan teaches the use of tetracyclines for treating Alzheimer's disease, Guillain Barre syndrome, adrenoleukodystrophy, Parkinson's disease and amyotrophic lateral sclerosis. Moreover, tetracyclines were used for their anti-inflammatory properties, either alone or in conjunction with another anti-inflammatory compound.

The Examiner notes that Duncan does not teach the use of an inhibitor of glutamate excitotoxicity, such as memantine, nor does Duncan teach the use of salicylates as an anti-inflammatory agent, and furthermore, Duncan does not teach the treatment of age-associated memory impairment, mild cognitive impairment, or cerebrovascular dementia. Applicant is in agreement with the Examiner on these points and thus asserts that Duncan does not teach the methods of the present invention as currently claimed. Furthermore, given the current amendments to the claims, there are now further distinct differences between the teachings of Duncan and the present application.

For example, claim 8, as currently amended recites:

“A method of treating age associated memory impairment (AAMI), mild cognitive impairment (MCI), cerebrovascular dementia (CVD) and related retrogenic diseases, comprising administering a therapeutically effective amount of:

- i) at least one first agent capable of inhibiting neuronal cell cycle progression;
- ii) at least one second agent capable of inhibiting activated microglial-induced mitogenic stimulation, wherein the at least one second agent is an anti-inflammatory agent selected from the group consisting of a non-steroidal anti-inflammatory agent (NSAIDS), a salicylate, a steroid and an immunophillin; and
- iii) at least one third agent capable of inhibiting glutamate-induced cytotoxicity, wherein the at least one third agent is selected from the group consisting of memantine, neramexane, amantadine, riluzole, MK801, ketamine, dextromethorphan, dextrophan, phencyclidine, and dexanabinol (HU-211).”

It is Applicant’s position that Duncan did not appreciate the complexity of the retrogenic diseases for which the methods of treatment are disclosed in the present application. Nor did Duncan contemplate the use of three agents having different modes of action for treating patients having such retrogenic diseases, in particular, AAMI, MCI and cerebrovascular dementia. In fact, it was only through the work of the present inventor that these treatment strategies have been taught.

Applicant asserts that while the first agent of the present invention is an agent that prevents neuronal cell cycle progression, and that tetracyclines have been shown to inhibit neuronal cell cycle progression, Applicant believes that the use of a combination of therapies may prove more beneficial in treating a retrogenic disease, such as those presently disclosed and claimed. More importantly, the combination of a first agent that inhibits neuronal cell cycle progression with a second agent that is an anti-inflammatory agent and a third agent that inhibits glutamate cytotoxicity, such as memantine, is not taught or suggested by Duncan. Moreover, Duncan does not teach or suggest the use of this combination of agents for treating age-associated memory impairment (AAMI), mild cognitive impairment (MCI) or cerebrovascular dementia.



In light of the foregoing claim amendments, Applicant respectfully requests withdrawal of the rejection.

Rejection under 35 U.S.C. §103(a)

Claims 8-9, 11-14, 23-30, 43-44, 46-49 and 56-63 are rejected under 35 U.S.C. §103(a) as being unpatentable over Duncan as applied to claims 8-9, 11, 13, 43-44, 46, 48, 56-60 and 62 for the reasons made of record above, and further in view of Lipton (WO 92/17168: 1992), Lee, *et al.* (U.S. Patent No. 6,043,224; 2000) and Gervais, *et al.* (U.S. Patent Application Publication 2005/0031651; 2005, priority to U.S. Provisional Application 60/482,214 filed June, 2003).

The Examiner notes that the differences between the Duncan reference and the presently claimed subject matter lie in that the reference does not teach:

- (i) the use of memantine as an inhibitor of glutamate-induced excitotoxicity;
- (ii) the use of salicylates as the anti-inflammatory agent; and
- (iii) the treatment of age-associated memory impairment, mild cognitive impairment, or cerebrovascular dementia.

However, the Examiner alleges that the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains because of the following:

- (i) It is acknowledged that Duncan is silent as to the use of the disclosed composition (i.e., comprising minocycline (or a tetracycline compound) and an anti-inflammatory agent) in further combination with memantine, an inhibitor of glutamate-induced excitotoxicity. However, the Examiner alleges that it was well known in the art that memantine was useful for the same therapeutic purpose of treating Alzheimer's disease by reducing neuronal damage by blocking NMDA receptor-operated channel activation by excitatory amino acids, such as glutamate-related compounds, when administered in an amount sufficient to block glutamates' effect on the NMDA receptor (see Lipton, *et al.*, page 4, lines 23-31, page 10, lines 14-17 and page 10, line 32, page 11, line 3). According to the Examiner, it would, therefore, have been obvious to a person of

ordinary skill in the art to employ memantine in combination with the composition disclosed by Duncan because each was known in the art to be successful for achieving the same therapeutic effect. The Examiner further alleges that the motivation to administer both compounds flows logically from the efficacy of each compound in treating Alzheimer's disease and other neurodegenerative diseases as demonstrated in the prior art and also because each compound has been previously administered for the same therapeutic endpoints. The Examiner alleges that in the absence of evidence to the contrary, it is generally *prima facie* obvious to use in combination two or more agents that have previously been used separately for the same purpose. See *In re Kerkhoven*, 205 USPQ1069 (CCPA).

(ii) Although Duncan broadly teaches the use of an anti-inflammatory agent, both steroidal and non-steroidal, in combination with minocycline or a tetracycline compound, the reference is silent as to the particular use of salicylates as the anti-inflammatory compound. However, the Examiner alleges that the use of salicylates in compositions effective for the treatment of neurodegenerative diseases, such as Alzheimer's disease, was well known in the art at the time of the invention. Lee, *et al.*, teaches compositions of non-steroidal anti-inflammatory agents, such as aspirin or salicylic acid or the salicylates Asacol, Disalcid, Pentasa, Salflex or Trilisate, for the treatment of Alzheimer's disease and other neurodegenerative diseases (col. 5, lines 10-16; col. 10, lines 38-42 and col. 18, lines 25-58). The Examiner thus alleges that it would therefore, have been obvious to a person of ordinary skill in the art to employ any one or more of these known salicylate-type non-steroidal anti-inflammatory agents in the composition disclosed by Duncan because each would have been reasonably expected to exert the same inflammation-reducing effects as those required by the reference.

(iii) Although the previously cited references are silent as to the treatment of conditions such as age-associated memory impairment, mild cognitive impairment or cerebrovascular dementia, such conditions were known in the art to be related to the peptide amyloid- β , as taught by Gervais, *et al.* (U.S. Patent Application Publication 2005/0031651; page 3, paragraph [0024]). In light of the fact that Duncan expressly

teaches the minocycline (or a tetracycline compound)/anti-inflammatory composition useful for the treatment of Alzheimer's disease, which he expressly states is characterized by amyloid plaques associated with upregulated microglial cell expression (see page 1, line 28, - page 2, line 2 and 23-24) and that memantine is also useful for the same therapeutic objective of treating Alzheimer's disease, the Examiner alleges that it would have been appreciated by the skilled artisan that such a combination of agents would have been useful in the treatment of other disorders associated with the amyloid peptide, such as age-associated memory impairment (synonymous with age-related cognitive decline), mild cognitive impairment or cerebrovascular dementia. Such a person would have been motivated to treat such conditions with this combination of agents because it would have been reasonable expected that such a combination would demonstrate the same or a similar level of efficacy against other amyloid peptide related diseases as it would against the treatment of Alzheimer's disease.

Moreover, age-associated memory impairment and mild cognitive impairment are further considered by the examiner to be progressive conditions that precede the development of mature Alzheimer's disease. Thus, because such a combination would have efficacy in treating the advanced condition of Alzheimer's disease, the Examiner asserts that it would also be reasonably expected that the same combination of agents would have efficacy in treating the less severe conditions directly associated to and preceding the development of advanced Alzheimer's.

The Examiner has the initial burden of establishing a *prima facie* case of obviousness. A finding of obviousness under § 103 requires a determination of the scope and content of the prior art, the differences between the claimed invention and the prior art, the level of ordinary skill in the art, and whether the differences are such that the claimed subject matter as a whole would have been obvious to one of ordinary skill in the art at the time the invention was made. Graham v. Deere, 383 US 1 (1966). Obviousness cannot be established by combining the teachings of the prior art to produce the claimed invention, absent some teaching or suggestion that the combination be made. In re Stencel, 828 F2d 751, 4 USPQ2d 1071 (Fed. Cir. 1987).

The Invention as Claimed

The present invention, as currently claimed, teaches the use of a combination of three types of agents for treating retrogenic diseases, in particular, age-associated memory impairment (AAMI), mild cognitive impairment (MCI) and cerebrovascular dementia. More importantly, the present invention teaches that the first class of agents inhibits neuronal cell cycle progression, and these may be selected from the group consisting of minocycline, any tetracycline capable of crossing the blood brain barrier, acetylsalicylic acid, any salicylate which inhibits early phase cell cycle progression, sirolimus, any sirolimus derivative capable of inhibiting early cell cycle progression, flavopiridol, ciclopirox, a paulone, indirubin, fascaplycin, olomoucine, roscovitine, Aragusterol A, valproate, N-(3-chloro-7-indolyl)-1,4-benzenedisulfamide, a farnesyl transferase inhibitor such as R115777, SCH66336 and BMS-214662, and sodium butyrate. The second class of agents may be selected from agents that inhibit activated microglial-induced mitogenic stimulation, wherein the at least one second agent is an anti-inflammatory agent selected from the group consisting of a non-steroidal anti-inflammatory agent (NSAIDS), a salicylate, a steroid and an immunophillin. The third agent is an inhibitor of glutamate induced excitotoxicity, and the agent may be selected from the group consisting of memantine, neramexane, amantadine, riluzole, MK801, ketamine, dextromethorphan, dextrophan, phencyclidine, and dexanabinol (HU-211).

The Duncan Reference

Applicant asserts that Duncan teaches the use of tetracyclines for treating neurological conditions such as Alzheimer's disease, Guillain Barre syndrome, adrenoleukodystrophy, Parkinson's disease and amyotrophic lateral sclerosis. Moreover, tetracyclines were used for their anti-inflammatory properties, either alone or in conjunction with another anti-inflammatory compound.

As noted by the Examiner, Duncan does not teach or suggest the use of an inhibitor of glutamate excitotoxicity, such as memantine. Nor does Duncan teach or suggest the use of a salicylate as the anti-inflammatory agent. Furthermore, Duncan does not teach the treatment of AAMI, MCI or cerebrovascular dementia.

Applicant asserts that Duncan does not teach or suggest the use of a composition comprising at least one first agent that is an inhibitor of neuronal cell cycle progression, such as tetracycline, combined with at least one second agent, such as an anti-inflammatory, combined with a third agent, such as an inhibitor of glutamate induced excitotoxicity, all as described in the present application, for the treatment of AAMI, MCI or cerebrovascular dementia.

The Lee et al. Reference

Lee et al. teach the use of compositions of non-steroidal anti-inflammatory agents, such as aspirin, or salicylic acid, or the salicylates Asacol, Disalcid, Pentasa, Salfex or Trilisate for the treatment of Alzheimer's disease.

Lee et al. do not teach or suggest the use of a composition comprising at least one first agent that is an inhibitor of neuronal cell cycle progression, such as tetracycline, combined with at least one second agent, such as an anti-inflammatory, combined with a third agent, such as an inhibitor of glutamate induced excitotoxicity, all as described in the present application, for the treatment of AAMI, MCI or cerebrovascular dementia.

The Lipton Reference

Lipton teaches the use of compositions containing NMDA receptor antagonists to treat neurodegenerative diseases. In particular, Lipton teaches the use of compounds including amantadine, rimantadine and memantine to treat hypoxia-ischemic encephalopathy, seizures, stroke, and AIDS dementia.

Lipton does not teach or suggest the use of a composition comprising at least one first agent that is an inhibitor of neuronal cell cycle progression, such as tetracycline, combined with at least one second agent, such as an anti-inflammatory, combined with a third agent, such as an inhibitor of glutamate induced excitotoxicity, all as described in the present application, for the treatment of AAMI, MCI or cerebrovascular dementia. In addition, Applicant asserts that the use of memantine for treating moderate to severe Alzheimer's disease was not approved by the FDA until September, 2003, following a study conducted by the present inventor, the results of which were not published until after the priority date of the present application. The article summarizing this study can

be found attached herewith as EXHIBIT K (New England Journal of Medicine April 3, 2003, pages 1333-1341).

The Gervais et al. Reference

Gervais et al. teach a method of concomitant therapy for patients suffering from diseases associated with enhanced beta amyloid peptide. The therapy includes the use of two agents, one being an agent that either blocks beta amyloid toxicity, or prevents or slows the rate of beta-amyloid formation. The second agent may be an anti-inflammatory agent or an acetyl cholinesterase inhibitor, or an agent that further prevents beta amyloid formation. Other neuroprotective second agents are also disclosed.

Gervais et al. do not teach or suggest the use of a composition comprising at least one first agent that is an inhibitor of neuronal cell cycle progression, such as tetracycline, combined with at least one second agent, such as an anti-inflammatory, combined with a third agent, such as an inhibitor of glutamate induced excitotoxicity, all as described in the present application, for the treatment of AAMI, MCI or cerebrovascular dementia.

The analysis under § 103(a).

Applicant asserts that the teachings of Duncan, when combined with the teachings of Lee, Lipton and Gervais do not teach or suggest the aspects of the present invention, as currently claimed. Moreover, it is Applicant's contention that one of skill in the art would not have been motivated to combine the teachings of Duncan with those of Lee, Lipton, and Gervais to result in the teachings of the present invention. Moreover, it is also Applicant's contention that, based on certain teachings in the art, as noted below, there would be no reasonable expectation of success if one were to combine the teachings of Duncan with those of Lee, Lipton, and Gervais.

While the Examiner alleges that absent express disclosure by Applicant stating that the at least one first agent cannot be the same as the at least one second agent, the Examiner considers the use of minocycline or a tetracycline compound as taught by Duncan to meet both the limitation of (i) at least one first agent capable of inhibiting neuronal cell cycle progression at or before an early phase and (ii) at least one second agent capable of inhibiting neuronal cell cycle progression. Applicant asserts that while the present application does not expressly state that the first and second agent cannot be

the same agent, in particular, a tetracycline, Applicant further asserts that the differences between the first and second agent, as currently claimed, is inherent in the teachings throughout the specification and the claims as filed.

As noted in the present application, (paragraphs [0049], [0050] and especially [0051] to [0053]) minocycline and other tetracycline family compounds are disclosed for use in treating AAMI, MCI and cerebrovascular dementia based on their cell cycle inhibitory effects and more particularly for their effects on early phase cell cycle progression.

Applicant notes in the application in paragraph [0051] that:

“These effects on the cell cycle likely account in part for other observed effects of minocycline including the recently uncovered effects on cerebral ischemia and inflammation (Yrjänheikki, et al. (1999) Proc. Natl. Acad. Sci. USA 96: 13496 – 13500; Yrjänheikki, et al. (1998) Proc. Natl. Acad. Sci. USA 95: 15769 – 15774).”

Duncan refers to prior art and knowledge regarding the hypothesis that anti-inflammatory treatment may be useful in Alzheimer’s disease. Duncan notes that “One proposed therapeutic approach is to minimize or prevent the inflammatory changes that occur in the brains of, e.g., Alzheimer’s patients (Selkoe 1999). It is thought that a key cell in the cellular cascade that results in neuronal death in Alzheimer’s disease is the microglial cell.” He also notes that in the prior art “It has been reported that minocycline, a tetracycline compound, reduced the microglial response in the area surrounding areas of brain ischemia in an experimental model of stroke, and that it protects neurons against death in tissue culture (Yrjänheikki, et al. 1999, 1998).”

The present application once again differs markedly from Duncan in that the present application suggests the utility of minocycline and related tetracycline compounds because of cell cycle events, not anti-inflammatory effects. Therefore, the dosage, endpoint and goal of administration are very different from those proposed by Duncan who suggests, “a method of providing ...an effective amount of a tetracycline compound sufficient to down regulate microglial expression, activation and production, and hence, prevent, reduce or minimize inflammation” (p.2, Duncan).

Regarding the Use of Anti-inflammatories eg. salicylates, for Treating Alzheimer’s Disease

Applicant respectfully traverses the Examiner’s allegations that it would be obvious to combine the teachings of Duncan on the use of tetracycline in combination with an anti-

inflammatory such as a salicylate, because it is known in the art that salicylates are useful for treating Alzheimer's disease. Applicant asserts that in fact, the anti-inflammatory approach to the treatment of Alzheimer's disease has not been successful as a first approach, which is in disagreement with the Examiner's allegations. For example, Aisen et al., found that the anti-inflammatory medications rofecoxib and naproxen do "not slow cognitive decline in patients with mild to moderate AD" (Aisen et al., 2003, EXHIBIT B). In another study of patients with mild cognitive impairment Thal et al. found that, "The results from this MCI study did not support the hypothesis that rofecoxib would delay a diagnosis of AD" (Thal et al., 2005, EXHIBIT C). Accordingly, no anti-inflammatory compounds have been approved by the FDA or other regulatory agencies for the treatment of AD, MCI, AAMI, cerebrovascular dementia or other retrogenic dementias.

Furthermore, the present application does not claim an anti-inflammatory agent as a first agent in the treatment of Alzheimer's disease (AD) or other named therapeutic entities. Accordingly, the present application is clearly different from the teachings of Duncan.

Regarding AAMI and MCI to be Progressive Conditions Leading to Alzheimer's Disease

In addition, the Examiner states (p.17), that "age associated memory impairment and mild cognitive impairment are further considered by the examiner to be progressive conditions that precede the development of mature Alzheimer's disease. Thus, because such a combination would have efficacy in treating the advanced condition of Alzheimer's disease, it would also reasonably be expected that the same combination agents would have efficacy in treating the less severe conditions directly associated to and preceding the development of advanced Alzheimer's."

The Applicant respectfully differs on each of these matters on the basis of current literature and knowledge. As noted by Wang et al., (EXHIBIT D), (2004) "it is unknown whether, and to what extent, self perceived decline is a perception of the normal aging process or might represent an early sign of dementia." Wang et al. note that most of the population based longitudinal studies have "included individuals with objective cognitive impairment and therefore were unable to address the question of whether subjective memory complaints were a sign of future dementia at an early stage when objective impairment was not evident."

Mild cognitive impairment (MCI) is also a condition which does not necessarily progress to frank dementia, nor does it necessarily progress to Alzheimer's disease. In the present application,

Applicant notes evidence from the work which the inventor has done which indicates that cell cycle inhibitors may nevertheless be useful in the treatment of this condition. For example, Roach, (EXHIBIT E), (2005) quotes Gauthier and Touchon as stating that “although individuals with MCI have an increased likelihood of progressing to frank dementia during a 10 years interval many of them remain stable and some even revert to normal during this period” (Gauthier, Touchon 2005, EXHIBIT F).

Regarding the Assumption that Medications Useful in Alzheimer’s Disease are Effective in Treating Other Retrogenic Diseases or Conditions

It should also be noted that there is universal agreement that medications which are effective for Alzheimer’s disease are not necessarily effective for MCI. Clearly, there is also universal agreement throughout the field that medications which are effective for Alzheimer’s disease are not necessarily, or even likely, to be effective for AAMI.

Even within the restricted range of Alzheimer’s disease, all of the medications which have been approved by the U.S. Food and Drug Administration have been approved for either treatment of mild to moderate Alzheimer’s disease exclusively, or, alternatively, moderate to severe Alzheimer’s disease exclusively. Furthermore, none of these medications are approved for treatment of mild cognitive impairment. In fact, despite studies in the mild cognitive impairment area, no medications have been approved at this time for the treatment of either mild cognitive impairment or age associated memory impairment.

More specifically, the cholinesterase inhibitors, donepezil, rivastigmine and galantamine have been approved exclusively for the treatment of mild to moderate Alzheimer’s disease (Physicians’ Desk Reference, pages 1197-1200; 2304- 2311; 1736- 1741 respectively, 2005, Thomson PDR). This limited approval obviously speaks volumes for the FDA’s belief that a medication approved for a specific severity range, even within the Alzheimer’s disease diagnosis, does not necessarily mean that the medication is effective for other severity ranges and certainly not for other diagnoses.

Memantine, an NMDA receptor antagonist, has been approved by the FDA for the treatment of moderate to severe Alzheimer’s disease (Physicians’ Desk Reference, pages 1288- 1291, 2005, Thomson PDR, EXHIBIT G). There has been an effort on the part of the clinical scientist to extend memantine’s approval to the range of mild Alzheimer’s disease. Recently, the

FDA rejected memantine for approval for the treatment of mild Alzheimer's disease (FDA rejects memantine for mild Alzheimer's, Alzheimer's Association 2005, EXHIBIT H).

There has been an effort to examine the efficacy of cholinesterase inhibitors in the treatment of mild cognitive impairment. Thus far, it is not clear whether any of them are effective in this disease entity. For example, Salloway et al. studied the efficacy of donepezil in a randomized placebo-controlled trial in mild cognitive impairment (Salloway et al., 2004, EXHIBIT I). They concluded that "significant treatment effects were not seen in the primary efficacy measures." However, they also concluded that, "outcome on secondary measures suggest promising directions for further evaluation of donepezil treatment in patients with MCI." (Salloway et al., Neurology 2004; 63:651-657, EXHIBIT I, quotes from the abstract page 651). Other current studies of cholinesterase inhibitors to date have either been negative or inconclusive.

Regarding the Obviousness to Combine Two Agents

The examiner also implies that if one medication is approved for an indication and another medication is also approved for that same indication, then naturally, it is obvious to everyone that both medications are better than one medication alone for that indication. Aside from errors in syllogistic logic, this reasoning in fact does not apply to Alzheimer's disease specifically, nor should it preclude patentability of the present invention. For example, if one prescribes a cholinesterase inhibitor for treating Alzheimer's disease, such as donepezil, adding another cholinesterase inhibitor does not increase efficacy. Such a procedure does however increase the potential for side effects. The reason for this is that there is a window of treatment efficacy for cholinesterase inhibitors in the approved mild to moderate Alzheimer's disease treatment range. If one exceeds this window then there is an excess of cholinergic side effects and not an increase in efficacy.

There has been one study of combining the two classes of AD treatments and examining efficacy and side effects. This study in patients with moderate to severe AD receiving stable doses of donepezil, a cholinesterase inhibitor found that memantine treatment "resulted in significantly better outcomes than placebo." The conclusion of the study was that, "these results, together with previous studies, suggest that memantine represents a new approach for the treatment of patients with moderate to severe AD" (Tariot et al., JAMA 2004; 291: 317-324, EXHIBIT J, quotes from abstract, page 317). This study did not indicate whether the combination of cholinesterase inhibitor

treatment with memantine (NMDA receptor blocker) treatment was better than either substance alone in the severity range of the study.

There is some evidence that the combination of a cholinesterase inhibitor treatment with NMDA receptor antagonism treatment may not be additive. This study suggests that both treatment modalities for Alzheimer's disease may be acting on a common disease mechanism. Specifically, De Sarno et al., in press, conclude that, "drugs in each class of therapeutic agents used for AD have the common property of increasing the regulatory serine-phosphorylation of GSK-3 within common pools of the enzyme" (De Sarno et al., *Neurobiology of Aging*, in press).

In the present application, Applicant proposes combinations of treatments which would have synergistic efficacy, based on Applicant's model of retrogenesis and cell cycle inhibition.

Accordingly, Applicant asserts that it is not obvious to combine the three treatment modalities of the present application for the reasons cited above.

Regarding Amyloid and Alzheimer's disease:

The relationship between amyloid and Alzheimer's disease is unclear at the present time. There are many who believe that the amyloid is a "scar" of the disease and does not have any particular causal relationship to AD. In the case of injury to the body, scar tissue may form. This association does not mean that the scar caused the injury. Similarly, there seems to be some association between the presence of amyloid (an amorphous proteinaceous material, comprised in part of a protein product known as beta amyloid [$A\beta$], which contains varying numbers of amino acids, e.g., either 40 or 42 amino acids) and the occurrence of Alzheimer's disease. However, "the number of $A\beta$ aggregates in the brain doesn't always correlate with the severity of dementia" (Anderson, M.W., *The Scientist*, October 25, 2004, pages 28-29). Indeed studies of the cerebrospinal fluid (CSF) have indicated either no change in $A\beta$ peptides or a reduction. Specifically, there is a "reduction in CSF $A\beta$ 42" however, "there is no change in CSF $A\beta$ 40 in AD" (Hampel and Blennow, *CSF tau and beta amyloid as biomarkers for mild cognitive impairment*, *Dialogues in Clinical Neuroscience*, 2004, Vol. 6, No.4, pages 379-390, quote from page 383). Naturally, if $A\beta$ is related to AD one would expect an increase in the CSF content of this substance rather than what is observed, namely, no change or a reduction. There are various speculations as to possible reasons for the observed findings.

The Examiner's attention is drawn to the fact that no treatments for Alzheimer's disease based on lowering, raising, or otherwise changing beta amyloid, or any other element of amyloid plaques, have been approved in the U.S. or anywhere else at the present time.

Regarding Amyloid and MCI

Studies of the relationship between the diagnosis of mild cognitive impairment (MCI) and A β have been inconsistent. For example, in the CSF studies, "one study found significant decrease in the CSF A β 1-42 in MCI subjects compared with controls. In another study in MCI patients who eventually developed AD, however, A β 1-42 levels did not differ significantly from age matched controls." (Hampel and Blennow, CSF tau and beta amyloid as biomarkers for mild cognitive impairment, Dialogues in Clinical Neuroscience, 2004, Vol. 6, No.4, pages 379-390, quote from page 383). Again, intuitively, one would expect an increase in A β CSF levels, if there were a direct causal relationship between beta amyloid and MCI. However, what is observed is no change or a decrease.

The Examiner's attention is drawn to the fact that no treatments for MCI based on lowering, raising, or otherwise changing beta amyloid, or any other element of amyloid plaques, have been approved in the U.S. or anywhere else at the present time.

Aging is believed to be associated with an increase in beta amyloid plaques in the brain. Little or nothing is presently known regarding the association between beta amyloid and AAMI.

The Examiner's attention is drawn to the fact that no treatments for AAMI based on lowering, raising, or otherwise changing beta amyloid, or any other element of amyloid plaques, have been approved in the U.S. or anywhere else at the present time.

The relationship between vascular dementia treatment and treatment of Alzheimer's disease.

The Examiner states that it is obvious that if a treatment is useful for Alzheimer's disease then it should be useful for vascular dementia. Logically, of course, this is not necessarily true. The literature also does not support this conclusion. For example, a prevention study of non steroidal anti-inflammatory drugs (NSAIDs) found that, "the long term use of NSAIDs may protect against Alzheimer's disease but not against vascular dementia" (Veld, et al., NEJM, 2001; Vol. 345:1515-21, quote from abstract page 1515).

The relationship between a medication which prevents an illness and a medication which treats the illness.

The examiner implies that a medication which treats an illness such as Alzheimer's disease is likely to be useful in preventing the condition.

Applicant respectfully asserts that this is not true. For example, in Alzheimer's disease, five medications have been approved for treatment in the U.S. None of these medications have been approved for preventing Alzheimer's disease. The same is true for other conditions. For example, another brain disease is schizophrenia. Many medications have been approved for the treatment of schizophrenia, but none of these medications have been approved for the prevention of schizophrenia.

Based on the foregoing, Applicant asserts that the present invention is still patentable over Duncan in view of Lee, Lipton and Gervais for at least the following reasons:

1. There simply is no teaching or suggestion to use a combination of three agents, one being a cell cycle inhibitor, one being an anti-inflammatory and the third being an NMDA antagonist, having the noted characteristics as currently claimed, for treating a retrogenic disease.
2. There simply is no teaching or suggestion to use a combination of three agents, one being a cell cycle inhibitor, one being an anti-inflammatory and the third being an NMDA antagonist, having the noted characteristics as currently claimed for treating AAMI, MCI and cerebrovascular dementia.

Based on the foregoing, withdrawal of the rejection is respectfully requested.


Fees

A check in the amount of \$325. is enclosed to cover the fee for three new independent claims and one new dependent claim over 20. No other fees are believed to be necessitated by the foregoing response. However, if this is in error, authorization is hereby given to charge Deposit Account No. 11-1153 for any underpayment, or credit any overages.

Conclusion

Applicants believe that the foregoing cancellation and amendments to the claims place the application in condition for allowance. Withdrawal of the rejections and objections is respectfully requested. If a discussion with the undersigned will be of assistance in resolving any remaining issues, the Examiner is invited to telephone the undersigned at (201) 487-5800, ext. 118, to effect a resolution.

Respectfully submitted,



Veronica Mallon, Ph.D.
Agent for Applicants
Registration No. 52,491

KLAUBER & JACKSON
411 Hackensack Avenue
Hackensack, New Jersey 07601
(201) 487-5800

Attachments: References listed as EXHIBITS A THROUGH K

Reprinted from

American Journal of Alzheimer's Disease® and Other Dementias

Volume 17, Number 4

July/August

2002

Evidence and mechanisms of retrogenesis in Alzheimer's and other dementias: Management and treatment import

Barry Reisberg, MD
Emile H. Franssen, MD
Liduïn E. M. Souren, RN, MSN
Stefanie R. Auer, PhD
Imran Akram, MD
Sunnie Kenowsky, DVM



American Journal of Alzheimer's Disease and Other Dementias ISSN 1533-3175 is published bimonthly by Prime National Publishing Corp., 470 Boston Post Road, Weston, MA 02493, 781-899-2702. Subscription price per volume: US Individual, \$136, US Institution/Library, \$193; Canadian Individual, \$172; Canadian Institution/Library, \$245; Foreign Individual, \$199; Foreign Institution/Library, \$304. Single issues: United States, \$35; Canada, \$40; Foreign, \$50 (Note: Prices are according to US currency). To subscribe, submit your complete name, address and zip code, attention: *American Journal of Alzheimer's Disease and Other Dementias*, Circulation Department, 470 Boston Post Road, Weston, MA 02493. Please enclose check, purchase order or credit card with authorization signature.

All rights reserved. Authorization to photocopy items

for internal or personal use of specific clients is granted by Prime National Publishing Corp., provided the appropriate fee is paid directly to Copyright Clearance Center (CCC), 222 Rosewood Drive, Danvers, MA 01923, USA, (978) 750-8400. CCC should also be contacted prior to photocopying items for educational classroom use. Multiple reprints of material published in *American Journal of Alzheimer's Disease and Other Dementias* can be obtained by filling out a reprint order form.

Printed in the United States of America.

©Copyright 2002 by Prime National Publishing Corp.

Disclaimer: The publisher and editors are not responsible for any opinions expressed by the authors of articles published in *American Journal of Alzheimer's Disease and Other Dementias*.

Evidence and mechanisms of retrogenesis in Alzheimer's and other dementias: Management and treatment import

Barry Reisberg, MD
Emile H. Franssen, MD
Liduín E.M. Souren, RN, MSN
Stefanie R. Auer, PhD
Imran Akram, MD
Sunnie Kenowsky, DVM

Abstract

Retrogenesis is the process by which degenerative mechanisms reverse the order of acquisition in normal development. Alzheimer's disease (AD) and related conditions in the senium have long been noted to resemble "a return to childhood." Previously, we noted that the functional stages of AD precisely and remarkably recapitulated the acquisition of

the same functional landmarks in normal human development. Subsequent work indicated that this developmental recapitulation also applied to the cognitive and related symptoms in AD. Remarkably, further investigations revealed that the same neurologic "infantile" reflexes, which mark the emergence from infancy in normal development, are equally robust indicators of corresponding stages in AD. Neuropathologic and biomolecular mechanisms for these retrogenic processes are now evident. For example, the pattern of myelin loss in AD appears to mirror the pattern of myelin acquisition in normal development. Also, recent findings indicate that mitogenic factors become reactivated in AD, and, consequently, the most actively "growing" brain regions are the most vulnerable. Because of this robust retrogenic process, the stages of AD can be translated into corresponding developmental ages (DAs). These DAs can account for the overall management and care needs of AD patients. A science of AD management can be formulated on the basis of the DA of the Alzheimer's patient, taking into consideration differences of AD from normal development as well as homologies.

Key words: Alzheimer's disease, care concepts, dementia, dementia management, retrogenesis

Historical background

Retrogenesis can be defined as the process by which degenerative mechanisms reverse the order of acquisition in normal development.¹ General relationships

Barry Reisberg, MD, Director, Zachary and Elizabeth M. Fisher Alzheimer's Disease Education and Resources Program; Clinical Director, William and Sylvia Silberstein Aging and Dementia Research Center; and Professor, Department of Psychiatry, New York University School of Medicine, New York, New York.

Emile H. Franssen, MD, Zachary and Elizabeth M. Fisher Alzheimer's Disease Education and Resources Program and the William and Sylvia Silberstein Aging and Dementia Research Center, New York University School of Medicine, New York, New York.

Liduín E.M. Souren, RN, MSN, Zachary and Elizabeth M. Fisher Alzheimer's Disease Education and Resources Program and the William and Sylvia Silberstein Aging and Dementia Research Center, New York University School of Medicine, New York, New York.

Stefanie R. Auer, PhD, Science and Development, Verein M.A.S. Bad Ischl, Austria.

Imran Akram, MD, Zachary and Elizabeth M. Fisher Alzheimer's Disease Education and Resources Program and the William and Sylvia Silberstein Aging and Dementia Research Center, New York University School of Medicine, New York, New York.

Sunnie Kenowsky, DVM, Zachary and Elizabeth M. Fisher Alzheimer's Disease Education and Resources Program, New York University School of Medicine, New York, New York.

between aging and development have long been noted by poets and playwrights,^{2,3} and are incorporated in the vocabulary of the English and other languages. More than two centuries ago, physicians also began to note similarities between aging and normal infant-child development.⁴ In the 1960s and 1970s, de Ajuriaguerra and associates observed that the decline of certain capacities in dementia appeared to reverse Piaget's developmental stages.⁵⁻⁷

Functional retrogenesis

Systematic empirical studies of the symptomatology of brain aging and progressive Alzheimer's disease (AD) in the 1980s resulted in the description of 16 successive functional stages and substages that were incorporated in the *functional assessment staging* (FAST) procedure.^{8,9} Simultaneously, this functional progression was recognized as reversing the order of acquisition of the same functions in normal human development (Table 1).¹⁰ Also immediately observed was that this functional progression was characteristic, but not pathognomonic, of the dementia of AD.^{11,12}

One concern of subsequent research was validating this characteristic functional progression of AD. Strong correlations with existing dementia measures (e.g., the Mini-Mental State Exam [MMSE]¹³) were immediately apparent;⁸ however, the final six to eight FAST substages occurred after the MMSE produced floor effects. Consequently, new measures needed to be developed for assessment of cognition in severe dementia, and the latter portion of the FAST had to be validated against these new measures. This was accomplished over the next decade.¹⁴ Other investigations concerned the "criterion validity" of the FAST progression of AD. The FAST progression showed unprecedented strong relationships to neuropathologic post-mortem assessments of hippocampal volume loss,¹⁵ cell loss,¹⁶ and neurofibrillary changes¹⁶ in AD. In another "criterion validity" study of the prospectively charted longitudinal course of AD, the FAST accounted for approximately twice the variance in temporal course of that explained by the MMSE.¹⁷

Consequently, these investigations indicated that:

- A characteristic functional course of AD was identifiable with the FAST staging procedure; and
- This characteristic course reversed normal human functional developmental acquisition.

Cognitive retrogenesis

The FAST staging procedure was developed from efforts

to systematically describe optimally concordant ordinal assessments of progressive changes in AD.¹⁸ These efforts, in turn, followed from the development of the Global Deterioration Scale (GDS).¹⁹ One of the concordant ordinal assessments developed was *praxic capacity*.²⁰ Studies demonstrated that the evolution of praxic deficit in AD followed the retrogenic pattern.²¹ Furthermore, if the FAST stages of AD and the corresponding praxis stages of loss in AD were each translated into corresponding developmental ages (DAs), *the loss of praxic capacity in AD occurred, on average, at the same DA as the loss of functional capacity*. In other words, cognitive retrogenesis in AD could be shown as concordantly mirroring functional retrogenesis.

Research from other centers also demonstrated the phenomenon of cognitive retrogenesis in AD. For example, a study of the MMSE, a measure developed for the assessment of cognition in AD and other dementias, was conducted in normal children. The investigators found an equally robust relationship between MMSE scores and mental age (MA), based on IQ scores in children, as has been observed between MMSE scores and any independent, objective dementia assessment in AD.²²

We adapted a Piagetian test battery, the Uzgris and Hunt Ordinal Scales of Psychological Development, for use in assessment of residual cognition in AD²³ and observed very strong relationships ($r = 0.8$) between the resulting measure (denoted M-OSPD) and FAST measurements of functional deterioration in AD.¹⁴ Thornbury compared performance on Piagetian tasks by AD patients with MMSE scores and obtained a similar correlation (i.e., $r = 0.8$) to that which we have noted with FAST scores and Piagetian measures.²⁴

Therefore, we determined the following:

- Cognitive retrogenesis appears to be as robust as functional retrogenesis. Specifically, *dementia measures appear to be just as good at measuring normal human development as they are at measuring dementia. Conversely, infant-child developmental measures appear to be just as good at measuring dementia in AD as in measuring cognitive development*; and
- Cognitive retrogenesis appears to occur, from certain perspectives, concordantly with AD-based functional retrogenesis. It appears that *cognition and functioning are linked in the same intimate qualitative and quantitative relationship in dementia as they are in normal development*.

Emotional retrogenesis

Once the functional and cognitive landmarks of

Table 1. Functional landmarks in normal human development and Alzheimer's disease*

Normal development (approximate total duration: 20 years) ↑	Approximate age		Approximate duration in development	Acquired abilities	Lost abilities	Alzheimer stage	Approximate duration in AD	Developmental age of patient	Alzheimer's degeneration (approximate total duration: 20 years) ↓
	Adolescence	13 – 19 yrs	7 yrs	hold a job	hold a job	3 (incipient)	7 yrs	19 – 13 yrs (adolescence)	
	Late childhood	8 – 12 yrs	5 yrs	handle simple finances	handle simple finances	4 (mild)	2 yrs	12 – 8 yrs (late childhood)	
	Middle childhood	5 – 7 yrs	2½ yrs	select proper clothing	select proper clothing	5 (moderate)	1½ yrs	7 – 5 yrs (middle childhood)	
	Early childhood	5 yrs 4 yrs 4 yrs 3 – 4½ yrs 2 – 3 yrs	4 yrs	put on clothes unaided shower unaided toilet unaided control urine control bowels	put on clothes unaided shower unaided toilet unaided control urine control bowels	6a (moderately severe) b c d e	2½ yrs	5 – 2 yrs (early childhood)	
	Infancy	15 mo 1 yr 1 yr 6 – 10 mo 2 – 4 mo 1 – 3 mo	1½ yrs	speak 5 – 6 words speak 1 word walk sit up smile hold up head	speak 5 – 6 words speak 1 word walk sit up smile hold up head	7a (severe) b c d e f	7 yrs	15 mo – birth (infancy)	

*Copyright © 1984, 1986, 2000 by Barry Reisberg, MD. All rights reserved.

degeneration in AD were established and the DAs corresponding to these landmarks were identified, other aspects of the retrogenic process could be studied. In a series of studies, we identified characteristic behavioral and psychological symptoms of dementia and the stages in the evolution of dementia at which these symptoms emerged.²⁵⁻²⁷ Interestingly and importantly, many of these symptoms occurred at precisely the point that would be anticipated from the corresponding DA of the AD patient.^{28,29}

Neurologic retrogenesis

Neurologic release signs, also referred to as primitive, developmental, or neonatal reflexes because of their occurrence in normal infancy, have long been known to

occur in elderly persons and dementia patients.³⁰⁻³³ Systematic studies precisely measured these reflexes and established the precise stages in the evolution of AD at which they emerged.³⁴⁻³⁶ *These studies indicated that many of the same reflexes that are used to mark the emergence from normal infancy to early childhood appear to be equally robust markers of the emergence of the infantile DA stage of AD (i.e., movement from the end of FAST stage 6, into FAST stage 7).*³⁶

Neuropathologic retrogenesis

A neuropathologic pattern of degeneration in AD, which appeared to reverse the normal human developmental pattern, was first observed by Brun and Gustafson a quarter century ago.³⁷ Subsequently, at least three

neuropathologic pattern in AD in terms of neuronal cell loss,³⁸ neurometabolic change,³⁸ neurofibrillary change,³⁹ and myelin vulnerability.^{39,40}

Biomolecular retrogenesis

The major apparent mechanism for the retrogenic phenomenon in AD and related dementing disorders has recently been elucidated. The response of the neuron to stressors, including beta-amyloid, is an attempt to regenerate.^{41,42} The result of this mitogenic stimulus is the activation of various mitogenic molecular markers, including the mitogen-activated protein kinase (MAPK) cascade, cyclins, and cyclin-dependent kinases.⁴³⁻⁵³ Activation of MAPK in the hippocampus has been shown to be linked to contextual and spatial memory formation in the hippocampus.⁵⁴⁻⁵⁷ Activation of these factors in some cases appears to precede the development of neuronal neurofibrillary changes.^{45,46} Some of these mitogenic factors have been related to the phosphorylation and hyperphosphorylation of tau, and, consequently, the development of neurofibrillary changes in AD.^{43,58,59} Reactivation of mitogenic factors has also been related to apolipoprotein (APP) processing into amyloidogenic elements.⁶⁰

Consequently, this biomolecular research in AD indicates that *the most metabolically active regions of the brain in AD, which are the most capable of responding to a mitogenic stimulus, are the most vulnerable in AD.*^{48,49} These phenomena clearly appear to account for the neuropathologic, neurologic, cognitive, functional, and emotional retrogenic phenomena reviewed in the preceding sections.

Risk factors for dementia and retrogenesis: Arboreal entropy

Some AD and dementia pathogenic factors currently have been associated with mitogenic factor activation. These include hypoxia and beta-amyloid (1-42).⁴² However, other risk factors for AD have, at this time, been related primarily to myelin disturbance. These include low vitamin B12 levels, folate deficiency, increased serum homocysteine levels, and increased serum methylmalonic acid levels.⁶¹⁻⁶³ Atherosclerosis and cerebrovascular disease are other risk factors associated with AD, which have been primarily associated with myelin disruption.⁶⁴⁻⁶⁶

Therefore, a full understanding of the retrogenic process in AD probably encompasses myelin as well as mitogenic neuronal reactivation. The process of myelination is now known to continue well into the latter portion of life. Neurons that are myelinated early in the life become

Possibly, myelin plays a role not only in the conduction of electrical impulses in the neuron, but also in protection and maintenance of the oligodendroglial, myelin, and axonal relationship.^{1,67} Consequently, the most recently affected, and, as a result, most thinly myelinated brain regions, may be the most vulnerable to injury.⁶⁷

A terminology that appears to describe the pattern of neuronal vulnerability accounting for the retrogenic observations is *arboreal entropy*. Just as the bark of a tree protects it from external injury and, to some extent, the thicker the bark, the greater the protection, the myelin protects the axon and its neuron. Hence, to some extent, we may say the thicker the myelin, the greater the protection. However, just as a tree can also be attacked from the inside, which ultimately rots the bark, the axon can be attacked from the inside. For example, neurofibrillary changes and neurotubular changes, secondary to mitogenic factor activation and tau hyperphosphorylation, can result in axonal destruction and resultant myelin loss. Note that the myelin is a living, metabolically active part of the axon of the neuron, with a membrane running through it, which is an extension of the cell (axonal) membrane. As a metabolically active cellular component, myelin is vulnerable to cellular and axonal injury.

Retrogenesis in non-AD dementias

At present, it is clear that the biomolecular factors and myelin vulnerability factors, which produce the phenomenon of retrogenesis in AD, frequently operate in other dementia disorders as well.

For example, investigators who have studied the M-phase mitotic markers cyclin B1 and cdc 2 in diverse dementing disorders have found increased activity in a variety of dementias, including, in addition to AD, Down's syndrome, frontotemporal dementia linked to chromosome-17, amyotrophic lateral sclerosis (ALS) dementia complex of Guam, Niemann Pick disease type-C, progressive supranuclear palsy, corticobulbar degeneration, and Pick's disease.⁵³ A common feature in most of these dementias is the occurrence of hyperphosphorylated tau, although not in a form necessarily associated with AD type neurofibrillary tangles. In contrast, these investigators found little or no increased activity of these particular M-phase mitotic markers in Parkinson's disease (PD), Huntington's disease (HD), ALS, multiple sclerosis (MS), pituitary adenoma, AIDS dementia, and fungal infection.⁵³

Also, conditions such as vitamin B12 deficiency and hyponatremia clearly can produce myelin disturbances that may result in dementing conditions.

The extent to which non-AD dementias show a retrogenic pattern of progressive clinical pathology is likely

dependent, in part, on the occurrence of retrogenic biomolecular and neuropathologic phenomena.

Management import of retrogenesis in AD

*An understanding of the phenomenon of retrogenesis in AD permits the foundation of a new science of AD care.*²⁹ This science is based on fundamental human needs, which apply to humans at all ages, and an understanding of the DA of the AD patient. Fundamental human needs include those for movement, socialization, love, and dignity. Typically, these fundamental needs are frequently not understood in the AD patient.

For example, in nursing homes in the United States, until recently, late-stage AD patients were routinely put into "geri-chairs," which prevented ambulation, to prevent falling. The situation would be analogous to locking infants in chairs to prevent falling. The AD patients commonly developed deformities, probably in large part because of their restricted movement.⁶⁸ Infants who were similarly restricted also would probably develop deformities.

To cite another example, although everyone knows infants need overt expressions of love, love is not frequently provided to AD patients. This is especially true of late-stage AD patients in nursing homes.

In many ways, the situation today is quite similar to the 18th century, when psychiatric patients were kept in chains and there was little understanding of mental illnesses. An understanding of retrogenesis provides a basis for more sophisticated principles of care.

A new science of AD care based on retrogenesis

The new science of AD care can be formulated into axioms, postulates, and caveats. *Axioms* in this context are self-evident basic human needs and desires, applicable at all ages. *Postulates* are testable hypotheses of AD patient care. Treatment, based on the postulates, is dependent on satisfaction of the axioms. *Caveats* are exceptions to the DA-retrogenesis model, based on the nature of human aging and AD. The science of AD management is the clinical science and art of AD patient care, based on the interaction of the axioms, postulates, and caveats.

Care axioms

Examples of the axioms of AD management include the following:

Axiom I: All human beings avoid trauma and humiliation. AD patients; at any stage; avoid or rebel against

experiences that are perceived as humiliating. The most prominent humiliating experience for the AD patient is appearing "stupid." Therefore, even early in the disease process, patients may avoid being questioned. As the disease progresses, patients may resist the humiliation of requiring a caregiver. So-called "delusions," such as that people are stealing things, have a psychologic basis insofar as AD patients at these stages prefer to accuse others of taking things, rather than accept the humiliation of admitting to themselves or others that they cannot remember.

Axiom II: All human beings seek a sense of accomplishment. In early AD (GDS stage 4), this sense of accomplishment can come from continued productivity. For example, an artist may continue to paint. A lawyer may continue the pretense of working on cases. One judge had his daughter write opinions. GDS stage 5 patients may continue to insist that they are working, even if they have been forced to retire. Later in AD, a sense of accomplishment may come from folding towels and other simple DA-appropriate activities.

This axiom has important corollaries (i.e., consequences or results which follow from the axiom), which are:

- All human beings resist losses;
- A sense of accomplishment can be fostered by beginning with what an AD patient can do and building upon this; and
- A sense of accomplishment comes from practicing an area of residual expertise or learning new things.

Axiom III: All human beings seek a sense of dignity and self-worth. This dignity and self-worth may come from practicing previously mastered skills. It may also come from optimal participation in "adult" activities. Or, it may also be fostered by introducing necessary caregivers as "friends." A corollary of this is that if an AD patient perceives an activity as "infantile" or "childish" and, therefore, as an affront to his or her dignity, they may become angry and refuse to participate.

Axiom IV: All human beings are social organisms. Therefore, the social needs of the AD patient remain throughout the illness process. Even in the late stage (GDS and FAST stage 7), patients continue to require interaction with caregivers and others for mental and physical health and well-being.

Axiom V: All human beings seek praise and acceptance. As social organisms, AD patients continue to require positive social reinforcement throughout the course of the illness process in order to maintain their motivation and skills.

learn. One aspect of this capacity to learn is that AD patients can be retrained in many of the skills they have lost by breaking the tasks down into small stages, which are achievable, and praising the patient for his or her accomplishments.

Axiom VII: *All human beings require love.* This is necessary for the emotional and physical health of the AD patient at all points in the illness process.

Axiom VIII: *All human beings have the capacity for happiness if basic needs are fulfilled.* This means that AD is a physiologically congruent process. As such, if proper care is provided and social, emotional, and other needs are met, AD patients need not suffer and they can derive satisfaction from their existence.

Axiom IX: *All human beings have the need for physical movement.* Indeed, movement is sometimes said to be a fundamental feature of animal life. As is the case of the other axioms, this fundamental need is frequently ignored or not recognized in AD patients to such an extent that, until recently, AD patients were routinely restrained in order to prevent falling. Naturally, this restraint actually increased falls in patients, who were made increasingly unstable from the restraints. Hence, the need for movement remains frequently unrecognized in the AD patient.

Axiom X: *All human beings have the capacity to remember.* As is true of many of the other axioms, this basic human capacity is frequently not recognized, particularly in the late-stage (GDS and FAST stage 7) AD patient. If the person's memory is placed in the context of the DA of the AD patient, his or her memory capacity becomes comprehensible and assessable. Just as a one-year-old child will forget people and events more quickly, a stage 7 AD patient will forget people and events more quickly than a healthy adult. Three weeks is a very long time for a stage 7 AD patient or a one-year-old child. As is true of all humans at all ages, emotional memories are particularly strong in AD.

Axiom XI: *All human beings have the capacity to think.* As with other basic human capacities, this one is sometimes not recognized in the AD patient—with deleterious consequences. For example, caregivers may sometimes speak unflatteringly about the late stage 6 and early stage 7 AD patient, thinking that, because the patient cannot articulate, he or she does not understand. The patient may become agitated in response to the caregiver's unempathetic comments. Unfortunately, the caregiver, not recognizing the patient's comprehension, may think the agitation is sporadic and arbitrary, rather than a response to the crude and critical remarks.

Axiom XII: *All human beings seek to influence their environment.* This basic human need and desire is also frequently underestimated in the stage 7 AD patient. For

pret a stage 7 AD patient's babbling, the patient seeks to make their desire known and many become agitated, or violent, if not "listened to."

Axiom XIII: *All human beings have a sense of "taste," i.e., likes and dislikes.* Just as infants will throw away a toy they do not like, an AD patient has preferences that are expressed and can be interpreted at any stage. These preferences are sometimes not recognized. For example, a case manager recommended "any ID bracelet" for an early stage 6 AD patient because "she won't notice." In actuality, this patient would have been highly insulted by an unattractive plastic bracelet and wished to maintain a sense of style and beauty.

Care postulates

The postulates are based on the retrogenesis observations and the DA model of the stages of AD, reviewed in the preceding sections. The validity of the postulates is subject to scientific investigation. A summary of some of the major postulates follows.

Postulate I: *The magnitude of care and supervision required by an AD patient, at a DA, is mirrored by the amount of care and supervision required by a child or infant at the corresponding DA.* For example, an AD patient at GDS and FAST stage 7, corresponding to an infancy DA, requires approximately the same amount of care and supervision as an infant.²⁹

Postulate II: *The kinds of activities enjoyed by an AD patient, at a particular DA, are mirrored by the kinds of activities enjoyed by children at a corresponding DA.* For example, just as child of age two to five years may enjoy working on puzzles, drawing with crayons, or assisting with simple household chores, similar activities are appropriate for the GDS and FAST stage 6 AD patient at a corresponding DA.

Corollaries of postulate II include the following:

- The kinds of activities that children find frightening or upsetting at a DA are mirrored by the kinds of activities AD patients find upsetting at a corresponding DA;
- The kinds of activities that a child considers "childish" or "babylike" at a particular DA, are mirrored by the kinds of activities an AD patient may find humiliating; and
- The kinds of activities that promote healthy and optimal motor development in children are similarly the kinds of activities which minimize motor degeneration in AD.

form in an area of residual expertise is dependent on the patient's DA. If a child at the particular DA can master the task, an AD patient at the corresponding DA potentially can retain the residual capacity. Variability in AD patients' loss of capacity is mirrored by a corresponding variability in children's ability to master the task. Therefore, expectations must be in accord with the DA of the AD patient. A corollary of this postulate is that as an AD patient approaches loss of capacity, they develop anticipatory anxieties. For example, AD patients commonly develop anxieties regarding toileting in FAST stage 6c, prior to loss of urinary continence in FAST stage 6d. Similarly, some patients develop anxieties about completing their income tax in GDS and FAST stage 4.

Postulate IV: Previous experiences may determine the kinds of activities enjoyed by an AD patient. For example, one GDS stage 6, FAST stage 6d AD patient was very anxious about having her front yard clean. Naturally, an urban patient might not be interested in cleaning a yard.

Postulate V: The emotional level of the AD patient is dependent on the DA. For example, just as a two to three year-old child may become "cranky" at various points in the day, a stage 6d AD patient at a corresponding DA may develop "mood swings."

Postulates VI: Life experiences appropriate to the DA become most relevant for AD patients at any particular stage. For example, a man's only interest may have been his work. However, at stage 5 (DA of five to seven years), much of the interest in his previous job (e.g., business, medicine, etc.) is lost. At this point, DA-appropriate activities can be introduced.

Postulate VII: Socialization of the AD patient is dependent on the DA. For example, just as an infant of a year old or less does not yet relate socially with other infants, a GDS and FAST stage 7 AD patient does not relate or socialize with other stage 7 patients.

Postulate VIII: Diversity in children's and infants' activities and interests is mirrored in diversity in AD patient's interests and activities at a corresponding DA. A corollary of this is that just as all normal, healthy, children at a given age clearly can be shown to have much in common, despite acknowledged diversity, all "uncomplicated" AD patients at a given stage have much in common, despite acknowledged diversity. For example, all normal, healthy, one-year-old children have much in common with each other in comparison with normal, healthy, 12-year-old children. Conversely, all GDS stage 7, FAST stage 7a, uncomplicated AD patients have much in common, as compared to all uncomplicated stage 4 AD patients.

at a DA are mirrored by the emotional changes observed in children at a corresponding DA. This is similar to postulate V, but concerns specifics. For example, delusions occur in AD patients that are very similar to childhood fantasies at the corresponding DAs.²⁷⁻²⁹

*Postulate X: Care settings appropriate to AD patients at a DA are mirrored by care settings appropriate to children at the corresponding DA.*²⁹ A corollary of this postulate is that just as institutions would be considered an inappropriate and undesirable care setting for infants and toddlers, institutions are inappropriate settings for the care of GDS and FAST stage 7 and stage 6 AD patients. Therefore, resources currently devoted to nursing home care probably should be shifted to care in community residences.

The analogy of nurseries and schools for children to day-care centers for AD patients also applies in this regard.

Postulate XI: Vulnerability (emotional, physical, and cognitive) of the AD patient at a DA is mirrored by the vulnerability of children at the corresponding DA. For example, just as an infant is vulnerable to social deprivation, poor care, and physical insult, a stage 7 AD patient is vulnerable to social deprivation, poor care, and physical insult. The result of these social, emotional, and physical injuries is excess disability.

Postulate XII: The need of an AD patient for physical movement is mirrored by the corresponding DA. A young child requires more than simply walks or running back and forth for optimal physical growth and attainment. Motor development in children is also dependent on the child dressing, developing eating skills, ball playing, and other skills. The same kinds of skills are required by the AD patient to prevent precipitous physical decline.

Postulate XIII: Just as one judges development in an infant or child by what the infant or child can do and has achieved, not by what the infant or child cannot do, the AD patient at any particular DA should be assessed in terms of his or her residual skills and accomplishments, what they have learned and relearned, not by what they cannot do.

Postulate XIV: The developmental analogy is sufficiently strong to trigger DA-appropriate childhood memories, beliefs, and anxieties in the AD patient. For example, most FAST stage 6c AD patients will state that their parents are alive. These DA-appropriate memories are the basis of the statement, incorporated in the Blessed *et al.* dementia scale, that AD patients tend to dwell in the past.⁶⁹

Postulate XV: The language changes of the AD patient are mirrored by the DA. For example, when

speech abilities break down in the AD patient at FAST stage 6c, patients commonly develop "verbigeration" and neologisms, which are very similar to the babbling of infants as they acquire speech at an equivalent DA.

Care caveats

Caveats which modify the retrogenic-DA model of AD make the care and management of the AD patient a complex art as well as a science. Some of the major caveats are enumerated below.

Caveat I: Development in infants and children is accompanied by increasing expectations, whereas AD at all stages is accompanied by progressively diminished expectations. These contrasting phenomena are accompanied by widely divergent social consequences. For example, the societal tendency is to praise children and to become frustrated with AD patients. However, growth, within the context of an individual's capacity for growth, is dependent on praise for one's accomplishments.

Caveat II: AD patients experience developmentally analogous brain changes; however, they do not undergo developmentally analogous physical changes. Therefore, AD patients are physically larger and more formidably appearing than children. Furthermore, until stage 7, AD patients have the physical habitus of a normal adult. These physical features produce special consequences for AD patients in comparison with children. For example, the AD patient's normal appearance conveys a level of sagacity and competence, which is not assumed to be present in children at the same DA.

Also, because of the absence of a physical retrogenesis, the physical capacities of the AD patient may sometimes exceed those of DA-comparable children and infants. For example, a very well cared for GDS stage 7, FAST stage 7b patient who receives retraining can be relatively dexterous as compared to a one-year-old. For example, one GDS 7, FAST-7b patient was able to lace, button, and slip on clothing.

Another consequence of the AD patients' size and strength in comparison with their DA peers is that a grasp reflex in a stage 7 AD patient can be much stronger and more difficult to release than an infant's grasp reflex, with consequences for the management of the AD patient.⁷⁰

Caveat III: AD patients can, to some extent, draw upon previously mastered skills, whereas infants and children may not have access to these skills. Consequently, AD patients may be relatively skilled and "precocious" in comparison with their chronologically younger DA peers. For example, even in late stage 7, long after serviceable speech has been lost, AD patients may occasionally (e.g., during their sleep, or in response

to startle or pain) utter seemingly forgotten words. Infants do not have access to such words.

Caveat IV: AD patients can, to some extent, draw upon previously mastered knowledge, whereas infants and children may not have access to this knowledge. For example, a GDS and FAST 7d AD patient who always had an immaculate household uttered "aagh!" when a caregiver dropped shoes in the middle of the floor. This same 7d AD patient would "whack" a caregiver who put her elbows on the table.

Caveat V: AD patients are older than their DA peers, and old age predisposes to various physical disabilities that influence the life and experience of an AD patient. For example, cataracts predispose AD patients to visual hallucinations.

Caveat VI: AD patients appear to be more prone to rigidity than their DA peers. The causes probably include the AD patients' brain disease in the absence of physical involution and the relative immobility of AD patients. The rigidity can greatly increase disability in the AD patient, ultimately resulting in contractures.

Caveat VII: AD patients can potentially concentrate on a task longer than infants or children at a corresponding DA. Conversely, infants and children are more distractible and impatient than AD patients. For example, one GDS 7, FAST 7c AD patient is known to stare at a newspaper for perhaps an hour. An 11-month-old infant will look at a book for a few minutes.

Caveat VIII: AD patients appear to be less fascinated by the world and less inquisitive than infants and children at a corresponding DA. For example, a two-year-old child may continuously ask questions, such as "What is this?" whereas an AD patient at GDS 6, FAST stage 6c is not inquisitive in this manner.

Care science

Therefore, a care science in AD can be firmly grounded in universal human, retrogenic, and AD-specific principles. The ingredients for the quality care recipe can now be described in considerable detail. These principles can potentially impact positively on the quality of life and excess disability of AD patients at this juncture. Potentially, much of the suffering and distress associated with AD can be relieved.

Pharmacologic treatment of the biomolecular retrogenic process

The nascent understanding of the role of mitogenic factors in the etiopathogenesis of AD has suggested new avenues for fundamental treatment investigations. In many ways, the mitogenic response of biomolecular retrogenesis to AD-associated stressors and toxins is similar to

the mitogenic response to stressors and toxins in non-nervous-system tissues.⁷¹⁻⁷³ In other organ systems and tissues, the stressors and toxins producing a mitogenic response are termed *carcinogens* and the mitogenic response is termed a *neoplastic response*. In the adult brain, despite the occasional occurrence of neurogenesis, an abortive neurogenesis or abortive mitosis is associated with neurofibrillary changes and, perhaps, apoptosis. Consequently, antineoplastic agents that suppress mitosis may have relevance for the treatment of AD.

Conclusion

In summary, retrogenesis appears to represent a new mechanism of disease with distinct biomolecular, pathologic, physiologic, and clinical manifestations. Distinct management and treatment principles and avenues for pharmacologic research follow from these retrogenic mechanisms and their manifestations. AD is the paradigmatic retrogenic disorder, but the process is applicable to a variable extent to many other dementing conditions.

Authors' note

The methodology described herein of a science of management, care, and treatment of Alzheimer's disease and related dementias is the subject of a pending patent application.

Acknowledgments

This work was supported in part by United States Department of Health and Human Services (DHHS) Grants AG03051, AG08051, and AG09127 from the National Institute on Aging of the US National Institutes of Health, by Grants 90AR2160 and 90AM2552 from the US DHHS Administration on Aging, by Grant M01 RR00096 from the General Clinical Research Center Program of the National Center for Research Resources of the US National Institutes of Health, and by the Fisher Center for Alzheimer's Disease Research Foundation.

References

1. Reisberg B, Franssen EH, Hasan SM, et al.: Retrogenesis: Clinical, physiologic and pathologic mechanisms in brain aging, Alzheimer's and other dementing processes. *European Archives of Psychiatry and Clinical Neuroscience*. 1999; 249 (Suppl 3): 28-36.
2. Aristophanes: The Clouds. In Oates WJ, O'Neil E (eds.): *The Complete Greek Drama*. New York: Random House, 1938 [translation]: 595.
3. Shakespeare W: "As you like it," Act 2, Scene 7. In Wells S, Taylor G (eds.): *William Shakespeare: The Complete Works*. (Compact Ed.) Oxford: Oxford University Press, 1988: 638.
4. Rush B: An account of the state of mind and body in old age. In: Rush B (ed.): *Medical Inquiries and Observations*. Philadelphia: Dobson, 1793: 311.
5. de Ajuriaguerra J, Rey M, Bellet-Muller M: A propos de quelques problèmes posés par le déficit opératoire des vieillards atteints de démence dégénérative en début d'évolution. *Cortex*. 1964; 1: 232-256.
6. de Ajuriaguerra J, Tissot R: Some aspects of psychoneurologic disintegration in senile dementia. In Mueller CH, Ciompi L (eds.): *Senile Dementia*. Switzerland: Huber, 1968: 69-79.
7. de Ajuriaguerra J, Tissot R: Some aspects of language in various forms of senile dementia (comparisons with language in childhood). In Lennenberg EH, Lennenberg E (eds.): *Foundations of Language Development* (Vol. 1). New York: Academic Press, 1975: 323-339.
8. Reisberg B, Ferris SH, Anand R, et al.: Functional staging of dementia of the Alzheimer's type. *Annals of the New York Academy of Sciences*. 1984; 435: 481-483.
9. Reisberg B: Functional assessment staging (FAST). *Psychopharmacology Bulletin*. 1988; 24: 653-659.
10. Reisberg B, Ferris SH, Franssen E: Functional degenerative stages in dementia of the Alzheimer's type appear to reverse normal human development. In Shagass C, Josiassen R, Bridger WH, et al. (eds.): *Biological Psychiatry*. 1985 (Vol. 7). New York: Elsevier Science, 1986: 1319-1321.
11. Reisberg B, Ferris SH, Franssen E: An ordinal functional assessment tool for Alzheimer's-type dementia. *Hospital & Community Psychiatry*. 1985; 36: 593-595.
12. Reisberg B: Dementia: A systematic approach to identifying reversible causes. *Geriatrics*. 1986; 41(4): 30-46.
13. Folstein MF, Folstein SE, McHugh PR: Mini-mental state: A practical method for grading the cognitive state of patients for the clinician. *J Psychiatric Research*. 1975; 12: 189-198.
14. Auer SR, Selan SG, Yaffee RA, Reisberg B: The neglected half of Alzheimer disease: Cognitive and functional concomitants of severe dementia. *J American Geriatrics Society*. 1994; 42: 1266-1272.
15. Bobinski M, Wegiel J, Wisniewski HM, et al.: Atrophy of hippocampal formation subdivisions correlates with stage and duration of Alzheimer disease. *Dementia*. 1995; 6: 205-210.
16. Bobinski M, Wegiel J, Tarnawski M, et al.: Relationships between regional neuronal loss and neurofibrillary changes in the hippocampal formation and duration and severity of Alzheimer disease. *J Neuropathology & Experimental Neurology*. 1997; 56: 414-420.
17. Reisberg B, Ferris SH, Franssen E, et al.: Mortality and temporal course of probable Alzheimer's disease: A five-year prospective study. *International Psychogeriatrics*. 1996; 8: 291-311.
18. Reisberg B, Schneck MK, Ferris SH, et al.: The brief cognitive rating scale (BCRS): Findings in primary degenerative dementia (PDD). *Psychopharmacology Bulletin*. 1983; 19: 47-50.
19. Reisberg B, Ferris SH, deLeon MJ, Crook T: The global deterioration scale for assessment of primary degenerative dementia. *American J Psychiatry*. 1982; 139: 1136-1139.
20. Reisberg B, Ferris SH, de Leon MJ: Senile dementia of the Alzheimer type: Diagnostic and differential diagnostic features with special reference to functional assessment staging. In Traber J, Gispén WH (eds.): *Senile Dementia of the Alzheimer Type* (Vol. 2). Berlin: Springer-Verlag, 1985: 18-37.
21. Reisberg B, Patschull-Furlan A, Franssen E, et al.: Cognition-related functional, praxis and feeding changes in CNS-aging and Alzheimer's disease and their developmental analogies. In Beyreuther KG, Schettler G (eds.): *Molecular Mechanisms of Aging*. Berlin: Springer-Verlag, 1990: 18-40.
22. Ouvrier RA, Goldsmith RF, Ouvrier S, Williams IC: The value of the mini-mental state examination in childhood: A preliminary study. *J Child Neurology*. 1993; 8: 145-148.
23. Selan SG, Foster JR, Reisberg B, et al.: Application of Piagetian measures of cognition in severe Alzheimer's disease. *Psychiatric J University of Ottawa*. 1990; 15: 221-226.
24. Thornbury MM: Cognitive performance on Piagetian tasks by Alzheimer's disease patients. *Research in Nursing & Health*. 1992; 15: 11-18.
25. Reisberg B, Ferris S: A clinical rating scale for symptoms of psychosis in Alzheimer's disease. *Psychopharmacology Bulletin*. 1985; 21: 101-104.

26. Reisberg B, Borenstein J, Salob SP, *et al.*: Behavioral symptoms in Alzheimer's disease: Phenomenology and treatment. *J Clinical Psychiatry*. 1987; 48 (Suppl): 9-15.
27. Reisberg B, Franssen E, Sclan SG, *et al.*: Stage specific incidence of potentially remediable behavioral symptoms in aging and Alzheimer's disease: A study of 120 patients using the BEHAVE-AD. *Bulletin of Clinical Neurosciences*. 1989; 54: 95-112.
28. Reisberg B, Auer SR, Monteiro I, *et al.*: A rational psychological approach to the treatment of behavioral disturbances and symptomatology in Alzheimer's disease based upon recognition of the developmental age. *International Academy for Biomedical and Drug Research*. 1998; 13: 102-109.
29. Reisberg B, Kenowsky S, Franssen EH, *et al.*: President's Report: Towards a science of Alzheimer's disease management: A model based upon current knowledge of retrogenesis. *International Psychogeriatrics*. 1999; 11: 7-23.
30. Paulson G, Gottlieb G: Developmental reflexes: The reappearance of fetal and neonatal reflexes in aged patients. *Brain*. 1968; 91: 37-52.
31. Jacobs L, Gossman MD: Three primitive reflexes in normal adults. *Neurology*. 1980; 30: 184-188.
32. Basavaraju NG, Silberstone F, Libow L, Paraskeros K: Primitive reflexes and perceptual sensory tests in the elderly—their usefulness in dementia. *J Chronic Disease*. 1981; 34: 367-377.
33. Huff FJ, Boller F, Luchelli F, *et al.*: The neurologic examination in patient's with probable Alzheimer's disease. *Archives of Neurology*. 1987; 44: 929-932.
34. Franssen EH, Reisberg B, Kluger A, *et al.*: Cognition independent neurologic symptoms in normal aging and probable Alzheimer's disease. *Archives of Neurology*. 1991; 48: 148-154.
35. Franssen EH, Souren LEM, Torossian CL, Reisberg B: The neurologic syndrome of severe Alzheimer's disease: Relationship to functional decline. *Archives of Neurology*. 1993; 50: 1029-1039.
36. Franssen EH, Souren LEM, Torossian CL, Reisberg B: Utility of developmental reflexes in the differential diagnosis and prognosis of incontinence in Alzheimer's disease. *J Geriatric Psychiatry & Neurology*. 1997; 10: 22-28.
37. Brun A, Gustafson L: Distribution of cerebral degeneration in Alzheimer's disease. *Archiv für Psychiatrie und Nervenkrankheiten*. 1976; 223: 15-33.
38. McGeer PL, McGeer EG, Akiyama H, *et al.*: Neuronal degeneration and memory loss in Alzheimer's disease and aging. *Experimental Brain Research*. 1990; (Supp. 21): 411-426.
39. Braak H, Braak E: Development of Alzheimer-related neurofibrillary changes in the neocortex inversely recapitulates cortical myelogenesis. *Acta Neuropathologica*. 1996; 92: 197-201.
40. Raz N: Aging of the brain and its impacts on cognitive performance: Integration of structural and functional findings. In Craik FIM, Salthouse TA (eds.): *Handbook of Aging & Cognition—II*. Mahwah, NJ: Erlbaum, 2000: 1-90.
41. Wu Q, Combs C, Cannady SB, Geldmacher DS, Herrup K: Beta-amyloid activated microglia induce cell cycling and cell death in cultured cortical neurons. *Neurobiology of Aging*. 2000; 21: 797-806.
42. Lee M-S, Kwon YT, Li M, *et al.*: Neurotoxicity induces cleavage of p35 to p25 by calpain. *Nature*. 2000; 405: 360-364.
43. Patrick GN, Zukerberg L, Nikolic M, *et al.*: Conversion of p35 to p25 deregulates Cdk5 activity and promotes neurodegeneration. *Nature*. 1999; 402: 615-622.
44. Nagy Z, Esiri MM, Smith AD: The cell division cycle and the pathophysiology of Alzheimer's disease. *Neuroscience*. 1998; 87: 731-739.
45. Nagy Z: Cell cycle regulatory failure in neurones causes and consequences. *Neurobiology of Aging*. 2000; 21: 761-769.
46. Busser J, Geldmacher DS, Herrup K: Ectopic cell cycle proteins predict the sites of neuronal cell death in Alzheimer's disease brain. *J Neuroscience*. 1998; 18: 2801-2807.
47. Arendt T, Holzer M, Ströbe A, *et al.*: Activated mitogenic signaling induces a process of dedifferentiation in Alzheimer's disease that eventually results in cell death. *Annals of the New York Academy of Sciences*. 2000; 920: 249-255.
48. Arendt T: Alzheimer's disease as a loss of differentiation control in a subset of neurons that retain immature features in the adult brain. *Neurobiology of Aging*. 2000; 21: 783-796.
49. Arendt T: Alzheimer's disease as a disorder of mechanisms underlying structural brain self-organization. *Neuroscience*. 2001; 102: 723-765.
50. Smith MZ, Nagy Z, Esiri MM: Cell cycle-related protein expression in vascular dementia and Alzheimer's disease. *Neuroscience Letters*. 1999; 271: 45-48.
51. Vincent I, Jicha G, Rosado M, Dickson DW: Aberrant expression of mitogenic Cdc2/cyclin B1 kinase in degenerating neurons of Alzheimer's disease brain. *J Neuroscience*. 1997; 17: 3588-3598.
52. Vincent I: Cycling to the finish. *Neurobiology of Aging*. 2000; 21: 757-760.
53. Husseman JW, Nochlin D, Vincent I: Mitotic activation: A convergent mechanism for a cohort of neurodegenerative diseases. *Neurobiology of Aging*. 2000; 21: 815-828.
54. Atkins CM, Selcher JC, Petraitis JJ, *et al.*: The MAPK cascade is required for mammalian associative learning. *Nature Neuroscience*. 1998; 1: 602-609.
55. Blum S, Moore AN, Adams F, Dash PK: A mitogen-activated protein kinase cascade in the CA1/CA2 subfield of the dorsal hippocampus is essential for long-term spatial memory. *J Neuroscience*. 1999; 19: 3535-3544.
56. Schafe GE, Nadel NV, Sullivan GM, *et al.*: Memory consolidation for contextual and auditory fear conditioning is dependent on protein synthesis, PKA, and MAP kinase. *Learning & Memory*. 1999; 6: 97-110.
57. Selcher JC, Atkins CM, Trzaskos JM, *et al.*: A necessity for MAP kinase activation in mammalian spatial learning. *Learning & Memory*. 1999; 6: 478-490.
58. Kobayashi S, Ishiguro K, Omori A, *et al.*: A cdc2-related kinase PSSALRE/edk5 is homologous with the 30kDa subunit of tau protein kinase II, a proline-directed protein kinase associated with microtubule. *Federation of European Biochemical Societies Letters*. 1993; 335: 171-175.
59. Ledesma MD, Correas I, Avila J, Diaznido J: Implication of brain Cdc2 and Map2 kinases in the phosphorylation of tau protein in Alzheimer's disease. *Federation of European Biochemical Societies Letters*. 1992; 308: 218-224.
60. Suzuki T, Oishi M, Marshak DR, *et al.*: Cell cycle-dependent regulation of the phosphorylation and metabolism of the Alzheimer amyloid precursor protein. *EMBO J*. 1994; 13: 1114-1122.
61. Clark R, Smith D, Jobst KA, *et al.*: Folate, vitamin B12, and serum total homocysteine levels in confirmed Alzheimer disease. *Archives of Neurology*. 1998; 55: 1449-1455.
62. Kristensen MO, Gulmann NC, Christensen JE, *et al.*: Serum cobalamin and methylmalonic acid in Alzheimer dementia. *Acta Neurologica Scandinavica*. 1993; 87: 475-481.
63. Diaz-Arrastia R: Hyperhomocysteinemia: A new risk factor for Alzheimer disease? [Editorial]. *Archives of Neurology*. 1998; 55: 1407-1408.
64. Hofman A, Ott A, Breteler MMB, *et al.*: Atherosclerosis, apolipoprotein E, and prevalence of dementia and Alzheimer's disease in the Rotterdam Study. *Lancet*. 1997; 349: 151-154.
65. Snowden DA, Greiner LH, Mortimer JA, *et al.*: Brain infarction and the clinical expression of Alzheimer's disease: The nun study. *JAMA*. 1997; 277: 813-817.

66. Esiri MM, Nagy Z, Smith MZ, *et al.*: Cerebrovascular disease and threshold for dementia in the early stages of Alzheimer's disease. *Lancet*. 1999; 354: 919-920.
67. Reisberg B, Franssen E, Shah MA, *et al.*: Clinical diagnosis of dementia. In Maj M, Sartorius N (eds.): *World Psychiatric Association Series. Evidence & Experience in Psychiatry: Dementia* (Vol. 3). Chichester, UK: John Wiley and Sons. 2000: 69-115.
68. Souren LEM, Franssen EM, Reisberg B: Contractures and loss of function in patients with Alzheimer's disease. *J American Geriatrics Society*. 1995; 43: 650-655.
69. Blessed G, Tomlinson BE, Roth M: The association between quantitative measures of dementia and senile change in the cerebral gray matter of elderly subjects. *British J Psychiatry*. 1968; 114: 797-811.

70. Souren LEM, Franssen EH, Reisberg B: Neuromotor changes in Alzheimer's disease: Implications for patient care. *J Geriatric Psychiatry & Neurology*. 1997; 10: 93-98.
71. Raina AK, Zhu X, Rottkamp CA, *et al.*: Cyclin toward dementia: Cell cycle abnormalities and abortive oncogenesis in Alzheimer disease. *J Neuroscience Research*. 2000; 61: 128-133.
72. Zhu X, Raina AK, Boux H, *et al.*: Activation of oncogenic pathways in degenerating neurons in Alzheimer disease. *International J Developmental Neuroscience*. 2000; 18: 433-437.
73. Raina AK, Zhu X, Monteiro M, *et al.*: Abortive oncogeny and cell cycle-mediated events in Alzheimer disease. In Meijer L, Jezequel A, Ducommun B (eds.): *Progress in Cell Cycle Research*. New York: Kluwer Academic/Plenum Publishers, 2000: 235-242.

Calendar

**National Conference of Gerontological
Nurse Practitioners
Annual Meeting
September 18-21, 2002
Chicago, Illinois**

For registration and information, contact:
National Conference
of Gerontological Nurse Practitioners
P.O. Box 232230
Centreville, VA 20120-2230
Tel: (703) 802-0088
Fax: (703) 802-1436
Web site: ncgnp@ncgnp.org

**1st International Conference
on Interdisciplinary Gerontology
September 19-21, 2002
Bad Ischl, Austria**

For registration and information, contact:
Verein M.A.S.
c/o Institut für
Interdisziplinäre Gerontologie
Wiesingerstrasse 4
4820 Bad Ischl
AUSTRIA
Tel./Fax: +43-6132-21 410
Email: congress@mas.or.at

**2nd International Conference
on Metals and the Brain
September 25-28, 2002
Fez, Morocco**

For registration and information, contact:
King's Holidays
Conference Services Office
Rue Ahmed Chbihi, Immb 3
Appt 4, Avenue des FAR
VN Fcs MAROC
Tel.: 00-212-55-93-04-99
Fax: 00-212-55-93-34-55
Email: kingcongres2002@iam.net.ma

**18th International Conference
on Alzheimer's Disease
October 23-26, 2002
Barcelona, Spain**

For registration and information, contact:
Information and registration
Suport Servicos
Calvet, 30
08021 Barcelona, Spain
Tel.: +34-93-201-7571
Fax: +34-93-201-9789
Email: suport@suportserveis.com
Web site: <http://www.alzheimer2002.com/eng/>

ORIGINAL CONTRIBUTION

Effects of Rofecoxib or Naproxen vs Placebo on Alzheimer Disease Progression

A Randomized Controlled Trial

Paul S. Aisen, MD

Kimberly A. Schafer, MS

Michael Grundman, MD, MPH

Eric Pfeiffer, MD

Mary Sano, PhD

Kenneth L. Davis, MD

Martin R. Farlow, MD

Shelia Jin, MPH

Ronald G. Thomas, PhD

Leon J. Thal, MD

for the Alzheimer's Disease Cooperative Study

ALZHEIMER DISEASE (AD) IS among the most important health problems of elderly persons, affecting more than 4 million people in the United States. In the last decade, cholinesterase inhibitors have been widely used to alleviate symptoms of cognitive dysfunction in AD. The use of anti-inflammatory drugs is among the strategies under active investigation for the development of effective disease-modifying treatment for AD. This approach is supported by a wealth of laboratory evidence that inflammatory mechanisms contribute to neuronal damage in AD.¹ Furthermore, many epidemiological studies suggest that anti-inflammatory drugs have a protective effect, reducing the incidence of AD.²

Results of small pilot clinical trials of nonsteroidal anti-inflammatory drug

Context Laboratory evidence that inflammatory mechanisms contribute to neuronal injury in Alzheimer disease (AD), along with epidemiological evidence, suggests that nonsteroidal anti-inflammatory drugs (NSAIDs) may favorably influence the course of the disease.

Objective To determine whether treatment with a selective cyclooxygenase (COX)-2 inhibitor (rofecoxib) or a traditional nonselective NSAID (naproxen) slows cognitive decline in patients with mild-to-moderate AD.

Design Multicenter, randomized, double-blind, placebo-controlled, parallel group trial, with 1-year exposure to study medications.

Setting Forty ambulatory treatment centers affiliated with the Alzheimer's Disease Cooperative Study consortium.

Participants Participants with mild-to-moderate AD (Mini-Mental State Examination score of 13-26) were recruited from December 1999 to November 2000 using clinic populations, referrals from community physicians, and local advertising. Stable use of cholinesterase inhibitors, estrogen, low-dose aspirin, and vitamin E was allowed. Participants with inflammatory diseases that might respond to the study medications were excluded. Of 474 participants screened, 351 were enrolled.

Interventions Once-daily rofecoxib, 25 mg, or twice-daily naproxen sodium, 220 mg, or placebo.

Main Outcome Measures The primary outcome measure was the 1-year change in the Alzheimer Disease Assessment Scale-Cognitive (ADAS-Cog) subscale score. Secondary outcome measures included the Clinical Dementia Rating scale sum-of-boxes, the Neuropsychiatric Inventory, the Quality of Life-AD, and the time to attainment of significant end points (4-point decline from baseline ADAS-Cog score, 1-step worsening on the global Clinical Dementia Rating scale, 15-point decline on the ADCS activities of daily living inventory, institutionalization, or death).

Results The 1-year mean (SD) change in ADAS-Cog scores in participants treated with naproxen (5.8 [8.0]) or rofecoxib (7.6 [7.7]) was not significantly different from the change in participants treated with placebo (5.7 [8.2]). Results of secondary analyses showed no consistent benefit of either treatment. Fatigue, dizziness, and hypertension were more commonly reported in the active drug groups, and more serious adverse events were found in the active treatment group than in the placebo group.

Conclusion The results of this study indicate that rofecoxib or low-dose naproxen does not slow cognitive decline in patients with mild-to-moderate AD.

JAMA. 2003;289:2819-2826

www.jama.com

For editorial comment see p 2865.

Author Affiliations and Participating members of this Alzheimer's Disease Cooperative Study are listed at the end of this article.
Corresponding Author and Reprints: Paul S. Aisen,

MD, Department of Neurology, Georgetown University Medical Center, Bldg D, Suite 207, 4000 Reservoir Rd, NW, Washington, DC 20057 (e-mail: psa@georgetown.edu).

JAMA, June 4, 2003—Vol 289, No. 21 2819

EFFECTS OF ROFECOXIB OR NAPROXEN VS PLACEBO ON ALZHEIMER DISEASE

(NSAID) efficacy in the treatment of AD have been encouraging. One report indicated a stabilizing effect of indomethacin treatment for 6 months,³ while a follow-up study showed a trend toward benefit with diclofenac treatment for 25 weeks.⁴ Neither study showed conclusive results, and the patients with AD showed poor tolerability of both drugs. Larger studies with better tolerated treatment regimens, such as low-dose prednisone for 1 year⁵ and hydroxychloroquine for 18 months,⁶ failed to demonstrate benefit in the treatment of AD.

Recent studies have provided strong support for the testing of efficacy of NSAIDs in the prevention of AD. A large epidemiologic study using computerized medical records to accurately indicate drug use corroborates earlier work, with evidence that use of NSAIDs sharply reduces AD risk.⁷ In transgenic mice with AD-type amyloid deposition, the NSAID ibuprofen reduced brain inflammation and levels of soluble and insoluble amyloid peptide, and improved behavior.^{8,9}

In this trial, we tested the hypothesis that NSAID treatment would slow the rate of cognitive decline in participants with mild-to-moderate AD. Selection of treatment regimens was based on consideration of the appropriate target in AD brain, as well as medication tolerability. Results of studies in cell culture systems,^{10,11} rodents,^{12,13} and human brain^{10,14} indicate that neuronal up-regulation of cyclooxygenase (COX)-2 may contribute to neurodegeneration in AD. However, the epidemiological data support the efficacy of long-term use of low doses of nonselective NSAIDs that inhibit COX-1 and COX-2,⁴ and microglial COX-1 may be an important target.¹⁵ Long-term treatment with full doses of nonselective NSAIDs has been associated with a substantial risk of serious gastrointestinal tract toxicity. Therefore, we used a standard dose of the selective COX-2 inhibitor rofecoxib and a low-dose of the nonselective NSAID naproxen. We chose a treatment period of 1 year based on evidence from a pilot study of NSAID efficacy with short-term treatment³ and the con-

cern about the risk of long-term exposure in individuals with AD.

METHODS

Study Design

The study used a randomized, double-blind, 3-group parallel design to compare rofecoxib or naproxen with placebo. The treatment period was 12 months, followed by a 2-month washout. Forty ambulatory treatment centers affiliated with the Alzheimer's Disease Cooperative Study (ADCS) consortium participated in this trial after obtaining approval from their local institutional review boards. Written informed consent was obtained from participants and/or legally authorized representatives, according to local guidelines.

Individuals with probable AD¹⁶ recruited from December 1999 to November 2000 using clinic populations, referrals from community physicians, and local advertising were eligible if they did not have comorbid conditions that increased the risk of adverse events associated with NSAID treatment (hypersensitivity to aspirin or NSAIDs, active peptic ulcer disease within 5 years, renal insufficiency [serum creatinine level >1.5 mg/dL or >132.6 μ mol/L], clinically significant liver disease, poorly controlled hypertension, congestive heart failure, or bleeding disorder). Individuals were excluded if they had comorbid conditions that might respond to NSAIDs (eg, inflammatory arthritis). Individuals also were excluded if within the prior 2 months they had regularly used anti-inflammatory medications (aspirin at a daily dose ≤ 325 mg was allowed), neuroleptics, antidepressants, sedatives, anti-Parkinsonian medications, or any investigational treatment for AD. Inclusion criteria included age older than 50 years and Mini-Mental State Examination (MMSE)¹⁷ score within the range of 13 to 26. Stable use of cholinesterase inhibitors was allowed.

Rofecoxib tablets, 25 mg, were over-encapsulated to allow preparation of an identical placebo capsule. Naproxen sodium tablets, 220 mg, and identical pla-

cebo tablets were supplied by Bayer Consumer, Inc (Morristown, NJ). Each participant received 2 bottles of study medication with coded labels at baseline and at the 3-, 6-, and 9-month visits: 1 bottle containing rofecoxib or identical placebo capsules, to be taken once daily; and 1 bottle of naproxen or identical placebo capsules, to be taken twice daily.

The randomization process used a permuted block design with block size of 3 stratified by site. The randomization sequence was generated by the ADCS data center. Scratch-off codebreakers were used so that instances of unblinding would be documented; all codebreakers were collected at the end of the trial. Adequacy of masking was assessed by questionnaires completed by participants, caregivers, psychometrists, and site investigators.

Safety assessments, including vital signs, physical examination, urinalysis, and hematology and chemistry blood tests, were performed at each visit. Cognitive and behavioral assessments were performed at baseline and at months 1, 3, 6, 9, 12, and 14.

Outcome Measures

The primary outcome measure for this trial was the 1-year change score on the Alzheimer Disease Assessment Scale-Cognitive (ADAS-Cog) subscale,¹⁸ an instrument that evaluates memory, attention, reasoning, language, orientation, and praxis. We considered a significant benefit in the active group to be a 50% reduction in cognitive decline as indicated by change in ADAS-Cog score compared with the placebo group. Data from a small pilot study³ have supported the hypothesis that an effect of this magnitude might be seen, and we considered that a smaller effect might not justify the risk of long-term NSAID therapy in the AD participants. Using longitudinal ADAS-Cog data from an earlier multicenter trial,⁵ we estimated that 100 participants were required in each treatment group (naproxen, rofecoxib, and placebo) for a power of 0.8. An overall drop-out rate of 20% was anticipated. Since we ex-

EFFECTS OF ROFECOXIB OR NAPROXEN VS PLACEBO ON ALZHEIMER DISEASE

pected more drop-outs from adverse drug effects in the active drug groups, fewer participants were randomly assigned to the placebo group (participants were randomly assigned using a ratio of 1.2 [naproxen]:1.2 [rofecoxib]: 1.1 [placebo]).


Secondary outcome measures included the Clinical Dementia Rating sum-of-boxes (CDR-SOB),¹⁹ Alzheimer's Disease Cooperative Study activities of daily living (ADCS-ADL) scale,²⁰ the Neuropsychiatric Inventory (NPI),²¹ the Quality of Life-AD (QOL-AD),²² and the time to attainment of significant end points (4-point decline from baseline ADAS-Cog score, 1-step worsening on the global CDR scale, 15-point decline on the ADCS-ADL, institutionalization, or death).

Statistical Analysis

The primary analysis was a comparison of the change in ADAS-Cog score between the naproxen and rofecoxib groups and the placebo group using analysis of covariance (ANCOVA), with the baseline ADAS-Cog score as a covariate. The statistical plan included assessment of the treatment groups for significant imbalance in age, sex, or apolipoprotein E genotype that might influence outcome; if an imbalance was present ($P < .15$) and the factor influenced ADAS-Cog change scores ($P < .10$), it would be included in the ANCOVA model.

The primary analysis was conducted on an intent-to-treat basis (ie, including all randomized participants). Because AD is a disease of progressive cognitive deterioration, the protocol specified an alternative imputation scheme rather than last-observation-carried-forward (LOCF) analysis. (LOCF analysis would overestimate a treatment effect if an excess of drop-outs occurred in the active drug group due to adverse effects.) To impute a participant's missing score at week 52, an estimate of change in ADAS-Cog score during the unobserved period based on all individuals with complete data from the participant's treatment group was

Figure. Patient Flow



applied to the participant's last observed score. A linear rate of progression was not assumed. We also conducted secondary analyses of change in ADAS-Cog scores using the LOCF imputation, and an analysis of completers only (ie, those who completed the week 52 visit). In addition, we conducted a longitudinal regression analysis using the method of generalized estimating equations.²³ For this analysis, a pair-wise comparison of the naproxen group vs the placebo group and the rofecoxib group vs the placebo group, the outcome was the ADAS-Cog score clustered by the baseline, 3-, 6-, 9-, and 12-month visits.

A planned secondary analysis of time to reach clinically significant end points was performed using a Cox proportional hazards model. Intent-to-treat analysis of secondary measures were conducted using both the slope imputation strategy described above and LOCF for missing week 52 values. No interim analysis was performed. Statistical analyses were performed using SAS version 8.2 (SAS Institute, Cary, NC).

RESULTS

The flow of participants through the study protocol is shown in FIGURE 1. From a total of 474 participants

screened, 351 met the study criteria and were randomly assigned to 1 of the 3 treatment groups (rofecoxib, naproxen, or placebo). Considering the likelihood that more participants in the active groups would drop out due to adverse effects, more participants were randomly assigned to rofecoxib ($n = 122$) and naproxen ($n = 118$) than to the placebo group ($n = 111$).

The primary outcome measure (change in ADAS-Cog score at week 52) was obtained for study completers: 90 (76%) participants in the naproxen group, 89 (73%) in the rofecoxib group, and 88 (79%) in the placebo group. The predominant reasons for early study discontinuation were caregiver issues (rofecoxib, $n = 12$; naproxen, $n = 11$; and placebo, $n = 10$) and adverse events (rofecoxib, $n = 14$; naproxen, $n = 11$; and placebo, $n = 9$).

The demographic and clinical characteristics of the treatment groups at baseline are shown in TABLE 1. No significant differences were found between the groups on any of the demographic or baseline characteristics. Eighty-seven participants (25%) were using low-dose aspirin during the trial: 33 (27%) in the rofecoxib group, 30 (25%) in the naproxen group, and 24 (22%) in the placebo group. Aspirin use

EFFECTS OF ROFECOXIB OR NAPROXEN VS PLACEBO ON ALZHEIMER DISEASE

Table 1. Participant Demographics and Assessment Scores at Baseline

	Placebo (n = 111)	Naproxen (n = 118)	P Value*	Rofecoxib (n = 122)	P Value*
Age, mean (SD), y	73.8 (8.0)	74.1 (7.8)	.78	73.7 (7.2)	.96
Sex, female, %	55.9	48.3	.31	54.9	.99
Education, mean (SD), y	14.3 (3.3)	13.8 (3.2)	.37	13.8 (3.2)	.24
Duration of disease, mean (SD), y	4.2 (3.3)	4.1 (2.3)	.92	4.1 (2.3)	.89
Users of cholinesterase inhibitors at baseline, No. (%)	74 (67)	78 (66)	.99	88 (70)	.57
Participants starting cholinesterase inhibitors during the study, No. (%)	6 (5)	10 (8)	.44	8 (7)	.79
Users of low-dose aspirin, No. (%)	24 (22)	30 (25)	.54	33 (27)	.36
Apolipoprotein E, No. (%) with ≥ 1 E4 allele	67.6	70.4	.68	68.1	.95
MMSE score, mean (SD)†	20.8 (3.6)	20.7 (3.6)	.81	21.2 (3.8)	.87
ADAS-Cog score, mean (SD)†	24.2 (9.4)	24.4 (10.2)	.92	23.4 (9.4)	.45
CDR sum-of-boxes score, mean (SD)†	5.5 (2.5)	6.0 (2.9)	.28	5.6 (2.5)	.82

Abbreviations: ADAS-Cog, Alzheimer Disease Assessment Scale-Cognitive subscale; CDR, Clinical Dementia Rating; MMSE, Mini-Mental State Examination.

*P values for comparison with placebo, using the Wilcoxon test for continuous variables and the Fisher exact test for frequencies.

†The range of scores for MMSE is 0 to 30 points; for ADAS-Cog, 0 to 70 points; and CDR sum-of-boxes, 0 to 18 points.

did not show a significant effect on ADAS-Cog scores for all groups or for the individual treatment groups.

Baseline characteristics were compared between participants who discontinued early and those who completed the study. In the naproxen group, participants who discontinued early had lower mean (SD) MMSE scores (19.6 [3.3] vs 21.0 [3.6]; $P = .04$; using Wilcoxon rank sum test) and higher mean (SD) CDR-SOB scores (7.2 [3.4] vs 5.7 [2.7]; $P = .03$). No significant differences were found between the rofecoxib and placebo groups.

A total of 238 (68%) participants were taking cholinesterase inhibitors at the time of enrollment. Twenty-four participants (rofecoxib group, $n = 8$; naproxen group, $n = 10$; and placebo group, $n = 6$) started treatment with cholinesterase inhibitors prior to the month 12 visit. These participants were included in the primary intent-to-treat analyses. In a secondary analysis of the main outcome measure excluding these 24 subjects, the results were similar (data not shown).

Analysis of Outcomes

Primary Outcome Measure. The effect of treatment on the primary and secondary measures in the intent-to-treat analysis is shown in TABLE 2. None of

the planned covariates (age, sex, and apolipoprotein E genotype) was included in the analysis, because there was neither imbalance nor association with the outcome.

Neither active treatment had a beneficial effect on the primary measure, the change in mean ADAS-Cog scores; the group treated with rofecoxib showed a trend toward greater cognitive decline compared with the other groups. For the comparison of naproxen with placebo, the SE of the difference between change scores was 1.07 (95% confidence interval for the difference in the change scores, -0.1 ± 2.14); therefore, it is highly unlikely that naproxen treatment reduces the 1-year decline in ADAS-Cog score by more than 2.04/5.7, or 36%. For the comparison of rofecoxib with placebo, the SE of the difference between change scores was 1.03; a similar calculation indicates it is highly unlikely that rofecoxib reduces the decline in ADAS-Cog score by more than 15%.

When LOCF imputation of the change in ADAS-Cog score was substituted for the primary imputation scheme, the result was similar to the primary analysis (mean [SD] ADAS-Cog change in score in the placebo group, 4.9 [8.2]; rofecoxib group, 5.9 [7.7], $P = .52$; and naproxen group, 5.0 [7.9],

$P = .94$). Analysis of participants taking study medication who completed the 52-week visit yielded similar results, with no significant difference in rate of decline for the mean (SD) ADAS-Cog score (placebo, 5.3 [8.5], $n = 79$; rofecoxib, 6.8 [7.7], $n = 78$, $P = .47$; naproxen, 5.3 [7.9], $n = 80$, $P = .9$). Regression analysis using the generalized estimating equations method demonstrated no effect of rofecoxib ($P = .77$) or naproxen ($P = .64$) on longitudinal ADAS-Cog scores.

Secondary Outcome Measures. The intent-to-treat analysis using the primary imputation scheme for missing values revealed no significant difference in decline for the CDR-SOB scores across all the groups (Table 2). Other analyses, using LOCF imputation or assessing completers only, showed similar results (data not shown). The ADCS-ADL scores showed a trend toward a beneficial effect with use of rofecoxib. However, none of the secondary analyses were adjusted for multiple comparisons because this was not specified in the original protocol. In the slope imputation analysis, use of rofecoxib reduced the decline in ADCS-ADL score by 22% (unadjusted $P = .09$); in the LOCF analysis, use of rofecoxib reduced the decline in ADCS-ADL score by 33% (unadjusted $P = .04$). In an analysis of study completers only, rofecoxib reduced the decline in ADCS-ADL score by 27% (unadjusted $P = .09$). Treatment with rofecoxib or naproxen had no effect on the NPI or QOL-AD scores.

The planned survival analysis considered the time interval from the baseline visit to the first among 5 possible end points: death, institutionalization, increase in global CDR score, 15-point decline on the ADCS-ADL scale, or 4-point decline on the ADAS-Cog scale. A total of 242 subjects (69%) reached at least 1 end point (placebo group, $n = 78$ [70%]; naproxen group, $n = 83$ [70%]; and rofecoxib, $n = 81$ [66%]). Rofecoxib ($P = .63$) or naproxen ($P = .50$) vs placebo did not differ in time to first end point. When time to individual end

EFFECTS OF ROFECOXIB OR NAPROXEN VS PLACEBO ON ALZHEIMER DISEASE

Table 2. One-Year Change in Assessment Scores in the Intent-to-Treat Analysis*

	Placebo (n = 111), Mean (SD)			Naproxen (n = 118), Mean (SD)			P Value†	Rofecoxib (n = 122), Mean (SD)			P Value‡
	Baseline	1 Year	Change	Baseline	1 Year	Change		Baseline	1 Year	Change	
ADAS-Cog	24.2 (9.6)	29.9 (13.7)	-5.7 (8.2)	24.4 (10.2)	30.2 (13.9)	-5.8 (8.0)	.96	23.4 (9.4)	31.0 (13.2)	-7.6 (7.7)	.09‡
CDR-SOB	5.5 (2.5)	7.7 (3.9)	-2.2 (2.3)	6.0 (2.9)	8.3 (4.0)	-2.3 (2.3)	.89	5.6 (2.5)	7.8 (3.5)	-2.2 (2.4)	.89
ADCS-ADL	82.8 (11.4)	51.3 (16.3)	-11.5 (11.2)	60.0 (13.1)	51.3 (16.3)	-8.7 (10.5)	.14	82.2 (11.4)	53.9 (14.4)	-8.3 (9.7)	.08
NPI	8.7 (10.6)	12.2 (12.8)	-3.4 (11.9)	9.4 (9.6)	13.1 (14.4)	-3.7 (12.5)	.76	9.2 (9.6)	13.1 (14.2)	-3.8 (10.9)	.76
QOL-AD	36.7 (5.8)	34.3 (6.0)	-2.4 (5.1)	37.0 (5.9)	34.5 (6.2)	-2.6 (5.5)	.93	36.7 (5.9)	33.8 (6.2)	-2.9 (5.6)	.85

Abbreviations: ADAS-Cog, Alzheimer Disease Assessment Scale-Cognitive subscale; ADCS-ADL, Alzheimer Disease Cooperative Study-Activities of Daily Living; CDR-SOB, Clinical Dementia Rating sum-of-boxes; NPI, Neuropsychiatric Inventory; QOL-AD, Quality of Life, Alzheimer Disease.

*For each scale, negative change scores indicate deterioration and positive change scores represent improvement. The range of points for ADAS-Cog is 0 to 70; CDR-SOB is 0 to 18; ADCS-ADL is 0 to 78; NPI is 0 to 144; and QOL-AD is 13 to 52.

†P values for comparison with placebo.

‡Holm adjustment applied to the P value for the ADAS-Cog. The purpose of the Holm adjustment is to account for planned multiple comparisons in the primary end point analysis, thus ensuring an overall type I error $\leq .05$; if either comparison yields $P \leq .05$, the smaller P value is doubled. The unadjusted P value was .044 for the rofecoxib-placebo comparison for ADAS-Cog.

points was examined, the only significant difference was between the rofecoxib and placebo groups for time to institutionalization, favoring the rofecoxib group ($P = .03$) (data not shown). However, this analysis was based on a small number of events (7 in the placebo group and 1 in the rofecoxib group), and the P value was not adjusted for multiple comparisons.

Adverse Events

Treatment emergent adverse events were grouped into categories for analysis. Naproxen and rofecoxib groups had more frequent adverse event reports ($P < .10$) (TABLE 3). Dizziness and fatigue were more common in the active drug groups than in the placebo group, and dry mouth was more common in the naproxen group than the placebo group. New cases of hypertension were reported as adverse events in the active drug groups, but none were reported in the placebo group (naproxen [$n = 6$] vs placebo [$n = 0$], $P = .03$ and rofecoxib [$n = 9$] vs placebo, $P = .004$).

In addition to adverse events reported on case report forms, study participants were asked about a list of symptoms that might arise with NSAID treatment. When the proportion of each treatment group reporting new symptoms (not present at baseline) was compared, the difference was significant compared with placebo for dry mouth (10% for naproxen vs 2% for placebo; $P = .01$) and muscle pain (13% for rofecoxib vs 4% for placebo; $P = .03$). The percentage of participants

Table 3. Adverse Events Reported More Frequently by Participants in the Active Treatment Groups

	Placebo, No. (%)	Naproxen, No. (%)	P Value*	Rofecoxib, No. (%)	P Value†
Fatigue	7 (6)	17 (14)	.08	18 (15)	.05
Dizziness	5 (5)	14 (12)	.06	14 (11)	.06
Dry mouth	1 (1)	6 (5)	.06	1 (1)	>.99
Hypertension	0	6 (5)	.03	9 (7)	.004

*For comparison of naproxen vs placebo.

†For comparison of rofecoxib vs placebo.

with new reports of any gastrointestinal tract symptom did not differ among the treatment groups (37% for naproxen, 31% for rofecoxib, and 30% for placebo).

Despite the cases of hypertension noted above, mean blood pressure did not differ significantly by treatment group. The effects of treatment group on mean (SD) change were minimal for hematocrit (placebo [$n = 87$]: -0.3% [2.2]; naproxen [$n = 88$]: -1.0% [2.2] vs placebo, $P = .01$; and rofecoxib [$n = 87$]: -1.2% [2.2] vs placebo; $P = .05$) and for serum creatinine (0.02 [0.13] mg/dL or 1.8 [11.5] $\mu\text{mol/L}$, $n = 88$; naproxen [$n = 87$]: 0.02 [0.13] mg/dL or 1.8 [11.5] $\mu\text{mol/L}$, vs placebo, $P = .90$; and rofecoxib [$n = 84$]: 0.08 [0.16] mg/dL or 7.1 [14.1] $\mu\text{mol/L}$ vs placebo, $P = .01$).

The total number of serious adverse events in each treatment group, along with number of events within specific categories, did not differ significantly (TABLE 4).

Success of Blinding

The randomization code was broken in 1 instance, based on clinical need for

the management of an acute medical problem. Study personnel involved in administering assessment tools were shielded from group assignment data in this instance.

The results of questionnaires administered at the month 12 visit indicated that the percentage of participants who believed they were taking active study medication also did not differ significantly across the treatment groups (rofecoxib group, 70%; naproxen group, 74%; and placebo group, 73%; $P = .82$, Fisher exact test), indicating that blinding was adequately maintained.

At the 12-month visit, the percentage of informants who believed that participants were taking active study medication also did not differ across the participants' treatment group (52% for rofecoxib group, 51% for naproxen group, and 59% for placebo group; $P = .10$, Fisher exact test). The differences also were not significant for study coordinators (54% for rofecoxib group, 50% for naproxen group, and 43% for placebo group; $P = .52$, Fisher exact test) and study physicians (56% for rofecoxib group, 40% for naproxen group,

EFFECTS OF ROFECOXIB OR NAPROXEN VS PLACEBO ON ALZHEIMER DISEASE

Table 4. Number of Serious Adverse Events

	Placebo	Naproxen	Rofecoxib
Total serious events	15	25	26
Deaths	1	1	2
Gastrointestinal tract event*	1	2	4
Stroke/transient ischemic attack	1	3	3
Subdural hematoma	0	2	1
Myocardial infarction	1	0	3

*Serious gastrointestinal tract events included 6 cases of gastrointestinal tract bleeding and 1 case of perforated small intestine (the latter occurred in the naproxen group).

and 40% for placebo group; $P=.19$, Fisher exact test).

COMMENT

This is the first large-scale trial, to our knowledge, to determine the efficacy of both nonselective and COX-2 selective NSAIDs in the treatment of AD. The result of the primary intent-to-treat analysis of this trial showed that neither naproxen nor rofecoxib slowed the rate of cognitive decline in comparison with placebo in participants with mild-to-moderate AD. Similar results were obtained when an LOCF analysis was substituted for the primary slope method of imputing missing 12-month data and when the data were analyzed using the generalized estimating equations method. The results of an analysis of study completers (those who were able to complete the trial while still taking study medication) also revealed no beneficial treatment effect of the active study drugs in comparison with placebo. Instead, the rofecoxib group showed a trend toward greater cognitive decline compared with placebo.

In the naproxen group, baseline MMSE and CDR-SOB scores differed between individuals who discontinued study participation early and those who completed the study. This difference raises the possibility that imputation of the data missing from non-completers may have influenced the outcome of the primary analysis. However, the nonsignificant finding of the primary analysis is in agreement with the nonsignificant findings of the LOCF analysis, the completers analysis, the generalized estimating equations regression analysis, and the clinical end

points survival analysis, which strongly support its validity.

Secondary analyses of the effect of treatment on a global measure of cognitive decline, behavior, ADL, and quality of life showed no consistent advantage of either treatment over placebo. Rofecoxib delayed the time to institutionalization, but this result was based on a small number of end points. Adjustment for multiple comparisons would have made this comparison nonsignificant.

In some analyses, there was a trend toward smaller decline in ADL with active treatment, particularly with rofecoxib. The trend toward greater cognitive decline with rofecoxib treatment suggests that the trend toward protection against decline in ADL may be related to an analgesic effect rather than an effect on AD.

Both active treatments were reasonably well tolerated, although participants who dropped out due to adverse events were more commonly from the active drug groups than from the placebo group. Surprisingly, the gastrointestinal tract symptoms were similar in the 3 groups. The small number of serious gastrointestinal tract events (primarily bleeding) in the active drug groups was consistent with expected risks. The occurrence of adverse events likely was reduced by excluding individuals at high risk of NSAID toxicity.

Two important issues must be raised when interpreting the lack of active treatment effect. The first issue concerns selection of drug, dose, and duration of treatment. Naproxen was selected because the epidemiological evidence supports the efficacy of non-

selective COX inhibitors, and naproxen has a long half-life (allowing for twice-daily dosing) and is relatively well tolerated. The naproxen dose was low because of concern that full-dose therapy with a nonselective COX inhibitor would be associated with more serious adverse events and a high drop-out rate compared with placebo. The nonsignificant results with naproxen could be explained by an insufficient dose; alternatively, other drugs in this same class, such as ibuprofen, may have greater activity in AD mediated by an effect on processing of the amyloid precursor protein.²⁴ Rofecoxib, 1 of 2 available selective COX-2 inhibitors at the time this study was initiated, was administered at a standard, full anti-inflammatory dose. It is possible that a higher dose is necessary to influence neuronal survival; a high dose of celecoxib, above the standard anti-inflammatory dose, is required to influence neoplastic transformation in the gastrointestinal tract.²⁵ It is also possible that a period of exposure to anti-inflammatory treatment greater than the 1 year of treatment in this trial is necessary to slow the disease process. Some epidemiological studies of AD prevention suggest that at least 2 years of exposure is necessary to optimally reduce risk of AD.^{26,27}

The second issue concerns the selection of participants. The most compelling epidemiological evidence suggests that long-term exposure to NSAIDs reduces risk of subsequent AD; on the other hand, some studies suggest a stabilizing effect of NSAID therapy in patients with clinically evident disease,^{3,28} supporting the design of the current trial. While COX-2 and other inflammatory mediators are elevated in mild-to-moderate AD brain,¹ it is nonetheless possible that when the disease is clinically apparent, the neuropathology is too advanced to be significantly attenuated by NSAID therapy. The results of this trial do not address the efficacy of NSAIDs in the prevention of AD. A primary prevention trial, in which elderly persons without dementia are randomly assigned to treat-

EFFECTS OF ROFECOXIB OR NAPROXEN VS PLACEBO ON ALZHEIMER DISEASE

ment with an NSAID or placebo, is necessary to test the utility of long-term NSAID therapy in AD risk reduction. Such a trial is now under way.²⁰

The results of the current study do not support the hypothesis that rofecoxib or naproxen can slow the progression of AD. Considering the risk of serious toxicity, such treatment should not be recommended. In view of evidence from cell culture and animal studies suggesting reduction of β -amyloid generation with other NSAIDs, such as ibuprofen (but not naproxen or selective COX-2 inhibitors),²⁴ additional treatment trials using other NSAIDs may be warranted. Recommendations regarding risk reduction with long-term NSAID therapy must await the results of a primary prevention trial.

Author Affiliations: Department of Neurology, Georgetown University Medical Center, Washington, DC (Dr Alsen); Department of Neurosciences, University of California, San Diego (Ms Schafer and Drs Grundman, Jin, and Thal); Suncoast Gerontology Center, University of South Florida College of Medicine, Tampa (Dr Pfeiffer); Department of Psychiatry (Dr Sano), Mount Sinai School of Medicine, Mount Sinai Medical Center (Dr Davis), New York, NY; Department of Neurology, School of Medicine, Indiana University, Indianapolis (Dr Farlow); Departments of Family and Preventive Medicine and Neurosciences, University of California, San Diego (Dr Thomas).

Author Contributions: Study concept and design: Alsen, Grundman, Sano, Davis, Thomas.

Acquisition of data: Alsen, Schafer, Grundman, Sano, Farlow, Thal.

Analysis and interpretation of data: Alsen, Grundman, Pfeiffer, Jin, Thomas, Thal.

Drafting of the manuscript: Alsen, Schafer, Grundman, Davis, Thomas, Thal.

Critical revision of the manuscript for important intellectual content: Alsen, Grundman, Pfeiffer, Sano, Farlow, Jin, Thomas, Thal.

Statistical expertise: Jin, Thomas.

Obtained funding: Alsen, Thomas, Thal.

Administrative, technical, or material support: Alsen, Schafer, Grundman, Pfeiffer, Farlow, Thomas, Thal.

Study supervision: Alsen, Davis, Thomas.

Participating Investigators and Staff: Oregon Health Sciences University, Portland: Jeff Kaye, MD, Joyce Lear, RN, MN, Joseph Quinn, MD; University of Southern California, Los Angeles: Lon Schneider, MD, Nansi Taggart, RN, BSN, MA, Sonia Pawluczyk, MD, Karen Stevens-Dagerman, MA; University of California, San Diego: Mary Margaret Pay, RNC, NP, John Olchney, MD, Ingrid Padilla BA, University of Michigan, Ann Arbor: Judith L. Heidebrink, MD, MS, Laurie A. Blumenthal, MS, RN, Norman L. Foster, MD; University of Washington, Seattle: Elaine R. Peskind, MD, Linda C. Mandelco, BS, Nancy A. Brown, CCRCC, Nghia T. Pham; Baylor College of Medicine, Houston: Rachelle Smith Doody, MD, PhD, Jamie Simms, Nikita Robinson, BS; Columbia University, New York: Karen Bell, MD, Ruth Tejada, MD, Gina Camilo, MD; Washington University, St. Louis: John C. Morris, MD, Christy Tomlinson, MSN, RN, Carol Dyer, PhD; Rush-Presbyterian-St. Luke's Medical Center, Chicago, Ill: Neelum Aggarwal, MD, Rose Marie Ferraro, LVN, Raj

Shah, MD; University of Minnesota, Minneapolis: David S. Knopman, MD, Margaret Patterson, Nancy Haukom, RN, BAN, Martha Mandarino, BA; Wein Center, Miami: Ranyan Duara, MD, Peggy Roberts, Peggy Imier, RN; University Hospitals of Cleveland, Ohio: David Goldmacher, MD, Lani Holland, MA, Alan Lemmer MD, Marianne Sanders, RN; University of South Florida, Tampa: Enc Pfeiffer, MD, Barbara Luhn, MSW, Dottie Baxter, MPH; New York University Medical Center, New York: Steven H. Ferris, PhD, Maria A. Vlasopoulos, BA, Emile Franssen, MD; University of Pennsylvania, Philadelphia: Christopher M. Clark, MD, Kris Gravanda, BA, Jessica Nuñez; University of Kentucky, Lexington: Frederick Schmitt, PhD, Eileen Tekrony, RNA, Charles Smith, MD; University of Pittsburgh, Pa: Steven DeKosky, MD, Jennifer Hagerty, MSN, CRNP, Timothy Shelley, MSW, LSW, Thomas Baumgartner, MSW, LSW; University of Rochester Medical Center, NY: Pierre Tariot, MD, Elizabeth Smith, BSN, J. Michael Ryan, MD, Bonnie Goldstein, MS, RNCS; University of California, Irvine: Ruth A. Mulnard, RN, DNSc, Catherine McAdams-Ortiz, RN, MSN, Hyunmi Kim, RN, MSN; University of Texas-Southwestern, Dallas: Doris Svetlik, BSN, MS, Stephen Vobach, MD, Myron Weiner, MD; Emory University, Atlanta, Ga: Allan Levey, MD, PhD, Janet Cellar, RN, MSN, Nancy Gauchman, BS; Kansas University, Kansas City, Mo: Grisel J. Lopez, MD, Jill Peltzer, RN, Phyllis Switzer; Vanderbilt University, Nashville, Tenn: Richard Margolin, MD, Diane Kent, RN; University of California, Los Angeles: Donna Masternan, MD, Michele Carter, RN, Susan O'Connor, RN; Mayo Clinic, Jacksonville, Fla: Neill R. Graff-Radford, MD, Francine C. Parfitt, MSC, CCRCC, Lauren M. Makarov; Indiana University, Indianapolis: Martin Farlow MD, Nicki Coleman RN, Jean Pearson BS; Memorial Hospital of Rhode Island, Pawtucket: Brian R. Ott, MD, Robert H. Paul, PhD, Ronald A. Cohen, PhD; Yale University School of Medicine, New Haven, Conn: Christopher H. van Dyck, MD, Martha G. MacAvoy, PhD, Tracy E. Righmer; University of California, Davis: Jean Coleman, RN, MSN, Ananth Acharya, MD, Bruce Reed, PhD; University of Arizona, Tucson: Geoffrey Ahern, MD, PhD, Janet Mar, RN; Baumer-Eisner Institute, Boca Raton, Fla: Larry S. Eisner, MD, Janella E. Crasto, MD, Rebecca Radzwill, MS; Baumer-Eisner Institute, Ft Lauderdale, Fla: Larry S. Eisner, MD, Maggie Szymczak, MD, Amy Johnson, MA; Baumer-Eisner Institute, Miami Beach, Fla: Juan Carlos Jaramillo, Rafael E. Comelio, MD, Damaris Alorson, MS; Southwestern Vermont Medical Center, Bennington: Paul Solomon, PhD, Cynthia Murphy, MBA, Betty Hathaway, Patty Flynn Rosenthal; University of Nevada School of Medicine, Las Vegas: Marie Stallbaum, RN, Donna Muncie, PhD, Gail Vranesh, RN; Barrow Neurology Group, Phoenix, Az: Charles L. Echols, MD, Marc Zomok, RN, Judith Lawrence, RN; Medical University of South Carolina, North Charleston: Jacobo Mintzer, MD, Don Bagwell, BS, Kern Costa, RN; ClinSearch, Summit: Mark Roffman, PhD, Ina Marntt, BS, Robert Moreines, MD; Premiere Neurological Group, Palm Beach, Fla: Carl Sadowsky, MD, Joe Piccinilli, RN, Walter Martinez, MD; Georgetown University, Washington DC: Carolyn Ward, MSPH, Mary Ann Cachola, Sarah Gustafson, BSN; Alzheimer's Disease Cooperative Study Coordinating Center, San Diego, Calif: Jackie Bochenek, MS, Anthony Camst, PhD, Karin Enstrom, MA, Rachelle Tractenberg, PhD, Julie Berg, MS, Feng He

Data and Safety Monitoring Board: Karl Kieburtz, MD, Trey Sunderland, MD, Bruce Miller, MD, Richard Kryscio, PhD.

Clinical Monitors: Kristin Broll, BA, Karen Croft, BA, Wes Houston, PhD, Virginia McClintock, MS, Natalie Melton, MS, Amy Rossi, MPH, Rebecca Ryan-Jones, PhD, Pamela Saunders, PhD, MaMo Schittini, MD, MPH.

Funding/Support: This study was supported by a grant

(U01-AG10483) from the National Institute on Aging, as well as grants (including 5 M01-RR13297 to Georgetown University) from the General Clinical Research Center Program of the National Center for Research Resources, National Institutes of Health. Naproxen and matching placebo were provided by Bayer Consumer, Inc.

Disclaimer: The study design was approved by an oversight committee of the National Institute on Aging. Bayer Consumer, Inc. supplied naproxen sodium, but did not participate in the design of the study, analysis or interpretation of the data, or preparation of the manuscript; a copy of the manuscript was submitted to Bayer Consumer, Inc. prior to publication.

REFERENCES

1. Akiyama H, Barger S, Barnum S, et al. Inflammation and Alzheimer's disease. *Neurobiol Aging*. 2000; 21:383-421.
2. McGeer PL, Schulzer M, McGeer EG. Arthritis and anti-inflammatory agents as possible protective factors for Alzheimer's disease: a review of 17 epidemiologic studies. *Neurology*. 1996;47:425-432.
3. Rogers J, Kirby LG, Hempelman SR, et al. Clinical trial of indomethacin in Alzheimer's disease. *Neurology*. 1993;43:1609-1611.
4. Scharf S, Mander A, Ugoni A, Valda F, Christophidis N. A double-blind, placebo-controlled trial of diclofenac/misoprostol in Alzheimer's disease. *Neurology*. 1999;53:197-201.
5. Alsen PS, Davis KL, Berg JD, et al. A randomized controlled trial of prednisone in Alzheimer's disease: Alzheimer's Disease Cooperative Study. *Neurology*. 2000;54:588-593.
6. Van Gool WA, Weinstein HC, Scheltens PK, Walstra GJ. Effect of hydroxychloroquine on progression of dementia in early Alzheimer's disease: an 18-month randomised, double-blind, placebo-controlled study. *Lancet*. 2001;358:455-460.
7. In 't Veld BA, Ruttenberg A, Hoffman A, et al. Non-steroidal antiinflammatory drugs and the risk of Alzheimer's disease. *N Engl J Med*. 2001;345:1515-1521.
8. Lim GP, Yang F, Chu T, et al. Ibuprofen suppresses plaque pathology and inflammation in a mouse model for Alzheimer's disease. *J Neurosci*. 2000;20:5709-5714.
9. Lim GP, Yang F, Chu T, et al. Ibuprofen effects on Alzheimer pathology and open field activity in APPsw transgenic mice. *Neurobiol Aging*. 2001;22:983-991.
10. Pasinetti GM, Alsen PS. Cyclooxygenase-2 expression is increased in frontal cortex of Alzheimer's disease brain. *Neuroscience*. 1998;87:319-324.
11. Bazan NG, Lukiw WJ. Cyclooxygenase-2 and presenilin-1 gene expression induced by interleukin-1 beta and amyloid beta 42 peptide is potentiated by hypoxia in primary human neural cells. *J Biol Chem*. 2002; 277:30359-30367.
12. Andreasson KI, Savonenko A, Videnko S, et al. Age-dependent cognitive deficits and neuronal apoptosis in cyclooxygenase-2 transgenic mice. *J Neurosci*. 2001;21:8198-8209.
13. Xiang Z, Ho L, Valdellon J, et al. Cyclooxygenase (COX)-2 and cell cycle activity in a transgenic mouse model of Alzheimer's disease neuropathology. *Neurobiol Aging*. 2002;23:327-334.
14. Ho L, Purohit D, Haroutunian V, et al. Neuronal cyclooxygenase 2 expression in the hippocampal formation as a function of the clinical progression of Alzheimer disease. *Arch Neurol*. 2001;58:487-492.
15. McGeer PL, McGeer EG. Inflammation, autotoxicity and Alzheimer disease. *Neurobiol Aging*. 2001; 22:799-809.
16. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group. *Neurology*. 1984;34:939-944.
17. Folstein M, Folstein S, McHugh P. The Mini-

A Randomized, Double-Blind, Study of Rofecoxib in Patients with Mild Cognitive Impairment

Leon J Thal¹, Steven H Ferris², Louis Kirby³, Gilbert A Block⁴, Christopher R Lines⁴, Eric Yuen⁴, Christopher Assaid⁴, Michael L Nessly⁴, Barbara A Norman⁴, Christine C Baranak⁴ and Scott A Reines⁴, on behalf of the Rofecoxib Protocol 078 study group⁵

¹University of California, San Diego, CA, USA; ²New York University School of Medicine, New York, NY, USA; ³Pivotal Research Centers, Peoria, AZ, USA; ⁴Merck Research Laboratories, West Point, PA, USA

Inflammatory mechanisms have been implicated in Alzheimer's disease (AD) and might be mediated via the COX-2 enzyme. Previous studies with the selective COX-2 inhibitors, rofecoxib and celecoxib, have shown that they do not alter the progression of AD. We conducted a double-blind study to investigate whether rofecoxib could delay a diagnosis of AD in patients with mild cognitive impairment (MCI), a group with an expected annual AD diagnosis rate of 10–15%. MCI patients ≥ 65 years were randomized to rofecoxib 25 mg ($N = 725$) or placebo ($N = 732$) daily for up to 4 years. The primary end point was the percentage of patients with a clinical diagnosis of AD. The estimated annual AD diagnosis rate was lower than the anticipated 10–15%; 6.4% in the rofecoxib group vs 4.5% in the placebo group (rofecoxib:placebo hazard ratio = 1.46 (95% CI: 1.09, 1.94), $p = 0.011$). Analyses of secondary end points, including measures of cognition (eg the cognitive subscale of the AD Assessment Scale (ADAS-Cog)) and global function (eg the Clinical Dementia Rating (CDR)), did not demonstrate differences between treatment groups. There was also no consistent evidence that rofecoxib differed from placebo in post hoc analyses comparing ADAS-Cog and CDR-sum of boxes scores in overlapping subgroups of patients who had Mini Mental State Exam scores of 24–26 in the present MCI study and in a previous AD treatment study with a similar design. The results from this MCI study did not support the hypothesis that rofecoxib would delay a diagnosis of AD. In conjunction with the lack of effects observed in previous AD studies, the findings suggest that inhibition of COX-2 is not a useful therapeutic approach in AD.

Neuropsychopharmacology (2005) 30, 1204–1215, advance online publication, 2 March 2005; doi:10.1038/sj.npp.1300690

Keywords: rofecoxib; nonsteroidal anti-inflammatory drugs; mild cognitive impairment; Alzheimer's disease; COX-2 inhibitors

INTRODUCTION

Neuropathological and epidemiological data suggest that inflammatory mediators and immune mechanisms may play a role in the pathogenesis of Alzheimer's disease (AD) (McGeer *et al*, 1996; Cagnin *et al*, 2001; Ho *et al*, 2001; in 't Veld *et al*, 2002; Elminan *et al*, 2003). It has been proposed that these effects may be mediated via the COX-2 enzyme (Pasinetti, 2001). These observations have led researchers to investigate whether nonsteroidal anti-inflammatory drugs (NSAIDs) might slow the progression of dementia. Initially, two small studies with the NSAIDs indomethacin (Rogers

et al, 1993) and diclofenac (Scharf *et al*, 1999) provided preliminary evidence for efficacy over 6 months in patients with AD. However, interpretation of the results from both studies was confounded by high dropout rates, mainly due to gastrointestinal side effects thought to be mediated via inhibition of COX-1 (nonselective NSAIDs inhibit both COX-1 and COX-2) (Warner *et al*, 1999). More recently, three larger randomized, controlled, 1-year clinical studies with the selective COX-2 inhibitors rofecoxib or celecoxib failed to show any effects of treatment on the progression of AD (Sainati *et al*, 2000; Aisen *et al*, 2003; Reines *et al*, 2004). One of these studies included the nonselective NSAID naproxen, which also failed to show any effects (Aisen *et al*, 2003). In addition, the anti-inflammatory agent hydroxychloroquine did not show any benefits in an 18-month trial (Van Gool *et al*, 2001).

A possible explanation for the lack of efficacy in the above studies is that the underlying pathology might be too advanced in patients with an established diagnosis of AD for an anti-inflammatory treatment to alter the course of the disease. Epidemiological evidence indicates that there may

*Correspondence: Dr CR Lines, Merck Research Laboratories, 10 Sentry Parkway, Blue Bell, PA 19422, USA, Tel: +1 484 344 2595, Fax: +1 484 344 4940, E-mail: chris_lines@merck.com

⁵Members of the Rofecoxib Protocol 078 study group are listed in Acknowledgments

Received 8 March 2004; revised 16 December 2004; accepted 29 December 2004

Online publication: 12 January 2005 at <http://www.acnp.org/citations/NPP011205040106/default.pdf>

be a critical period, 2 or more years before the onset of dementia, during which exposure to NSAIDs protects against AD (in 't Veld *et al*, 2001). Selective COX-2 inhibitors offer the potential for treatments that may have beneficial protective effects in AD while being well tolerated in long-term use. We therefore conducted a study to determine whether treatment with rofecoxib could delay a diagnosis of AD. Rather than evaluating a general elderly population with a relatively low incidence rate of AD, we sought to recruit elderly patients with mild cognitive impairment (MCI) (Petersen *et al*, 2001a, b). These patients were expected to have an annual AD diagnosis rate of 10–15% vs a rate of 1–2% for the general elderly population (Petersen *et al*, 2001a, b). The study also provided the opportunity to gather important placebo-controlled, long-term safety data on rofecoxib in an elderly population.

METHODS

Patients

Patients aged 65 years or older who had completed at least 8 grades of education, and had a reliable informant who could accompany them to each clinic visit, were recruited at 46 study sites in the United States from April 1998 to March 2000. Potentially eligible patients were initially identified by investigators (by any means available) or via a centralized telephone prescreening process (Lines *et al*, 2003). Patients were screened at the study sites to determine if they met all the following criteria for MCI: patient reports memory problem; or informant reports that patient has memory problem; informant reports that patient's memory has declined in the past year; Mini Mental State Exam (MMSE) (Folstein *et al*, 1975) score ≥ 24 ; Clinical Dementia Rating (CDR) (Morris, 1993) global score ≤ 0.5 with memory domain score ≥ 0.5 ; Blessed Dementia Rating Scale (BDRS) (Morris *et al*, 1988) total score ≤ 3.5 , with no part 1 item score > 0.5 ; Auditory Verbal Learning Test (AVLT) (Schmidt, 1996) total score ≤ 37 . The cut score on the AVLT corresponds to a score ≤ 1 standard deviation (SD) below the mean for normal elderly subjects; for the first 6 months of study enrollment, age-adjusted cut scores 1.5 SDs below the means for separate age bands were used (see Appendix A1). Further details of these tests of cognition and function are given in Appendix A1. The CDR assessment was completed on the basis of interviews with the patient and informant by a rater who was blinded to the results of psychometric tests (AVLT and MMSE).

Patients were excluded if they had: dementia; inadequate motor or sensory capacities to comply with testing; a modified Hachinski Ischemic Scale (Rosen *et al*, 1980) score > 4 (to exclude patients whose cognitive impairment may have been related to a vascular condition); a Hamilton Depression Scale (17-item version) (Hamilton, 1960) score > 13 (to exclude patients whose cognitive impairment may have been related to depression); a history of angina or congestive heart failure with symptoms that occurred at rest; uncontrolled hypertension; a history within the past year of myocardial infarction, coronary artery bypass, angioplasty, or stent placement; a history within the past 2 years of stroke, multiple lacunar infarcts, or transient ischemic events; a history within the past 3 months of

Rofecoxib in mild cognitive impairment

LJ Thal *et al*

1205

gastrointestinal bleeding; an expected therapeutic need for chronic NSAID or estrogen replacement therapy during the study. Patients taking NSAIDs on a chronic basis (≥ 7 days/month for the 2 months prior to study entry), estrogen replacement therapy (excluding topical ointments) within 2 months of study entry, or cholinesterase inhibitors within 1 month of study entry were also excluded. No more than 20% of patients at each study site could be taking vitamin E > 400 IU at the time of study entry. Concomitant use of the above medications during the study was discouraged, but patients who did take them were not discontinued for this reason. Patients who developed a need for cardio-protective doses of aspirin after randomization were permitted to use aspirin ≤ 100 mg/day; clopidogrel was also allowed. Each site received the approval of its local institutional review board to perform the study, and informed consent was obtained from each subject.

Design

This was a randomized, double-blind, placebo-controlled study with parallel groups. After screening, eligible patients were randomly assigned to receive rofecoxib 25 mg once daily or placebo once daily for up to 4 years. Rofecoxib 25 mg was the maximum recommended dose for approved chronic indications (Merck & Co. Inc., 2003; on September 30, 2004, Merck & Co. Inc., announced the voluntary worldwide withdrawal of rofecoxib from the market). Randomization of patients at each study site was determined by a computer-generated allocation schedule and was stratified according to MMSE score (24–26, > 26). The allocation schedule was generated by a statistician at Merck Research Laboratories according to in-house blinding conditions. The rofecoxib and placebo tablets were visually identical. The original intention was that the study would be event-rate driven and would continue until 220 patients had received an AD diagnosis, which it was anticipated would take approximately 2 years. The study was extended to 4 years due to lower than expected AD diagnosis rates. A decision was made in October 2002 to terminate the study in April 2003, 11 months earlier than the scheduled March 2004 termination (ie 4 years after the last patient was enrolled). Based on the diagnosis and discontinuation rates observed as of July 2002, it was estimated that 219 end points should have been observed by April 2003. In fact, only 189 diagnoses of AD had been made at this time, and discontinuation rates had continued to increase. The combination of the low diagnosis rates and increasing discontinuation rates made it unlikely that the target number of events could have been achieved and it was therefore considered that continuation of the study until its planned termination would not have been productive. The decision was made prior to unblinding.

Procedure

After randomization, patients were scheduled to attend the clinic at months 1 and 4, and then every 4 months until the study was completed or the patient was diagnosed with dementia. The study was conducted using an intent-to-treat approach; that is, patients who discontinued treatment, but who had not developed dementia, were asked to return to

the clinic for all remaining visits and assessments. The following assessments were administered at baseline and every 4 months, or at discontinuation from the study: CDR, MMSE, Selective Reminding Test (SRT; see Appendix A1 for details) (Buschke, 1973). The cognitive subscale of the AD Assessment Scale (ADAS-Cog; see Appendix A1 for details) (Rosen et al, 1984) was administered at baseline and then every 12 months, or at discontinuation from the study. The BDRS was administered at baseline and at 24, 36, and 48 months, or at discontinuation from the study. Any patient who received a global CDR score ≥ 1 at a routine clinic visit was suspected of having converted to dementia and administered a CT or MRI scan of the brain and any psychometric tests that were not already scheduled for that visit. The patient was continued on treatment and returned for an end point confirmation visit 2 months later, at which time all the psychometric evaluations were repeated. If the global CDR score was still ≥ 1 at this visit, the patient was considered to have reached the end point of dementia and was discontinued from the study (regardless of the outcome of the adjudication process described below). In some cases, a patient was determined by an investigator to have developed dementia despite maintaining a global CDR score of 0.5 at the trigger and confirmation visits, and these patients were also counted as end points and were discontinued from the study. The investigator determined the type of dementia: possible or probable AD according to NINCDS-ADRDA criteria (McKhann et al, 1984) or other, for example, vascular dementia. For patients who reached the end point of clinically diagnosed dementia, all relevant data were sent to an independent blinded adjudication committee consisting of three experts. Each adjudicator reviewed the data independently and indicated whether or not they concurred with the investigator's diagnosis. In order to qualify as an event for the primary analysis, a majority decision in ≥ 2 of the three adjudicators that the patient met criteria for possible or probable AD was required (ie ≥ 2 adjudicators classified the event as probable AD, ≥ 2 adjudicators classified the event as possible AD, or one adjudicator classified the event as possible AD and one adjudicator classified it as probable AD).

Any adverse experiences occurring during the study were recorded and rated by the investigator, while still blinded to the treatment that the patient was receiving, as to seriousness (death, life threatening, resulting in persistent or significant disability, resulting in hospitalization, prolonging an existing hospitalization, any other important medical event), drug-relatedness (possibly, probably or definitely drug-related, probably not or definitely not drug-related), and intensity (mild, moderate, or severe). All serious vascular events (including cardiac, peripheral vascular, and cerebrovascular events) and upper gastrointestinal perforations, ulcers, and bleeds were reviewed by independent blinded adjudication committees, who determined if they were confirmed events according to prespecified case definitions (confirmed events) (Bombardier et al, 2000; Konstam et al, 2001).

Statistical Analysis

The primary efficacy analysis compared the cumulative incidence of possible or probable AD according to NINCDS-

ADRDA criteria (McKhann et al, 1984); patients with dementia of other cause were censored in the analysis at the time of diagnosis. AD diagnoses confirmed by the end point adjudication committee were the only end points included in the primary analysis. The analysis was based on a Cox proportional hazards model of time-to-event data (based on the initial diagnosis of AD) using an intention-to-treat approach, which included all randomized patients regardless of whether or not they were taking study medication. The model included terms for treatment, region within the United States (North East, South East, South, Midwest, West), and baseline MMSE strata (baseline MMSE score ≤ 26 vs > 26). Region was intended to be a surrogate for investigating the influence of study site, since there were too few events at individual study sites to include that as a factor. An additional prespecified on-drug analysis was restricted to patients who converted to AD within 14 days of being on study medication.

The calculation for the power statement assumed that the incidence of possible or probable AD over 2 years in the placebo group would be 30% and that the discontinuation rate would be 20%. Based on these considerations and a planned sample size of 520 evaluable patients per group, the study had 90% power to detect a one-third reduction in the incidence of AD in the rofecoxib group vs the placebo group with two-tailed $\alpha = 0.05$.

Analyses of prespecified secondary measures (SRT-summed recall score, SRT-delayed recall score, MMSE score, ADAS-Cog score, CDR-sum of boxes score, BDRS score) were based on available data for evaluable patients (ie patients who had a baseline score and at least one postrandomization score). The annual rates of change from baseline to a given time point (slopes), and slope differences between groups, were estimated using an intention-to-treat approach (ie including all data regardless of whether a patient discontinued from therapy or not) and analyses were performed using longitudinal repeated measures models for the comparison, under the assumption that missing data were missing at random (ie ignorable missingness). Additional sensitivity analyses were also performed using a last-observation-carried forward approach, as well as with other models with different assumptions about the missing data structure.

The present study had a broadly similar design (randomized, double-blind, placebo-controlled) and utilized some similar assessments (ADAS-Cog and CDR) to a previous 1-year AD treatment study of rofecoxib 25 mg in patients who had MMSE scores of 14–26 (Reines et al, 2004). Since there were overlapping subgroups of patients who had MMSE scores of 24–26 in both studies, indicating a similar level of cognitive impairment despite the difference in diagnosis, we also conducted a *post hoc* analysis to compare change from baseline scores on measures of cognition (ADAS-Cog) and global function (CDR-sum of boxes) in the overlapping subgroups in the two studies. The analysis looked at estimated annual slope differences using the same methods as described above. For the subgroup from the previous AD treatment trial, the annual slope estimates were derived from data over 12 months. For the subgroup from the present study, the annual slope estimates were derived from data over 48 months.

The assessment of tolerability included on-drug adverse experiences with onset up to 14 days after patients stopped taking test medication, and was based on the population of patients who took at least one dose of study medication. Prespecified groupings of adverse experiences (eg the number of patients with one or more adverse experience) were analyzed using Fisher's exact test.

RESULTS

The study profile is shown in the study flowchart (Figure 1). A total of 1457 patients were randomized. The study was terminated after 189 confirmed diagnoses of AD had been made. The median duration of study participation was 115 weeks in the rofecoxib group and 130 weeks in the placebo group. The median duration patients took study medication was 94 weeks in the rofecoxib group and 105 weeks in the

placebo group. Estimates of treatment compliance based on counts of the number of tablets in medication bottles at each clinic visit and calculated as ((number of days on therapy/number of days in study) × 100) indicated reasonable compliance; 61.0% of the 725 patients randomized to rofecoxib and 70.8% of the 732 patients randomized to placebo had ≥ 80% compliance during the time they were in the study. Approximately 45% of patients discontinued the study prematurely, while 40% completed the study on-drug (including patients who had not completed 48 months of treatment but were still in the study at the time of termination), and 15% completed the study off-drug. Reasons for discontinuation were generally similar across the treatment groups, although there were some small differences; for example, a greater proportion of patients in the placebo group discontinued due to withdrawal of consent (see Figure 1). The time course of discontinuations was similar between the treatment groups over the duration




Figure 1 Study flowchart. Patients were prescreened for eligibility before being screened in the clinic. Data on the precise number of subjects prescreened were not collected, but the number was > 17000. The total who completed the study includes patients who completed 48 months of study participation ($N = 115$ for rofecoxib, $N = 146$ for placebo), patients who were diagnosed with dementia of any cause ($N = 112$ for rofecoxib, $N = 83$ for placebo), patients who completed less than 48 months of study participation but were still in the study at the time it was terminated ($N = 121$ for rofecoxib, $N = 123$ for placebo), and patients who were discontinued because the study site closed ($N = 52$ for rofecoxib, $N = 49$ for placebo).

Rofecoxib in mild cognitive impairment

LJ Thal et al

1208

of the study (data not shown). Of those who discontinued, placebo patients had higher baseline ADAS-Cog scores, indicative of greater cognitive impairment and possibly greater risk of progression to AD, than patients randomized to rofecoxib ($p=0.034$ in a logistic regression model); the mean baseline ADAS-Cog score was 9.6 in placebo discontinuers vs 9.1 in rofecoxib discontinuers. The change from baseline to last evaluation time point scores on the ADAS-Cog, MMSE, and CDR-sum of boxes in the subgroup of discontinued patients who had at least one on-treatment evaluation was similar between the treatment groups (data not shown).

Table 1 shows the baseline demographic characteristics and baseline test scores for randomized patients. The groups were generally comparable with regard to baseline characteristics and test scores. The distribution of secondary diagnoses at baseline was generally similar between the groups. The most common pre-existing medical conditions were hypertension (37.7% in the rofecoxib group and 34.3% in the placebo group) and osteoarthritis (22.2% in the

rofecoxib group and 24.3% in the placebo group). The percentages of patients with reported prior use (for any length of time in the 2 months prior to study entry for NSAIDs and estrogen, and 1 month prior to study entry for other drugs) of drugs claimed to have an influence on dementia were generally similar in the rofecoxib vs placebo groups for NSAIDs (8.4 vs 10.2%), ginkgo (12.6 vs 11.2%), statins (14.9 vs 13.0%), estrogen (0.4 vs 1.4%), and vitamin E >400 IU daily (5.8 vs 5.6%). The percentages of patients with reported concomitant use (for any length of time during the study) of drugs claimed to have an influence on dementia were lower in the rofecoxib group than the placebo group for NSAIDs (30.9 vs 34.7%) and estrogen (2.6 vs 4.8%), higher for cholinesterase inhibitors (11.2 vs 8.6%), and similar for statins (24.8 vs 24.9%), ginkgo (13.7 vs 12.8%), and vitamin E >400 IU daily (7.3 vs 7.5%). The median duration of reported concomitant NSAID use was lower in the rofecoxib group (5.6 weeks) than the placebo group (7.4 weeks). The percentages of patients with reported aspirin use in the 2 months prior to the study were 10.1% in the rofecoxib group and 9.0% in the placebo group. The percentages of patients with reported aspirin during the study were 31.7% in the rofecoxib group and 29.9% in the placebo group.

A total of 195 investigator diagnoses of dementia (190 diagnoses of possible or probable AD and five diagnoses of non-AD dementia) were evaluated by the end point adjudication committee. The primary end point of clinically diagnosed AD included the 189 patients who were confirmed to have developed possible or probable AD by ≥ 2 members of the committee. Two events adjudicated to be non-AD dementia, two events adjudicated to be nondementia, and two events in which there was no agreement between adjudicators (one adjudicator classified the event as AD, one classified it as non-AD dementia, and one classified it as nondementia) were not included as end points in the primary analysis. In the 189 patients with adjudicator-confirmed clinically diagnosed AD, global CDR scores at the last clinic visit were 0.5 (questionable dementia) in 21 patients, one (mild dementia) in 159 patients, two (moderate dementia) in eight patients, and three (severe dementia) in one patient.

In the rofecoxib group, 107 of 725 (14.8%) patients had clinically diagnosed AD over the 4-year study period vs 82 of 732 (11.2%) placebo patients over 4 years. The estimated hazard ratio (rofecoxib: placebo), adjusting for the effects of baseline MMSE stratum and region, was 1.46 (95% CI: 1.09, 1.94), which was statistically significant ($p=0.011$, Wald χ^2 test) in favor of placebo. The treatment-by-time interaction was not significant ($p=0.260$), indicating that the proportional hazards assumption was reasonably met. Baseline MMSE stratum (score ≤ 26 ; score >26) had a highly statistically significant effect on outcome with an estimated hazard ratio of 3.23 (95% CI: 2.42, 4.32) ($p<0.001$), indicating that patients in the lower stratum were much more likely to progress to AD irrespective of treatment assignment. Treatment effects were consistent across MMSE stratum levels, as well as geographic regions. Kaplan-Meier estimated proportions of patients with clinically diagnosed AD at 4-month increments are shown in Table 2. A separation in event rates between treatment groups was evident at the earliest time point (4 months), a gap that was

Table 1 Baseline Patient Characteristics

Characteristic	Placebo (N = 732) ^a	Rofecoxib 25 mg (N = 725) ^a
% Women	31.1%	34.3%
% Family history of AD	29.5%	31.2%
% Apolipoprotein <i>ε</i> 4	35.8%	34.6%
Years of education		
11 or less	9.8%	11.2%
12-17	77.7%	74.7%
18 or more	12.4%	13.4%
Mean (SD) years of age	74.8 (6.0)	75.1 (6.0)
Mean (SD) AVLT score ^b	28.6 (5.8)	28.6 (5.7)
Mean (SD) MMSE score ^c	27.3 (1.7)	27.4 (1.7)
Mean (SD) CDR-sum of boxes score ^d	1.4 (0.8)	1.4 (0.8)
Mean (SD) ADAS-Cog score ^e	9.1 (3.8)	9.2 (4.0)
Mean (SD) SRT-summated recall score ^f	33.6 (9.1)	33.4 (9.0)
Mean (SD) SRT-delayed recall score ^g	4.5 (2.4)	4.4 (2.5)
Mean (SD) BDRS score ^h	1.1 (0.7)	1.1 (0.8)

^aSample sizes varied for some measures due to missing data. For apolipoprotein genotyping, sample sizes were 669 for placebo and 652 for rofecoxib.

^bAuditory Verbal Learning Test. Range = 0-75; lower score indicates greater impairment.

^cMini Mental State Exam. Range = 0-30; lower score indicates greater impairment.

^dClinical Dementia Rating. Range = 0-18; higher score indicates greater impairment.

^eCognitive subscale of the AD Assessment Scale. Range = 0-70; higher score indicates greater impairment.

^fSelective Reminding Test (immediate recall component). Range = 0-72; lower score indicates greater impairment.

^gSelective Reminding Test (delayed recall component). Range = 0-12; lower score indicates greater impairment.

^hBlessed Dementia Rating Scale. Range = 0-17; higher score indicates greater impairment.

Table 2 Kaplan-Meier Estimated Proportions of Clinically Diagnosed AD at 4-Month Intervals

Month after treatment	Placebo		Rofecoxib	
	N ^a	Estimated proportion	N ^a	Estimated proportion
0	732	0.0	725	0.0
4	707	0.3	682	0.9
8	666	1.4	642	3.1
12	625	3.1	582	5.1
16	581	4.2	535	7.4
20	528	6.4	486	9.2
24	458	9.2	415	12.3
28	390	10.3	344	14.2
32	359	11.0	311	15.7
36	310	12.5	258	17.9
40	240	13.7	214	19.3
44	177	15.6	159	21.0
48	86	17.4	68	23.0

^aNumber of patients at risk (number who had not been censored nor had an event up to, but not including, that time point).

maintained through the last follow-up time point of 48 months as shown by the nonsignificant *p*-value for the test of proportionality of treatment-specific hazard rates (see above). Estimated annual diagnosis rates were 6.4% (95% CI: 5.3%, 7.7%) in the rofecoxib group and 4.5% (95% CI: 3.6%, 5.6%) in the placebo group. The above analysis included patients whether or not they were taking study medication. In the prespecified analysis looking at events that occurred on-drug (*N*=723 for rofecoxib and *N*=728 for placebo), there was no evidence for an increased hazard ratio (1.49 (95% CI: 1.08, 2.05), *p*=0.014) compared with the hazard ratio of 1.46 observed in the primary analysis, which included patients who had stopped taking study treatment.

To further explore the unexpected finding favoring placebo in the primary analysis, we conducted a *post hoc* analysis to adjust for factors that showed an effect, at a significance level of *p*<0.10, on progression to AD. Those factors correlated with greater likelihood of progression to AD were lower baseline MMSE score stratum (24–26), female gender, age >75, and prior ginkgo use. Factors associated with a decreased risk of progressing to AD were longer duration of concomitant NSAID use, and concomitant use of statins. In the analysis that adjusted for these factors, the statistical significance of the hazard ratio was reduced (1.31 (95% CI: 0.98, 1.75) *p*=0.065). Presence of the apolipoprotein ε4 allele was a risk factor but was not included in the model because information was missing for a substantial proportion of patients. Based on a significance level of *p*<0.10, family history of AD did not appear to be a risk factor in this study.

Because rofecoxib and other NSAIDs can cause small mean increases in blood pressure (Gertz et al, 2002; Schwartz et al, 2002), and there is some evidence to suggest

that increased blood pressure might be associated with an increased risk of dementia (Skoog, 1997; Birkenhager et al, 2001), we also performed two *post hoc* analyses to evaluate whether the rofecoxib:placebo risk ratio for diagnosis of AD increased as a function of increased blood pressure change. In the first analysis, change from baseline in mean arterial blood pressure (defined as (2 × diastolic blood pressure + systolic blood pressure)/3) at month 4 was calculated for each patient. The rofecoxib:placebo odds ratios for diagnosis of AD were then calculated for three categories of patients: those with no change or a decrease (odds ratio = 1.43), those with an increase ≤ 5 mmHg (odds ratio = 1.18), and those with an increase > 5 mmHg (odds ratio = 1.47). The Breslow-Day test of homogeneity of the odds ratios across categories indicated no significant differences (*p*=0.895). In the second *post hoc* analysis, we looked at a predefined limit of change in systolic blood pressure, which was prespecified as a postrandomization value that was ≥ 180 mmHg and showed a ≥ 20 mmHg increase from baseline. The rofecoxib:placebo hazard ratios for diagnosis of AD were similar in those patients who did not meet the predefined limit of change criteria (hazard ratio = 1.42 (95% CI: 1.06, 1.92)), compared with those who did meet the criteria (hazard ratio = 1.53 (95% CI: 0.49, 4.81)).

The least squares mean scores for the secondary end points of cognition and function at 1-year intervals using the repeated measures models approach described in Methods are summarized in Table 3. The prespecified analysis looked at the estimated annual slope difference (placebo minus rofecoxib) for each measure based on data over the entire 4-year period. In contrast to the primary end point, there were no significant differences between treatment groups on estimated annual slope differences for SRT-summed recall score (slope difference = 0.026 (95% CI: -0.307, 0.359), *p*=0.878), SRT-delayed recall score (slope difference = -0.012 (95% CI: -0.104, 0.081), *p*=0.806), MMSE score (slope difference = -0.002 (95% CI: -0.095, 0.090), *p*=0.959), ADAS-Cog score (slope difference = -0.098 (95% CI: -0.287, 0.091), *p*=0.311), or BDRS score (slope difference = 0.079 (95% CI: -0.081, 0.238), *p*=0.333). The slope difference for the CDR-sum of boxes score showed a nonsignificant trend in favor of placebo (slope difference = -0.068 (95% CI: -0.139, 0.002), *p*=0.058); this would be expected to be highly correlated with the primary end point since the global score on the CDR was used to trigger the diagnosis of AD. Additional sensitivity analyses for the secondary end points were also performed using a last-observation-carried forward approach. The results, indicating a lack of treatment differences, were similar across these analyses, which were based on different assumptions about the missing data structure.

The *post hoc* analysis comparing test scores in overlapping subgroups of patients with MMSE scores of 24–26 in the present study (the subgroup with the worst prognosis) and a previous AD treatment study (Reines et al, 2004) included a total of 405 patients (rofecoxib *N*=189, placebo *N*=216) from the present MCI study, and 205 patients (rofecoxib *N*=91, placebo *N*=114) from the previous AD treatment study. There was no consistent evidence of a differential treatment effect of rofecoxib in the

Rofecoxib in mild cognitive impairment

LJ Thal et al

Table 3 Secondary End Points: Least Squares Mean (Standard Error) Scores with Difference and 95% Confidence Interval

Measure	1-Year			2-Year			3-Year			4-Year		
	Placebo ^a (N ^c = 659)	Rofecoxib ^b (N ^c = 624)	Difference ^{a,b} (95% CI)	Placebo ^a (N ^c = 509)	Rofecoxib ^b (N ^c = 460)	Difference ^{a,b} (95% CI)	Placebo ^a (N ^c = 344)	Rofecoxib ^b (N ^c = 302)	Difference ^{a,b} (95% CI)	Placebo ^a (N ^c = 195)	Rofecoxib ^b (N ^c = 176)	Difference ^{a,b} (95% CI)
SRT-summed	35.7 (0.3)	35.6 (0.4)	-0.2 (-1.1,0.7)	36.5 (0.4)	35.7 (0.4)	-0.8 (-1.9,0.3)	36.1 (0.5)	35.7 (0.5)	-0.3 (-1.7,1.1)	36.0 (0.7)	35.6 (0.7)	-0.4 (-2.3,1.5)
SRT-delayed	4.8 (0.1)	4.7 (0.1)	-0.0 (-0.3,0.2)	4.9 (0.1)	4.8 (0.1)	-0.1 (-0.5,0.2)	4.9 (0.1)	4.7 (0.1)	-0.2 (-0.6,0.1)	4.5 (0.2)	4.5 (0.2)	0.1 (-0.5,0.6)
ADAS-Cog	8.9 (0.1)	9.3 (0.1)	0.3 (-0.0,0.7)	9.4 (0.2)	9.7 (0.2)	0.3 (-0.3,0.8)	9.8 (0.3)	10.1 (0.3)	0.3 (-0.4,1.0)	10.3 (0.3)	10.7 (0.3)	0.3 (-0.4,1.1)
MMSE	27.3 (0.1)	27.2 (0.1)	-0.1 (-0.3,0.1)	27.2 (0.1)	27.3 (0.1)	0.1 (-0.2,0.4)	27.1 (0.1)	27.1 (0.1)	-0.1 (-0.4,0.3)	26.7 (0.2)	26.5 (0.2)	-0.2 (-0.7,0.3)
CDR-sum of boxes	1.5 (0.0)	1.6 (0.0)	0.1 (-0.0,0.2)	1.8 (0.1)	2.0 (0.1)	0.2 (-0.0,0.4)	2.1 (0.1)	2.2 (0.1)	0.2 (-0.0,0.4)	2.2 (0.1)	2.4 (0.1)	0.2 (-0.0,0.5)
BDRS	—	—	—	1.4 (0.1)	1.4 (0.1)	0.0 (-0.1,0.2)	1.4 (0.1)	1.5 (0.1)	0.1 (-0.1,0.3)	1.4 (0.1)	1.6 (0.1)	0.1 (-0.1,0.4)

Results based on repeated measures model.

^aDifference is rofecoxib-placebo.^bSample size (number of patients with any secondary efficacy measure by time point).

See Table 1 for key to test abbreviations and interpretation of direction of scores. 1-year data are not provided for the BDRS because it was not scheduled to be administered until the 2-year time point.

overlapping subgroups in the two studies. The estimated annual slope difference for the ADAS-Cog score was -0.240 (95% CI: -0.736, 0.255), $p=0.340$ in the MMSE 24-26 subgroup from the present study, and 0.447 (95% CI: -1.284, 2.178), $p=0.611$ in the MMSE 24-26 subgroup from the previous AD treatment study. The estimated annual slope difference for the CDR-sum of boxes score was -0.180 (95% CI: -0.350, -0.009), $p=0.039$ in the MMSE 24-26 subgroup from the present study, and -0.053 (95% CI: -0.560, 0.454), $p=0.837$ in the MMSE 24-26 subgroup from the previous AD treatment study.

A total of 1451 patients were included in the on-drug safety analysis (723 in the rofecoxib group and 728 in the placebo group). There were slightly fewer patients in the safety analysis than in the efficacy analysis because six patients who never took study medication were excluded from the safety analysis. The adverse experience profile is summarized in Table 4. The rofecoxib and placebo groups were similar with regard to the percentages of patients with any adverse experience, any serious adverse experience, and who discontinued treatment due to an adverse experience. There was a significant increase for rofecoxib in the number of patients with adverse experiences that were considered possibly, probably, or definitely drug-related by the investigators. The most common drug-related adverse experiences are shown in Table 4. There was no particular individual adverse experience that contributed to the overall difference between rofecoxib and placebo for drug-related adverse experiences, and relatively few patients discontinued study treatment due to drug-related adverse experiences (58 or 8.0% for rofecoxib and 41 or 5.6% for placebo). The number of patients with confirmed upper gastrointestinal perforations, ulcers, or bleeds was 14 in the rofecoxib group and four in the placebo group. A total of 39 deaths occurred in patients who were taking study treatment or from fatal adverse events that started within 14 days of the last dose (24 or 3.3% for rofecoxib and 15 or 2.1% for placebo). Patients died from a range of causes that were consistent with expectations for an elderly population, and there was no specific pattern as to the cause of death in either treatment group. The only specific fatal adverse events with more than one patient per treatment group were myocardial infarction (four patients on rofecoxib and three on placebo), cardiac arrest (two patients on rofecoxib and none on placebo), pneumonia (two patients on rofecoxib and none on placebo), and renal failure (one patient on rofecoxib and two on placebo). (An individual patient may have had more than one adverse event associated with death.) Off-drug follow-up mortality data were available for less than half of the patients ($N=356$ for rofecoxib, $N=307$ for placebo); the median duration of off-drug follow-up in these patients was 29 weeks in the rofecoxib group and 20 weeks in the placebo group. There were an additional 22 deaths in the off-drug period (17 in patients assigned to rofecoxib and five in patients assigned to placebo); 12 of these (11 in the rofecoxib group and one in the placebo group) occurred more than 48 weeks after treatment discontinuation. Compared with the on-drug period, there were an additional five patients with off-drug myocardial infarction fatal adverse events (five rofecoxib, none placebo), and an additional two patients with each of cardiac arrest (two rofecoxib, none placebo), pneumonia

Table 4 Summary of Clinical Adverse Experiences

Number (%) of patients with	Placebo (N = 720)	Rofecoxib 25 mg (N = 723)
≥ 1 adverse experience	670 (92.0%)	651 (90.0%)
≥ 1 drug-related adverse experience ^a	173 (23.8%)	211 (29.2%)*
Serious adverse experience ^b	236 (32.4%)	217 (30.0%)
Discontinued treatment due to adverse experience	147 (20.2%)	156 (21.6%)
Most common drug related adverse experiences ^c		
Diarrhea	16 (2.2%)	11 (1.5%)
Dyspepsia	17 (2.3%)	24 (3.3%)
Nausea	15 (2.1%)	14 (1.9%)
Dizziness	14 (1.9%)	18 (2.5%)
Peripheral edema	11 (1.5%)	23 (3.2%)
Increased blood pressure	14 (1.9%)	16 (2.2%)
Hypertension	25 (3.4%)	29 (4.0%)

^aExperiences that were rated as possibly, probably, or definitely drug-related by the investigator.

^bDeath, cancer, any adverse event that was life threatening, resulted in persistent or significant disability, resulted in or prolonged an existing hospitalization, or any other important medical event.

^cIncidence > 2.0% in either treatment group.

*Significantly different from placebo, $p < 0.05$.

(two rofecoxib, none placebo), and renal failure (two rofecoxib, none placebo) off-drug fatal adverse events. The number of patients with confirmed serious thrombotic vascular events on-drug was similar in the two groups (38 in the rofecoxib group and 36 in the placebo group). There were a total of six patients with confirmed ischemic strokes and one patient with hemorrhagic stroke in the rofecoxib group compared to 13 patients with confirmed ischemic strokes and two patients with hemorrhagic strokes in the placebo group. Thirteen patients in the rofecoxib group had confirmed nonfatal myocardial infarctions vs 10 in the placebo group.

DISCUSSION

In this 4-year study of 1457 patients with MCI, there was no evidence that rofecoxib delayed a diagnosis of AD. A treatment difference in favor of placebo was observed on the primary end point of time to clinically diagnosed AD. This finding was not confirmed by secondary measures of cognition (ADAS-Cog, SRT, MMSE) or global function (BDRS, CDR-sum of boxes), which found no statistically significant or clinically meaningful differences between treatment groups. The possibility that rofecoxib might be inferior to placebo was also not supported by data from two large previous AD treatment studies, which included patients with an MMSE score up to 26 (thereby partly overlapping with patients in the present study who had an MMSE score ≥ 24), and found that rofecoxib had no significant effect on the progression of cognitive or functional decline (Aisen et al, 2003; Reines et al, 2004).

Given the unexpected nature of the primary finding, it was thought important to compare the present results with those from an independent database. We therefore conducted a *post hoc* analysis of test scores in subgroups of patients with MMSE scores of 24–26 in the present study (ie those patients who were most likely to receive a diagnosis of AD) and in a previous AD treatment study (Reines et al, 2004). These studies had similar designs and utilized similar assessments. Since MCI is hypothesized to be on a continuum of cognitive impairment ranging from very mild impairment (ie MCI) through mild, moderate, and severe dementia, it is plausible that many patients with MMSE scores of 24–26 in the two studies may have been biologically similar, even though one group was diagnosed with dementia and the other was not. There was no evidence to suggest differences between treatments on assessments of cognition (ADAS-Cog) in this subgroup in either study. In the subgroup analysis from the present study, there was a significant difference between treatments for the CDR-sum of boxes score. This finding was not surprising given that the global score on the CDR was the trigger for diagnosing AD. Indeed, because of its close relation with the primary end point of AD diagnosis, the CDR sum-of boxes score can be viewed as a surrogate for the primary end point. There was no difference between rofecoxib and placebo on the CDR sum-of-boxes score in the subgroup analysis from the previous AD treatment study.

An intriguing observation in the present study was that the separation between treatment groups in rates of diagnosis of AD was apparent from 4 months (the earliest time point assessed) but did not increase over time, raising the possibility that there may have been an imbalance between the groups at baseline. Although there was no clear evidence of a major imbalance for measured baseline variables, including severity of impairment, the model which adjusted for covariates that were risk factors for receiving a diagnosis of AD showed the smallest treatment effect and was not statistically significant. The finding that the treatment difference was not further increased in the analysis restricted to the on-drug population was also not supportive of a true treatment effect.

Since this is the first report of a completed randomized controlled study examining progression to an AD diagnosis in MCI patients, it is important to consider aspects of the design or conduct of the study that could have had an influence on the results. An obvious concern in a study with a long duration is that differential discontinuation rates may have influenced the results. A discontinuation rate of approximately 45% occurred over the course of the study and only 40% of patients completed the study on-drug. These findings are not surprising given the duration of the study and the elderly population being investigated. Overall discontinuation rates, time course of discontinuations, and change from baseline test scores at the time of discontinuation were similar between the treatment groups, although there were some small differences between the groups with regard to specific reasons for discontinuation; for example, a greater proportion of patients in the placebo group discontinued due to withdrawal of consent. The relatively high proportion of patients who withdrew consent may have been a consequence of the fact that patients were required to sign an additional consent form after 2 years,

owing to the study being extended beyond the original timeline, and were unwilling to go beyond their initial commitment. There was also some evidence that patients who discontinued from the placebo group were more impaired at baseline on the ADAS-Cog than patients who discontinued from the rofecoxib group. If the more impaired patients who discontinued were at higher risk of conversion to AD, then differential dropouts may have had an influence on the results.

Another notable observation in the present study was that the overall annual AD diagnosis rate of 5–6% was lower than the anticipated 10–15% annual diagnosis rate reported in previous observational or natural history studies (Petersen et al, 2001a,b). There are several possible explanations for this disparity. Firstly, the diagnosis of MCI is heavily dependent on clinical judgment (Petersen, 2003), and the particular tests and score thresholds used to provide objective evidence of memory impairment are not standardized (Chertkow, 2002; Petersen, 2003). It is therefore possible that the population of patients included in the study was more heterogeneous than, or different from, MCI populations previously described. Indeed, the population contained a lower proportion of women and patients with the apolipoprotein $\epsilon 4$ allele than others have reported (Petersen et al, 1999; Jack et al, 1999; Morris et al, 2001). Another possible contributing factor to the low diagnosis rates was the change in the cut-score criteria on the AVLT (the memory test used to provide objective evidence of memory impairment) during the study. After 6 months of enrollment, the cutoff was changed from age-adjusted 1.5 SD cut scores to a single cut score 1 SD below the mean for normal elderly subjects (see Appendix A1). We retrospectively looked at diagnosis rates in patients who met the original 1.5 SD criteria ($N=203$ for rofecoxib and $N=203$ for placebo) and the estimated annual diagnosis rates (11.3% for rofecoxib and 8.3% for placebo) were more in line with previous data, although still lower than anticipated for the placebo group. Finally, the fact that the MCI concept was relatively new at the time the study was initiated, along with the source of the patients (advertising campaigns as well as direct clinical referrals), may have also resulted in greater heterogeneity with regard to underlying etiology in the recruited patients.

To summarize the efficacy data, the unexpected primary finding in the present study suggesting that rofecoxib might accelerate the rate of AD diagnosis in patients with MCI was not supported by data on secondary measures in the same study, nor by other data from previous studies that assessed cognitive and functional decline in AD patients (Aisen et al, 2003; Reines et al, 2004). These observations suggest that the finding may not be indicative of a true treatment effect. If not a true effect, then the most likely explanations are a chance occurrence, or an imbalance between the treatment groups that existed at baseline or arose during the study due to differential discontinuations.

We cannot exclude the possibility that rofecoxib might accelerate conversion of MCI patients to AD, although this possibility would be at variance with the prior epidemiological and clinical data cited above, as well as the body of experimental evidence suggesting that COX-2 inhibition might attenuate neuronal death in several disease states including Parkinson's disease, multiple sclerosis, and

amyotrophic lateral sclerosis, in addition to AD (Andreasson et al, 2001; Jain et al, 2002; Xiang et al, 2002; Giovannini et al, 2003; Scali et al, 2003; Teismann et al, 2003; Qin et al, 2003; Consilvio et al, 2004; Rose et al, 2004). The early onset of the treatment difference (and the lack of progressive widening of the difference over the course of the trial) further argues against a direct effect of rofecoxib on the underlying pathophysiology in AD. It could be speculated that the apparent increased conversion to AD might be secondary to a non-AD-specific aspect of rofecoxib's biological effects. For example, rofecoxib and other NSAIDs can cause small mean increases in blood pressure (Gertz et al, 2002; Schwartz et al, 2002). It has been suggested that increased blood pressure might be associated with an increased risk of dementia, although the evidence is inconsistent (Skoog, 1997; Birkenhager et al, 2001). We investigated this possibility by performing *post hoc* analyses to determine if the rofecoxib:placebo risk ratio for diagnosis of AD increased as a function of increased blood pressure change. There was no evidence for such a relationship. A proposed mechanism by which increased blood pressure could lead to dementia is due to an increased risk for cardiovascular adverse events such as strokes or infarcts. As noted below, there was no evidence for an increase in these types of events in patients taking rofecoxib. Furthermore, CT or MRI brain scans were performed in all patients diagnosed with dementia, to exclude the possibility of vascular dementia.

In addition to evaluating efficacy, the present study provided important placebo-controlled data on the safety of rofecoxib 25 mg over periods of up to 4 years in an elderly population. The median duration of exposure to study medication was approximately 2 years. The mean age of patients was 75 years and approximately 50% were at least 75 years old. Previous safety data on rofecoxib come largely from osteoarthritis studies looking at a relatively younger population with a mean age around 60 years (Langman et al, 1999; Reicin et al, 2002). Rofecoxib was generally well tolerated by the elderly patients in the study, consistent with results from prior clinical studies in osteoarthritis (Langman et al, 1999; Reicin et al, 2002) and AD (Reines et al, 2004). The overall incidence of adverse experiences, serious adverse experiences, and discontinuations due to adverse experiences were similar or only slightly increased for rofecoxib vs placebo. Not surprisingly, there was an increase in the number of adverse experiences thought to be drug-related by the investigators for rofecoxib vs placebo, but relatively few patients discontinued treatment due to these adverse experiences. The increase was mainly due to small increases in adverse experiences known to be associated with NSAID use and previously reported for rofecoxib such as dyspepsia, hypertension, and peripheral edema. Despite the high mean age of the study population, few patients had confirmed upper gastrointestinal perforations, ulcers, or bleeds, although there were numerically more in the rofecoxib group than in the placebo group. Elderly patients are at increased risk for serious vascular events. Rofecoxib did not appear to increase the risk in this study, since the number of confirmed serious thrombotic vascular events was similar in each treatment group. The procedure used for determining confirmed cardiovascular events, such as heart attack and stroke, was the same as for the recent

3-year APPROVe (Adenomatous Polyp Prevention on Vioxx) study in 2586 patients (62% male, mean age 59 years) with a history of colorectal adenomas, which found an approximately two-fold increased relative risk for these events in patients taking rofecoxib vs placebo, beginning after 18 months of treatment (Bresalier *et al*, 2004). The reason for the discrepancy between the present findings and those from APPROVe is unclear. No striking differences were noted between the treatment groups with regard to other, nonserious, vascular adverse events in the present study. The total number of deaths and causes of death on-drug were consistent with expectations for an elderly population, and there was no specific pattern as to the cause of death in either treatment group. The off-drug mortality data are difficult to assess since follow-up information was available for less than half the patients, there was an imbalance in the extent of follow-up data between the two groups, and the majority of deaths occurred more than 48 weeks after treatment had been discontinued.

In conclusion, the finding that rofecoxib did not delay a diagnosis of AD in the present study, or slow the progression of AD in previous studies (Aisen *et al*, 2003; Reines *et al*, 2004), suggests that inhibition of COX-2 is not a useful therapeutic approach in AD. It is possible that nonselective NSAIDs may still be found to have beneficial effects in treating or delaying the progression of symptoms in AD, due to factors other than COX-2 inhibition. However, the only long-term results from an adequately designed study with a nonselective NSAID (naproxen) available to date are not encouraging (Aisen *et al*, 2003). Trials evaluating alternative, non-anti-inflammatory, approaches to delaying a diagnosis of AD in patients with MCI are underway (Petersen, 2003).

ACKNOWLEDGEMENTS

This study was funded by Merck Research Laboratories. SH Ferris was a paid consultant for Merck Research Laboratories on this study. L Kirby received funding from Merck Research Laboratories to participate in this study. GA Block, CR Lines, E Yuen, C Assaid, ML Nessler, BA Norman, CC Baranak, and SA Reines were employees of Merck Research Laboratories at the time the study was performed (GA Block is currently an employee of Astra Zeneca, and SA Reines and E Yuen are currently employees of Johnson & Johnson). These data were previously presented at the 42nd annual meeting of the American College of Neuropsychopharmacology, Puerto Rico, December 7–11, 2003.

Participating investigators in the Rofecoxib Protocol 078 study group were as follows: Piero Antuono, Froedtert Lutheran Memorial Hospital, Milwaukee, WI; Barry Baumel, Baumel-Eisner Neuromedical Institute, Miami, FL; Karen Bell, Columbia University, New York, NY; B Ian Blatt, Southeastern Pennsylvania Medical Institute, Havertown, PA; Martin Conway, Lovelace Scientific Resources Inc., Albuquerque, NM; Howard S Cummins, The Institute for Advanced Clinical Research, Elkins Park, PA; Mary J Derbenwick, Atlantic Institute of Clinical Research, Daytona Beach, FL; P Murali Doraiswamy, Duke University Medical Center, Durham, NC; Eugene DuBoff, Summit Research Network Inc., Denver, CO; Keith Edwards, Bennington, VT;

Larry Eisner, Baumel-Eisner Neuromedical Institute, Jay Ellis, Neuroscience Research, Pittsfield, MA; Donald England, Radiant Research, Eugene, OR; James Ferguson, Pharmacological Research Corp, Salt Lake City, UT; Jerome Goldstein, San Francisco Headache Clinic, San Francisco, CA; James Hartford, Summit Research Network, Cincinnati, OH; Jon Heiser, Pharmacology Research Institute, Northridge, CA; Daniel Hier, University of Illinois at Chicago, Chicago, IL; Bruce Hodges, Midwest Center for Clinical Studies, Kansas City, MO; Richard Holub, Neurological Associates of Albany, Albany, NY; Lawrence R Jenkyn, Dartmouth-Hitchcock Medical Center, Lebanon, NH; F Cleveland Kinney, University of Alabama at Birmingham, Birmingham, AL; Louis Kirby, Pivotal Research Centers, Peoria, AZ; Peter D Lundborg, Seattle, WA; Elliott Mancall, Headache Center, Philadelphia, PA; David Margolin, Margolin Brain Institute, Fresno, CA; James McCarthy, Comprehensive Neuroscience Inc., South Yarmouth, MA; William McEntee, Comprehensive Neuroscience Inc., Sarasota, FL; Margarita Nunez, Comprehensive Neuroscience Inc., St Petersburg, FL; Jorg Pahl, Pahl Brain Associates PC, Oklahoma City, OK; Anton Porsteinsson, University of Rochester, Rochester, NY; Eric Pfeiffer, USF Suncoast Gerontology Center, Tampa, FL; Norman Relkin, New York Presbyterian Hospital, New York, NY; Joel S Ross, Long Branch, NJ; Barry Rovner, Philadelphia, PA; Ward Tolbert Smith, Summit Research, Portland, OR; John A Stoukides, Comprehensive Neuroscience Inc., East Providence, RI; Joseph Tate, DeKalb-Gwinnett, OB/GYN, PC, Norcross, GA; Stephen Thein Jr, Pacific Research Network, San Diego, CA; Harvey Tilker, Four Rivers Clinical Research Center, Paducah, KY; Jerry R Tindel, Radiant Research, Austin, TX; Jack Tomlinson, Alzheimer's Diagnostic Clinic, Wichita Falls, TX; Richard Weisler, Raleigh, NC; Jeanette Wendt, Neurological Associates of Tucson, Tucson, AZ; Kerri Wilks, Innovative Medical Research, Towson, MD; Parvaneh Zolnouri, California Clinical Trials, Beverly Hills, CA.

Members of the Rofecoxib Protocol 078 Endpoint Adjudication Committee were as follows: Steven Ferris, New York University School of Medicine, New York, NY; Mary Sano, Mount Sinai School of Medicine, Bronx Veterans Medical Research Center, New York, NY; Peter Whitehouse, University Hospitals of Cleveland, Cleveland, OH.

REFERENCES

- Aisen PS, Schafer KA, Grundman M, Pfeiffer E, Sano M, Davis KL *et al* (2003). Effects of rofecoxib or naproxen vs placebo on Alzheimer's disease progression: a randomized controlled trial. *JAMA* 289: 2819–2826.
- Andreasson KI, Savonenko A, Videnky S, Goellner JJ, Zhang Y, Shaffer A *et al* (2001). Age-dependent cognitive deficits and neuronal apoptosis in cyclooxygenase-2 transgenic mice. *J Neurosci* 21: 8198–8209.
- Birkenhager WH, Forette F, Scux ML, Wang JG, Staessen JA (2001). Blood pressure, cognitive functions, and prevention of dementia in older patients with hypertension. *Arch Intern Med* 161: 152–156.
- Bombardier C, Laine L, Reicin A, Shapiro D, Burgos-Vargas R, Davis B *et al* (2000). Comparison of upper gastrointestinal toxicity of rofecoxib and naproxen in patients with rheumatoid arthritis. *N Engl J Med* 343: 1520–1528.

- Bresalier RS, Sandler R, Riddell R, Lanus A, Morton D, Reicin A et al (2004). Preliminary Cardiovascular Safety Data from the APPROVe Study, Presented at the annual meeting of the American College of Rheumatology, San Antonio, Texas, October 18. Abstract available at: http://www.rheumatology.org/annual/press/APPROVesession_announce.asp.
- Buschke H (1973). Selective reminding for analysis of memory and learning. *J Verbal Learn Verbal Behav* 12: 543-550.
- Cagnin A, Brooks DJ, Kennedy AM, Gunn RN, Myers R, Turkheimer FE et al (2001). In-vivo measurement of activated microglia in dementia. *Lancet* 358: 461-467.
- Chertkow H (2002). Mild cognitive impairment. *Curr Opin Neurol* 15: 401-407.
- Consilvio C, Vincent AM, Feldman EL (2004). Neuroinflammation, COX-2, and ALS—a dual role? *Exp Neurol* 187: 1-10.
- Eminan M, Gill S, Samii A (2003). Effect of non-steroidal anti-inflammatory drugs on risk of Alzheimer's disease: systematic review and meta-analysis of observational studies. *BMJ* 327: 128.
- Folstein MF, Folstein SE, McHugh PR (1975). 'Mini-Mental State': a practical method for grading the cognitive state of patients for the clinician. *J Psychiatric Res* 12: 189-198.
- Gertz BJ, Krupa D, Bolognese JA, Sperling RS, Reicin A (2002). A comparison of adverse renovascular experiences among osteoarthritis patients treated with rofecoxib and comparator non-selective non-steroidal anti-inflammatory agents. *Curr Med Res Opin* 18: 82-91.
- Giovannini MG, Scali C, Prosperi C, Belluci A, Pepeu G, Casamenti F (2003). Experimental brain inflammation and neurodegeneration as model of Alzheimer's disease: protective effects of selective COX-2 inhibitors. *Int J Immunopathol Pharmacol* 16(Suppl 2): 31-40.
- Hamilton M (1960). A rating scale for depression. *J Neurol Neurosurg Psychiatry* 23: 56-62.
- Ho L, Purhoit D, Haroutunian V, Luteran JD, Willis F, Naslund J et al (2001). Neuronal cyclooxygenase 2 expression in the hippocampal formation as a function of the clinical progression of Alzheimer disease. *Arch Neurol* 58: 487-492.
- in 't Veld BA, Launer LJ, Breteler MMB, Hofman A, Stricker BH Ch (2002). Pharmacological agents associated with a preventive effect on Alzheimer's disease: a review of the epidemiologic data. *Epidemiol Rev* 24: 248-268.
- in 't Veld BA, Ruitenberg A, Hofman A, Launer LJ, van Duijn CK, Stijnen T et al (2001). Nonsteroidal antiinflammatory drugs and the risk of Alzheimer's disease. *N Engl J Med* 345: 1515-1521.
- Ivnik RJ, Malec JF, Tangalos EG, Petersen RC, Kokmen E, Kurland LT (1990). The auditory-verbal learning test (AVLT): norms for ages 55 years and older. *Psychol Assess: J Consult Clin Psychol* 2: 304-312.
- Jack Jr CR, Petersen RC, Xu YC, O'Brien PC, Smith GE, Ivnik RJ et al (1999). Prediction of AD with MRI-based hippocampal volume in mild cognitive impairment. *Neurology* 52: 1397-1403.
- Jain NK, Patil CS, Kulkarni SK, Singh A (2002). Modulatory role of cyclooxygenase inhibitors in aging- and scopolamine or lipopolysaccharide-induced cognitive dysfunction in mice. *Behav Brain Res* 133: 369-376.
- Konstam MA, Weir MR, Reicin A, Shapiro D, Sperling RS, Barr E et al (2001). Cardiovascular thrombotic events in controlled, clinical trials of rofecoxib. *Circulation* 104: 2280-2288.
- Langman MJ, Jensen DM, Watson DJ, Harper SE, Zhao PL, Quan H et al (1999). Adverse upper gastrointestinal effects of rofecoxib compared with NSAIDs. *JAMA* 282: 1929-1933.
- Lines CR, McCarroll KA, Lipton RB, Block GA (2003). Telephone screening for amnesic mild cognitive impairment. *Neurology* 60: 261-266.
- McGeer PL, Schulzer M, McGeer EG (1996). Arthritis and anti-inflammatory agents as possible neuroprotective factors for Alzheimer's disease: a review of 17 epidemiological studies. *Neurology* 47: 425-432.
- McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM (1984). Clinical diagnosis of Alzheimer's disease: report of the NINCDS/ADRDA work group under the auspices of the Department of Health and Human Services task force on Alzheimer's disease. *Neurology* 34: 939-944.
- Merck & Co. Inc. (2003). VIOXX[®] (rofecoxib tablets and oral suspension). In: *Physicians' Desk Reference*[®]. Thomson PDR: Montvale, NJ. pp 2120-2125.
- Morris JC (1993). The clinical dementia rating (CDR): current version and scoring rules. *Neurology* 43: 2412-2414.
- Morris JC, Mohs RC, Rogers H, Fillenbaum G, Heyman A (1988). Consortium to establish a registry for Alzheimer's disease (CERAD). Clinical and neuropsychological assessment of Alzheimer's disease. *Psychopharmacol Bull* 24: 641-652.
- Morris JC, Storandt M, Miller JP, McKeel DW, Price JL, Rubin BH et al (2001). Mild cognitive impairment represents early-stage Alzheimer disease. *Arch Neurol* 58: 397-405.
- Pasinetti GM (2001). Cyclooxygenase and inflammation in the clinical progression of Alzheimer's disease. In: Vane JR, Botting RM (eds). *Therapeutic Roles of Selective COX-2 Inhibitors*. William Harvey Press: London. pp 191-205.
- Petersen RC (2003). Mild cognitive impairment clinical trials. *Nat Rev: Drug Discov* 2: 646-653.
- Petersen RC, Doody R, Kurz A, Mohs RC, Morris JC, Rabins PV et al (2001a). Current concepts in mild cognitive impairment. *Arch Neurol* 58: 1985-1992.
- Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E (1999). Mild cognitive impairment: clinical characterization and outcome. *Arch Neurol* 56: 303-308.
- Petersen RC, Stevens JC, Ganguli M, Tangalos EG, Cummings JL, DeKosky ST (2001b). Practice parameter: early detection of dementia: mild cognitive impairment (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 56: 1133-1142.
- Qin W, Ho L, Pompl PN, Peng Y, Zhao Z, Xiang Z et al (2003). Cyclooxygenase (COX)-2 and COX-1 potentiate β -amyloid peptide generation through mechanisms that involve γ -secretase activity. *J Biol Chem* 278: 50970-50977.
- Reicin AS, Shapiro D, Sperling R, Barr E, Yu Q (2002). Comparison of cardiovascular thrombotic events in patients with osteoarthritis treated with rofecoxib versus nonselective nonsteroidal anti-inflammatory drugs (ibuprofen, diclofenac, and nabumetone). *Am J Cardiol* 89: 204-209.
- Reines SA, Block GA, Morris JC, Liu G, Nessly ML, Lines CR et al (2004). Rofecoxib: no effect on Alzheimer's disease in a 1-year, randomized, blinded, controlled study. *Neurology* 62: 66-71.
- Rogers J, Kirby LC, Hempelman SR, Berry DL, McGeer PL, Kaszniak AW et al (1993). Clinical trial of indomethacin in Alzheimer's disease. *Neurology* 43: 1609-1611.
- Rose JW, Hill KE, Watt HE, Carlson NG (2004). Inflammatory cell expression of cyclooxygenase-2 in the multiple sclerosis lesion. *J Neuroimmunol* 149: 40-49.
- Rosen WG, Mohs RC, Davis KL (1984). A new rating scale for Alzheimer's disease. *Am J Psychiatry* 141: 1356-1364.
- Rosen WG, Terry RD, Fuld PA, Katzman R, Peck A (1980). Pathological verification of ischemic score in differentiation of dementias. *Ann Neurol* 7: 486-488.
- Sainati S, Ingram D, Talwalker S, Geiss G (2000). Results of a double-blind, randomized, placebo-controlled study of celecoxib in the treatment of the progression of Alzheimer's disease. Sixth International Stockholm/Springfield Symposium on Advances in Alzheimer Therapy, 180 [Abstract].
- Scali C, Giovannini MG, Prosperi C, Belucci A, Pepeu G, Casamenti F (2003). The selective cyclooxygenase-2 inhibitor rofecoxib suppresses brain inflammation and protects cholinergic neurons from excitotoxic degeneration in vivo. *Neuroscience* 117: 909-919.

- Scharf, Mander E, Ugoni A, Vajda F, Christophidis N (1999). A double-blind, placebo-controlled trial of diclofenac/misoprostol in Alzheimer's disease. *Neurology* 53: 197-201.
- Schmidt M (1996). *Key Auditory Verbal Learning Test. A Handbook*. Western Psychological Services: Los Angeles.
- Schwartz JI, Vandormael K, Malice MP, Kalyani RN, Lasserer KC, Holmes GB et al (2002). Comparison of rofecoxib, celecoxib, and naproxen on renal function in elderly subjects receiving a normal-salt diet. *Clin Pharmacol Ther* 72: 50-61.
- Skoog I (1997). The relationship between blood pressure and dementia: a review. *Biomed Pharmacother* 51: 367-375.
- Teismann P, Vila M, Choi DK, Tieu K, Wu DC, Jackson-Lewis V et al (2003). COX-2 and neurodegeneration in Parkinson's disease. *Ann NY Acad Sci* 991: 272-277.
- Van Gool WA, Weinstein HC, Scheltens P, Walstra GJ, Scheltens PK (2001). Effect of hydroxychloroquine on progression of dementia in early Alzheimer's disease: an 18-month randomised, double-blind, placebo-controlled study. *Lancet* 358: 455-460.
- Warner TD, Giuliano F, Vojnovic I, Bukasa A, Mitchell JA, Vane JR (1999). Nonsteroid drug selectivities for cyclo-oxygenase-1 rather than cyclo-oxygenase-2 are associated with human gastrointestinal toxicity: a full *in vitro* analysis. *Proc Natl Acad Sci USA* 96: 7563-7568.
- Xiang Z, Ho L, Yemul S, Zhao Z, Qin W, Pompl P et al (2002). Cyclooxygenase-2 promotes amyloid plaque deposition in a mouse model of Alzheimer's disease neuropathology. *Gene Expr* 10: 271-278.

APPENDIX A1

Details of Tests Used

MMSE (Folstein et al, 1975): This is a brief test of overall cognitive function (range of scores = 0-30, lower score indicates greater impairment). Patients who scored <24 were excluded because of the possibility that they might have dementia.

AVLT (Schmidt, 1996): This is a 15-item, five-trial, word recall test (range of scores = 0-75, lower score indicates greater impairment). The interference trial and delayed recall trial, which usually follow the initial five-trial learning phase, were not administered. Patients had to score ≤ 37 words recalled correctly over five trials as objective evidence of memory impairment. This corresponds to a score ≤ 1 SD below the mean for normal elderly subjects aged 65-69 (Ivnik et al, 1990). For the first 6 months of study enrollment, age-adjusted cut scores 1.5 SDs below the means for separate age bands were used (eg there was a cut

score for those aged 67-69, another cut score for those aged 70-72, etc), but the criteria were modified after 6 months due to concerns that the original criteria may have been too strict and were conflicting with investigator's clinical judgment (see Petersen, 2003), resulting in low enrollment.

CDR (Morris, 1993): This is a global dementia rating scale, based on a semistructured interview with the patient and an informant. Patients are rated on dementia severity (0 = none, 0.5 = questionable dementia, 1 = mild dementia, 2 = moderate dementia, 3 = severe dementia) in six domains or 'boxes' (memory, orientation, judgment and problem solving, community affairs, home and hobbies, personal care). The global CDR score (range = 0-3, higher score indicates greater impairment) was determined according to a computerized version of the published scoring algorithm (<http://www.biostat.wustl.edu/~adrc/cdrpgm/index.html>). The sum of scores on the individual boxes was also calculated (range = 0-18, higher score indicates greater impairment). Patients had to have a global CDR score of 0.5 (questionable dementia) with a score of at least 0.5 in the memory domain to qualify for study entry.

ADAS-Cog (Rosen et al, 1984): This consists of 11 performance-based subtests of cognition, including measures of memory and praxis. The error scores on each subtest are added to give the total score (range = 0-70, higher score indicates greater impairment).

SRT (Buschke, 1973): This is a 12-item, six-trial, word list recall test with selective reminding of words not correctly recalled on the immediately preceding trial. Following the sixth trial, there is a 5 min delay, after which the patients are again asked to recall the list of 12 words. Two scores were derived—the sum of the number of words recalled over each of the six selective reminding trials (SRT-summed recall; range = 0-72, lower score indicates greater impairment) and the number of words recalled after the 5 min delay (SRT-delayed recall; range = 0-12, lower score indicates greater impairment).

BDRS (Morris et al, 1988): The version from the CERAD battery was used. This is an informant-based rating of a patient's ability to perform activities of daily living such as household tasks (Part 1) and self-care (Part 2) (range of scores = 0-17, higher score indicates greater impairment). Patients who scored >3.5 with any Part 1 item score >0.5 (indicating severe impairment) were excluded because of the possibility of dementia.

THE WILLIAM AND SYLVIA SILVERSTEIN
AGING AND DEMENTIA RESEARCH CENTER
Department of Psychiatry (THN 312B)
New York University Medical Center
550 First Avenue
New York, NY 10016
USA

FAX TRANSMITTAL FORM

Our FAX number: (212) 263-6991
E-mail address: BARRY.REISBERG@MED.NYU.EDU
Telephone number: (212) 263-8550

FROM: BARRY REISBERG, M.D.

To: RONNIE MALLON

FAX number: 201-343-1684

Telephone number: 201-487-5800 Ext 118

Country: USA

Number of pages (including this cover sheet): 02

Comments:

Date: 10/11/05

Time: 5:05 PM

- Scharf, Mander E, Ugoni A, Vajda F, Christophidis N (1999). A double-blind, placebo-controlled trial of diclofenac/misoprostol in Alzheimer's disease. *Neurology* 53: 197-201.
- Schmidt M (1996). *Rey Auditory Verbal Learning Test. A Handbook*. Western Psychological Services: Los Angeles.
- Schwartz JL, Vandormael K, Malice MP, Kalyani RN, Lasseter KC, Holmes GB et al (2002). Comparison of rofecoxib, celecoxib, and naproxen on renal function in elderly subjects receiving a normal-salt diet. *Clin Pharmacol Ther* 72: 50-61.
- Skoog I (1997). The relationship between blood pressure and dementia: a review. *Biomed Pharmacother* 51: 367-375.
- Teismann P, Vila M, Choi DK, Tieu K, Wu DC, Jackson-Lewis V et al (2003). COX-2 and neurodegeneration in Parkinson's disease. *Ann NY Acad Sci* 991: 272-277.
- Van Gool WA, Weinstein HC, Scheltens P, Walstra GJ, Scheltens PK (2001). Effect of hydroxychloroquine on progression of dementia in early Alzheimer's disease: an 18-month randomised, double-blind, placebo-controlled study. *Lancet* 358: 455-460.
- Warner TD, Giuliano F, Vojnovic I, Bukasa A, Mitchell JA, Vane JR (1999). Nonsteroid drug selectivities for cyclo-oxygenase-1 rather than cyclo-oxygenase-2 are associated with human gastrointestinal toxicity: a full *in vitro* analysis. *Proc Natl Acad Sci USA* 96: 7563-7568.
- Xiang Z, Ho L, Yemul S, Zhao Z, Qin W, Pompl P et al (2002). Cyclooxygenase-2 promotes amyloid plaque deposition in a mouse model of Alzheimer's disease neuropathology. *Gene Expr* 10: 271-278.

APPENDIX A1

Details of Tests Used

MMSE (Folstein et al, 1975): This is a brief test of overall cognitive function (range of scores = 0-30, lower score indicates greater impairment). Patients who scored <24 were excluded because of the possibility that they might have dementia.

AVLT (Schmidt, 1996): This is a 15-item, five-trial, word recall test (range of scores = 0-75, lower score indicates greater impairment). The interference trial and delayed recall trial, which usually follow the initial five-trial learning phase, were not administered. Patients had to score ≤ 37 words recalled correctly over five trials as objective evidence of memory impairment. This corresponds to a score ≤ 1 SD below the mean for normal elderly subjects aged 65-69 (Ivnik et al, 1990). For the first 6 months of study enrollment, age-adjusted cut scores 1.5 SDs below the means for separate age bands were used (eg there was a cut

score for those aged 67-69, another cut score for those aged 70-72, etc), but the criteria were modified after 6 months due to concerns that the original criteria may have been too strict and were conflicting with investigator's clinical judgment (see Petersen, 2003), resulting in low enrollment.

CDR (Morris, 1993): This is a global dementia rating scale, based on a semistructured interview with the patient and an informant. Patients are rated on dementia severity (0 = none, 0.5 = questionable dementia, 1 = mild dementia, 2 = moderate dementia, 3 = severe dementia) in six domains or 'boxes' (memory, orientation, judgment and problem solving, community affairs, home and hobbies, personal care). The global CDR score (range = 0-3, higher score indicates greater impairment) was determined according to a computerized version of the published scoring algorithm (<http://www.biostat.wustl.edu/~adrc/cdrpgm/index.html>). The sum of scores on the individual boxes was also calculated (range = 0-18, higher score indicates greater impairment). Patients had to have a global CDR score of 0.5 (questionable dementia) with a score of at least 0.5 in the memory domain to qualify for study entry.

ADAS-Cog (Rosen et al, 1984): This consists of 11 performance-based subtests of cognition, including measures of memory and praxis. The error scores on each subtest are added to give the total score (range = 0-70, higher score indicates greater impairment).

SRT (Buschke, 1973): This is a 12-item, six-trial, word list recall test with selective reminding of words not correctly recalled on the immediately preceding trial. Following the sixth trial, there is a 5 min delay, after which the patients are again asked to recall the list of 12 words. Two scores were derived--the sum of the number of words recalled over each of the six selective reminding trials (SRT-summed recall; range = 0-72, lower score indicates greater impairment) and the number of words recalled after the 5 min delay (SRT-delayed recall; range = 0-12, lower score indicates greater impairment).

BDRS (Morris et al, 1988): The version from the CERAD battery was used. This is an informant-based rating of a patient's ability to perform activities of daily living such as household tasks (Part 1) and self-care (Part 2) (range of scores = 0-17, higher score indicates greater impairment). Patients who scored >3.5 with any Part 1 item score >0.5 (indicating severe impairment) were excluded because of the possibility of dementia.

Subjective Memory Deterioration and Future Dementia in People Aged 65 and Older

Li Wang, MS,* Gerald van Belle, PhD,^{†‡} Paul K. Crane, MD, MPH,* Walter A. Kukull, PhD,[§] James D. Bowen, MD,^{||} Wayne C. McCormick, MD, MPH,* and Eric B. Larson, MD, MPH*[¶]

OBJECTIVES: To study whether subjective memory deterioration is associated with future dementia in older people.

DESIGN: A population-based prospective cohort study begun in 1994 with biennial follow-up interviews.

SETTING: Community-based members of Group Health Cooperative, a large health maintenance organization in the Seattle area.

PARTICIPANTS: A sample of 1,883 subjects, dementia free, aged 65 and older, who scored 91 or higher on the 100-point Cognitive Ability Screening Instrument (CASI) at study entry.

MEASUREMENTS: Subjective memory was assessed by asking whether memory had changed on 5-point Likert scales (e.g., 1 = definitely improved, 3 = no change, 5 = definitely deteriorated) with regard to five items: remembering names, faces, friends, and appointments and judging the time. The items were summed for a possible total score ranging from 5 to 25. Subjective memory deterioration was defined as present if the total score was 20 or above. Cognitive performance was measured using the CASI. Incident dementia cases were identified using *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition* criteria.

RESULTS: Of 1,883 subjects, 126 developed dementia during 5 years of follow-up. Subjective memory deterioration was associated with cognitive decline and incident dementia. Age modified the association between subjective memory deterioration and future dementia. For persons reporting subjective memory deterioration at the ages of 70, 75, and 80, the hazard ratios of developing dementia were 6.0 (95% confidence interval (CI) = 2.1–18), 3.2 (95% CI = 1.6–6.2) and 1.6 (95% CI = 0.86–3.1), respectively.

CONCLUSION: Subjective memory deterioration was found to precede the development of dementia in older people with normal cognitive screening results. These findings suggest that a high level of subjective memory deterioration in persons with normal objective cognitive function may identify a subset of individuals at greater risk for developing dementia. *J Am Geriatr Soc* 52:2045–2051, 2004.

Key words: subjective memory deterioration; incident dementia

The introduction of interventions to slow down the development of Alzheimer's disease (AD) and other types of dementia stresses the need to detect dementia at an early stage. Therefore, it is important to search for early signs of dementia. Many older people report subjective memory decline or so called memory complaints.^{1–4} Self-perceived memory deterioration may be a harbinger of future dementia in older people.

Development of dementia, particularly AD, often follows a gradual course that may take years before clinical dementia can be observed. During its course, people might perceive their own cognitive decline, manifested as memory deterioration, at an early period when they still appear normal, but it is unknown whether, and to what extent, self-perceived decline is a perception of the normal aging process or might represent an early sign of dementia.

Several population-based longitudinal studies found that subjective memory complaints were associated with cognitive decline or dementia,^{4–10} but other studies found no association.^{11,12} Most of these studies included individuals with evidence of objective cognitive impairment and therefore were unable to address the question of whether subjective memory complaints were a sign of future dementia at an early stage when objective impairment was not evident. Few studies have investigated the temporal relationship between subjective memory complaints and future dementia in nondemented older people who had no objective evidence of cognitive impairment on a screening test. Only two reports, both from the Amsterdam Study of the

From the Departments of *Medicine, †Biostatistics, ‡Environmental Health, §Epidemiology, and ||Neurology, University of Washington, Seattle, Washington; and ¶Group Health Cooperative Center for Health Studies, Seattle, Washington.

The research on which this article is based was performed pursuant to cooperative agreement U01 AG06781 with the National Institute on Aging (NIA), National Institutes of Health (NIH). Drs. van Belle and Kukull also acknowledge support of U01 AG16976 with the NIA, NIH. Dr. Crane was supported by Grant K08 AG 022232–01 from the NIA.

Address correspondence to Gerald van Belle, Box 357232, University of Washington, Seattle WA 98195. E-mail: vanbelle@u.washington.edu

Elderly (ASE), found that memory complaints were associated with incident AD⁷ and accelerated cognitive decline⁹ in older people in whom cognitive impairment was not yet apparent. In the ASE, memory complaints were assessed using a general question regarding whether a person had complaints about memory. The question assessed the prevalence of memory problems, but people could complain about their memory for reasons that were not necessarily related to worsening in memory.^{13,14} By contrast, the current study focused on perceived change in memory as assessed using five specific questions about whether a person's memory had changed, because perceived worsening in memory might be an early sign in the development of dementia with insidious onset like AD. It was hypothesized that a high level of self-perceived memory decline is associated with future dementia in older people whose cognition was in the normal range at baseline.

METHODS

Subjects

The subjects were selected from the Adult Changes in Thought (ACT) study. The study design of the ACT has been described in more detail elsewhere.^{15,16} In brief, the study population was sampled from Group Health Cooperative members aged 65 and older in the Seattle area from 1994 to 1996. Subjects received the Cognitive Ability Screening Instrument (CASI)¹⁷ as the initial cognitive screen and were interviewed with structured questionnaires to obtain data on demographics, medical history, memory and general functioning, and potential epidemiological risk factors. The CASI provides quantitative assessment of attention, concentration, orientation, short-term memory, long-term memory, language ability, visual construction, list-generating fluency, abstraction, and judgment.¹⁷ The CASI score ranges from 0 to 100. Scores of the Mini-Mental State Examination and the modified Mini-Mental State Examination can also be estimated from subsets of the CASI items.¹⁷ Subjects scoring 86 or higher were directly entered into the ACT cohort as cognitively intact. Those with a score lower than 86 or an invalid CASI score underwent additional medical record review and further evaluation. Those who did not meet established criteria for dementia¹⁸ were included in the ACT inception cohort. Of a random sample of 6,782 subjects, 1,360 were ineligible because they were current residents of a nursing home, were participating in another research study, had existing dementia, or met criteria for dementia as a result of screening and examinations. Of 5,422 eligible subjects, 2,841 chose not to participate for a variety of medical, personal, and other reasons before they were formally approached and asked for consent; 2,581 enrolled in the study. Nonresponse has been described elsewhere.¹⁵ Briefly, refusal was more common in the oldest age group (≥ 85), women, and African-American and minority groups. Dementia and AD incidence rates from the ACT study have been published. These rates were in accordance with rates reported in U.S. and European cohort studies.¹⁵

The present study sample was selected from the 2,581 ACT participants. One thousand eight hundred ninety-four subjects whose CASI scores were above the 25th percentile (CASI scores 91–100) were selected. Six hundred eighty-

seven subjects whose CASI scores were in the lowest quartile (CASI scores 62–90 or invalid scores) were excluded because the lowest quartile group might include subjects who had cognitive impairment or impending dementia, and thus, it could not be ascertained that subjective memory deterioration preceded the development of dementia in this group. Eleven subjects who did not have baseline data on subjective memory change were also excluded. Of 1,883 subjects included in this study, 1,739 had at least one follow-up visit; 144 died or discontinued right after their baseline visits.

Incident Dementia

Biennial examinations were conducted to identify incident dementia cases. Subjects were rescreened using the CASI at biennial examinations. Those who scored 86 or higher on the CASI remained in the ACT cohort. CASI scores of less than 86 at follow-up triggered a full standardized clinical examination. The results of rescreening using the CASI and the clinical examination were reviewed at a consensus diagnosis conference that included at least the examining physician, a neuropsychologist, another study physician, and the study nurse. Subjects who did not meet the criteria for dementia remained in the ACT cohort, receiving follow-up with biennial examinations, whereas subjects who met the criteria for dementia were considered as incident dementia cases and were followed annually to confirm the accuracy of the dementia diagnosis. Subjects who met the *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition* (DSM-IV) criteria¹⁸ for dementia were considered incident dementia cases. Dementia type was determined using the National Institute of Neurological and Communicative Disease and Stroke/Alzheimer's Disease and Related Disorder Association criteria¹⁹ for AD and DSM-IV criteria¹⁸ for other types of dementia.

Cognitive Function

Overall cognitive ability was measured using the CASI. Memory performance was measured using the short-term memory recall (STM), which is the three-word recall within the CASI test. Cognitive decline was defined as a drop in CASI scores to 86 or lower at follow-up, because the cutoff CASI score value of 86 was the threshold used to trigger clinical examinations.

Assessment of Subjective Memory Change

Subjective memory change, as a part of memory and functioning evaluation, was assessed using five items. "In the past 10 to 20 years or since the last visit, do you think: (1) Your ability to remember the names of people you have just met has changed? (2) Your ability to remember the faces of people you have just met has changed? (3) Your ability to remember the names of close friends or relatives has changed? (4) Your ability to remember appointments correctly has changed? (5) Your ability to judge the passage of time and guessing the time of day without looking at a clock or the sun has changed?" Response categories for each item were 1 = definitely improved, 2 = slightly improved, 3 = no change, 4 = slightly deteriorated, and 5 = definitely deteriorated. The Subjective Memory Rating Scale (SMRS) was composed by summing the five items and ranged from 5 to 25. A score of 15 on the SMRS indicated no change on

average, and a score of 20 indicated slight deterioration in all items on average. Subjective memory deterioration was defined as an SMRS score of 20 and above.

Other Variables

Demographic variables were age, sex, ethnicity, and years of education. Family history of AD was defined as having a first-degree relative (parent or sibling) with AD. Depression was measured using the 11-item Center for Epidemiologic Studies Depression Scale (CES-D)²⁰ at baseline. CES-D scores ranged from 0 to 33, with higher scores representing more depressive symptoms. A blood sample was drawn from consenting subjects at baseline to obtain apolipoprotein E (ApoE) genotype.

Data Analysis

Cronbach alpha test was performed to assess the internal consistency of the SMRS. Exploratory factor analysis was used to describe relationships between the five memory items.

Because many factors, such as depression,^{2-4,6,9} education level,^{2,15} ApoE genotype,^{9,21} and a family history of AD, could confound self-reported subjective memory, confounders needed to be examined and adjusted. Therefore, logistic regression was used to examine factors that might be associated with subjective memory deterioration at baseline. The analysis used the factors age; sex; years of education; family history of AD; ApoE genotype; and baseline CES-D, CASI, and STM scores.

To investigate whether subjective memory deterioration was associated with cognitive decline, logistic regression was performed using cognitive decline as the response variable. Not only the temporal relationship of subjective memory deterioration preceding cognitive decline, but also the co-occurrence of subjective memory deterioration and performance decline was examined to assess the validity of subjective memory deterioration. In this analysis, subjective memory deterioration was classified as subjective memory deterioration at baseline or new subjective memory deterioration at follow-up. The variables adjusted for were age; sex; years of education; family history of AD; ApoE genotype; and baseline CES-D, CASI, and STM scores.

To evaluate the temporal relationship between subjective memory deterioration and incident dementia, Cox regression was performed using years of age for the time scale, left truncated at age of entry into the study. The outcome of interest was the time from the age of entry into the study to the age of dementia. The predictor of interest was subjective memory deterioration at baseline. Hazard ratios (HRs) in the Cox model were calculated at years of age during the study. Because onset of clinical dementia, particularly AD, is not a discretely timed event, it could only be determined that a subject developed dementia during the study period between the time of dementia diagnosis and the previous examination. The halfway point between diagnosis and the previous examination was taken as the event time for dementia. Subjects who died or discontinued the study before developing dementia were considered censored at the last visit date. Subjects who did not develop dementia during the study were considered censored at the last follow-up date. Variables adjusted for were baseline cognitive performance, ApoE genotype, family history of AD, baseline

CES-D scores, sex, and education. The Schoenfeld residual test²² was used to check the proportional hazard assumption. If the assumption failed, time-varying covariates were introduced.

To investigate potential effect modifications, the interactions between subjective memory deterioration and other covariates were tested in these regression models. Effect modification was considered to be present if the coefficient for the interaction was found to be statistically significant at $P = .05$.

RESULTS

Study participants were followed from 1994 to 2002, with a mean \pm standard deviation follow-up of 5.2 ± 2.1 years. Of 1,883 subjects, 1,288 remained dementia free, 326 died, 143 dropped out, and 126 developed dementia (83 AD). Table 1 shows the baseline characteristics of study participants who remained dementia free, developed dementia, dropped out or died. Table 2 shows the baseline characteristics of study population by levels of subjective memory deterioration.

Over 5 years of follow-up, 15% of persons with baseline subjective memory deterioration developed dementia, in comparison with 13% of persons with family history of AD and 11% of persons with ApoE $\epsilon 3/4$ or $\epsilon 4/4$ alleles. By contrast, dementia occurred in 6% of persons without baseline subjective memory deterioration, 6% of persons without family history of AD, and 6% of persons with ApoE $\epsilon 3/3$ alleles.

The SMRS had a Cronbach alpha coefficient of 0.6. Exploratory factor analysis found one common factor with an eigen value greater than 1 and factor loadings from 0.4 to 0.5 for the five items of remembering names, faces, friends and relatives, and appointments and judging time.

Factors associated with subjective memory deterioration at baseline were older age, male sex, high CES-D scores, and presence of the ApoE $\epsilon 2/4$ alleles. Odds ratios of subjective memory deterioration for 5-year increments of age, male sex, one-point increase of CES-D score, and presence of ApoE $\epsilon 2/4$ alleles versus $\epsilon 3/3$ alleles were 1.28 (95% confidence interval (CI) = 1.07–1.54), 1.96 (95% CI = 1.24–3.10), 1.10 (95% CI = 1.06–1.15), and 3.14 (95% CI = 1.10–8.99), respectively. Baseline subjective memory deterioration was not associated with performance on the cognitive and memory tests using CASI and STM. Although it was expected that a family history of AD might lead to concerns about memory deterioration, the data showed that family history of AD was not associated with subjective memory deterioration.

Subjective memory deterioration was associated with cognitive decline at follow-up. Of 1,739 subjects who had been followed, 1,599 reported no subjective memory deterioration at baseline and follow-up, 79 reported subjective memory deterioration at baseline, and 61 reported new subjective memory deterioration at follow-up. During follow-up, 202 subjects declined in their cognitive performance, with a drop in CASI score to 86 or lower: 165 (10.3%) of those without subjective memory deterioration, 21 (34.4%) of those with subjective memory deterioration at follow-up, and 16 (20.3%) of those with subjective memory deterioration at baseline. Odds ratios of cognitive decline

Table 1. Baseline Characteristics of Study Participants by Follow-Up Status

Baseline Variables	Nondemented (n = 1,288; 68%)	Demented (n = 126; 7%)	Dropped Out or Died (n = 469; 25%)	Total (N = 1,883; 100%)
Age, mean \pm SD	73.5 \pm 5.2	78.4 \pm 5.5	76.5 \pm 6.6	74.6 \pm 5.8
Sex, n (%)				
Male	489 (64)	55 (7)	215 (28)	759
Female	799 (71)	71 (6)	254 (23)	1,124
Ethnicity, n (%)				
White	1,208 (68)	122 (7)	442 (25)	1,772
Nonwhite	79 (72)	4 (4)	27 (25)	110
Education, years				
<12, n (%)	93 (57)	12 (7)	57 (35)	162
\geq 12, n (%)	1,193 (69)	113 (6)	412 (24)	1,718
Mean \pm SD	14.3 \pm 2.7	14.3 \pm 2.8	13.9 \pm 2.9	14.2 \pm 2.7
Apolipoprotein E alleles, n (%)				
e3/3	738 (70)	61 (6)	254 (24)	1,053
e2/4	28 (82)	3 (9)	3 (9)	34
e2/2 or e2/3	169 (69)	8 (3)	68 (28)	245
e3/4 or e4/4	242 (64)	43 (11)	96 (25)	381
Missing	111 (65)	11 (6)	48 (28)	170
Family history of Alzheimer's disease, n (%)				
No	1,207 (68)	110 (6)	451 (26)	1,768
Yes	82 (71)	15 (13)	18 (16)	115
Subjective memory rating scale (5–25)				
\leq 15, n (%)	373 (70)	26 (5)	44 (8)	536
16–19, n (%)	865 (69)	87 (7)	90 (7)	1,260
\geq 20, n (%)	50 (57)	13 (15)	9 (10)	87
Score, mean \pm SD	16.4 \pm 1.7	16.9 \pm 2.0	16.3 \pm 2.1	16.4 \pm 1.8
Short-term memory recall score (0–12)				
\leq 9, n (%)	122 (57)	29 (14)	14 (7)	214
10–12, n (%)	1,165 (70)	97 (6)	129 (8)	1,669
Mean \pm SD	11.1 \pm 1.1	10.5 \pm 1.4	10.9 \pm 1.1	11.0 \pm 1.2
Cognitive ability screening instrument score (0–100)				
91–93, n (%)	309 (59)	50 (10)	56 (11)	520
94–97, n (%)	676 (68)	64 (6)	65 (7)	988
98–100, n (%)	303 (81)	12 (3)	22 (6)	375
Mean \pm SD	95.4 \pm 2.4	94.3 \pm 2.3	94.7 \pm 2.3	95.2 \pm 2.4
Depression				
Center for Epidemiologic Studies Depression scale score, mean \pm SD (range 0–33)	3.7 \pm 3.9	5.3 \pm 4.8	4.7 \pm 4.8	4.1 \pm 4.2

SD = standard deviation.

were 2.7 (95% CI = 1.45–4.98) and 3.5 (95% CI = 1.92–6.28), respectively, for subjects who reported subjective memory deterioration at baseline and who reported new subjective memory deterioration at follow-up, after adjusting for baseline performance on CASI and STM, baseline CES-D scores, age of reporting subjective memory deterioration, ApoE genotype, and family history of AD.

Subjective memory deterioration at baseline was associated with incident dementia during follow-up, as illustrated in Figure 1. Of 1,739 participants who had at least 1 year of follow-up, 126 developed dementia, and 1,613 were censored as no dementia at the last follow-up. Age modified the association between subjective memory deterioration and incident dementia, as indicated by the statistically significant interaction between subjective memory deterioration and age at baseline in the Cox model. For example, for persons reporting subjective memory deterioration at the ages of 70, 75, and 80, the HRs of developing dementia were 6.0 (95% CI = 2.1–18), 3.2 (95% CI = 1.6–6.2), and 1.6 (95% CI = 0.86–3.1), respectively. These HRs were adjusted for age, ApoE genotype, family history of AD, and baseline STM and CES-D scores. The baseline CASI score was no longer a significant predictor for incident dementia after including the baseline STM score in the Cox model. No subject entering the study at the age of 65 developed dementia during the study period.

Table 2. Baseline Characteristics by Levels of Subjective Memory Deterioration

Baseline Characteristic	Subjective Memory Rating Scale Score			
	15* (n = 536; 28%)	16-19† (n = 1,260; 67%)	20‡ (n = 87; 5%)	Total (N = 1,883; 100%)
Age, mean \pm SD	73.6 \pm 5.5	74.9 \pm 5.9	76.1 \pm 6.0	74.6 \pm 5.8
Sex, n (%)				
Male	217 (29)	494 (65)	48 (6)	759
Female	319 (28)	766 (68)	39 (3)	1,124
Ethnicity, n (%)				
White	503 (28)	1,195 (67)	74 (4)	1,772
Nonwhite	33 (30)	64 (58)	13 (12)	110
Apolipoprotein E alleles, n (%)				
e3/3	306 (29)	694 (66)	53 (5)	1,053
e2/4	4 (12)	25 (74)	5 (15)	34
e2/2 or e2/3	63 (26)	167 (68)	15 (6)	245
e3/4 or e4/4	104 (27)	265 (70)	12 (3)	381
Missing	59 (35)	109 (64)	2 (1)	170
Family history of Alzheimer's disease, n (%)				
No	502 (28)	1,183 (67)	79 (5)	1,764
Yes	34 (29)	77 (65)	8 (7)	119
Short-term memory recall score (0-12)				
≤ 9 , n (%)	41 (19)	184 (77)	9 (4)	214
10-12, n (%)	495 (30)	1,096 (66)	78 (5)	1,669
Mean \pm SD	11.1 \pm 1.1	10.9 \pm 1.2	10.95 \pm 1.0	11.0 \pm 1.2
Cognitive ability screening instrument score (0-100)				
91-93, n (%)	117 (23)	378 (73)	25 (5)	520
94-97, n (%)	311 (31)	632 (64)	45 (5)	988
98-100, n (%)	108 (29)	250 (67)	17 (5)	375
Mean \pm SD	95.4 \pm 2.3	95.1 \pm 2.4	95.1 \pm 2.4	95.2 \pm 2.4
Education, years				
< 12, n (%)	51 (31)	101 (62)	10 (6)	162
≥ 12 , n (%)	485 (28)	1,156 (68)	77 (4)	1,718
Mean \pm SD	14.3 \pm 2.7	14.2 \pm 2.7	14.6 \pm 3.1	14.2 \pm 2.7
Depression				
Center for Epidemiologic Studies Depression scale score, mean \pm SD (range 0-33)	3.0 \pm 3.4	4.3 \pm 4.4	6.2 \pm 5.0	4.1 \pm 4.2

* No self-perceived deterioration.

† Some deterioration.

‡ Definite deterioration.

SD = standard deviation.

To assess the potential biases that subjects who died or were lost to follow-up might introduce, logistic regression was performed to examine whether subjective memory deterioration was associated with death and dropout. It was found that subjective memory deterioration at baseline was not associated with death and dropout. Therefore, the influence of death and dropout on the estimates for the associations of subjective memory deterioration with cognitive decline and incident dementia was minimal in this study.

DISCUSSION

It was found that older people with subjective memory deterioration at baseline were at greater risk of developing dementia during 5 years of follow up, even though at base-

line they performed as well as those without subjective memory deterioration on the CASI and the short-term memory recall. These findings support the hypothesis that subjective memory deterioration provides additional information about future dementia at a time when a screening test does not show objective evidence of cognitive impairment. A screening test administered at one point measures levels of cognitive function but not the change in cognitive function of a person over time. In contrast, subjective memory deterioration assesses self-perceived change over time in a person, but longitudinal assessments including better measures of subjective memory deterioration and objective memory function are necessary to determine the validity of self-reported memory changes over time. Nevertheless, it is reasonable to speculate that high-performing individuals may be aware of decline in their memory, when




Figure 1. Kaplan-Meier survival estimates for the probabilities of being dementia free. Subjects who reported subjective memory deterioration (SMD) but scored normal on the Cognitive Ability Screening Instrument at baseline were less likely to be dementia free than those who reported no SMD.

they are still able to perform in the normal range on a screening test. Subjective memory deterioration might indicate mild cognitive impairments that could be detected if a more comprehensive assessment were used.

One of the strengths of this study is its prospective, longitudinal nature. This design avoids some of the pitfalls inherent in cross-sectional measurement of longitudinally developing processes.²³ The principal findings of this study are that a high level of subjective memory deterioration was associated with cognitive decline and future dementia in older people who did not have objective cognitive impairment according to a screening test; age modified the association between subjective memory deterioration and future dementia, with younger persons reporting subjective memory deterioration having a higher risk of dementia than older persons reporting subjective memory deterioration. One report from the ASE found that, for 2,169 subjects with normal baseline cognition, memory complaints were associated with incident AD in 3 years of follow-up.⁷ Another ASE report found that memory complaints were associated with accelerated cognitive decline over 6 years of follow-up in cognitively normal elderly.⁹ These two studies used a single question to assess subjective memory: "Do you have complaints about your memory?" A smaller longitudinal study with 331 subjects (20 subjects with dementia) and 7 to 8 years follow-up showed that memory complaints reflected perceptions of past memory performance and an early manifestation of memory impairment.⁵ The association between complaints and future memory performance was present even when dementia cases were excluded, but the magnitude of the association decreased. In this study,⁵ memory complaints were assessed using four specific questions regarding whether a person's memory had become worse (1) remembering recent things, (2) remembering where belongings were located, (3) recalling conversations, and (4) remembering appointments and social arrangements) and compared with earlier in life. Although the differences in subjective memory measures and study methods make results difficult to compare across the studies, the re-

sults of the current study are consistent with these longitudinal studies^{5,7,9} and provide evidence that subjective memory deterioration can be associated with subsequent cognitive decline and future dementia.

Subjective memory deterioration was associated with a higher risk of developing dementia in younger persons than in older persons. Although these analyses showed that the magnitude of this effect modification by age was large, the finding needs to be confirmed and replicated in other studies. It is possible that small numbers of persons with subjective memory deterioration who developed dementia in various age groups may generate an unstable estimate for the interaction of subjective memory deterioration with age. If these findings are confirmed, there are several possible explanations for this effect modification. One possibility is that younger old persons may assess their memory changes more correctly than older persons. Because executive impairments develop with age, self-assessment of subjective memory may become less reliable in older age. Another possible explanation is that memory problems are more common in older age groups (≥ 80), so it is more difficult to distinguish whether subjective memory deterioration is an early sign of dementia or so-called normal aging. This is probably related to the fact that cognitive function changes, like so many age-related changes, have increased variance with age and increased frequency with age; thus, the distinction between normal and abnormal becomes more difficult at older ages.

The data showed that subjective memory deterioration at baseline and new subjective memory deterioration at follow-up were strongly associated with decline in cognitive performance. The subjective memory deterioration measure has adequate face validity, but the subjective memory deterioration scale was imprecise and had a low predictive value for dementia, which implies that, in its present form, it would likely have limited clinical utility. This measure is not proposed as a diagnostic test for detecting early dementia. Better measurement of subjective memory may help improve the ability to identify a subgroup of individuals at high risk for developing dementia. The need for instrument development in assessing changes in subjective memory, which combined with clinical tests, might have potential to be a practical way of identifying individuals in an early stage of developing dementia, is stressed. Of the five memory items, the judgment of time reflects orientation rather than memory. Thus, subjective memory deterioration as defined in this study might assess a broader range of cognitive decline than memory per se.

The relationship between subjective memory complaints and depression has been reported.^{24,25} It was found in the current study that subjective memory deterioration at baseline was associated with higher CES-D scores. After adjusting for CES-D score, there was still a significant relationship between baseline subjective memory deterioration and incident dementia, but because extensive psychometric measurements were not performed on persons who were in the normal range on the screening test, it cannot be ascertained whether subjective memory deterioration was due to subtle but real impairments or to some other causes like depression.

Studies have reported that education levels were associated with memory complaints^{2,15} and cognitive decline

and dementia,²⁶⁻²⁸ although other studies had different results.^{29,30} It was found that education levels were not associated with subjective memory deterioration at baseline or incident dementia at follow-up in this well-educated cohort. The analyses might have been underpowered to detect such associations if they exist, because there were few subjects with low levels of education (2% with ≤ 8 years and 6% with 9-11 years of education). These findings may apply only to relatively well-educated populations.

It may be important to identify individuals at high risk of developing AD and other types of dementia at an early stage. When effective interventions become available, early treatment may slow the development of disease and prevent further irreversible damage from occurring. The current study suggests that some older individuals with normal cognitive functioning who go on to develop dementia appear to be able to discern changes in their memory before those changes are apparent on cognitive screening tests. This study adds to the growing body of evidence that suggests that a high level of subjective memory deterioration may represent an early sign of dementia in a subset of individuals. Present research tools are too blunt to apply widely, and future work should develop better tools to measure this area.

ACKNOWLEDGMENTS

The authors would like to thank all members of the University of Washington Alzheimer's Disease Patient Registry for their contributions made to the project. We also thank Dr. Linda Teri for reviewing an early draft of the manuscript. We acknowledge the contribution of the Honolulu Asian Aging Study, which was the source of the subjective memory change questions.

REFERENCES

- Jonker C, Launer LJ, Hooijer C et al. Memory complaints and memory impairment in older individuals. *J Am Geriatr Soc* 1996;44:44-49.
- Gagnon M, Darrigues JF, Mazaux JM et al. Self-reported memory complaints and memory performance in elderly French community residents: Results of the PAQUID research program. *Neuroepidemiology* 1994;13:145-154.
- O'Connor DW, Pollitt PA, Roth M et al. Memory complaints and impairment in normal, depressed, and demented elderly persons identified in a community survey. *Arch Gen Psychiatry* 1990;47:224-227.
- Tobiansky R, Blizard R, Livingston G et al. The Gospel Oak Study stage IV. The clinical relevance of subjective memory impairment in older people. *Psychol Med* 1995;25:779-786.
- Jorm AF, Christensen H, Korten AE et al. Memory complaints as a precursor of memory impairment in older people: A longitudinal analysis over 7 to 8 years. *Psychol Med* 2001;31:441-449.
- Schmand B, Jonker C, Geerlings MI et al. Subjective memory complaints in the elderly: Depressive symptoms and future dementia. *Br J Psychiatry* 1997;171:373-376.
- Geerlings MI, Jonker C, Bouter LM et al. Association between memory complaints and incident Alzheimer's disease in elderly people with normal baseline cognition. *Am J Psychiatry* 1999;156:531-537.
- Schofield PW, Marder K, Dooneief G et al. Association of subjective memory complaints with subsequent cognitive decline in community-dwelling elderly individuals with baseline cognitive impairment. *Am J Psychiatry* 1997;154:609-615.
- Dik MG, Jonker C, Comijs HC et al. Memory complaints and APOE- $\epsilon 4$ accelerate cognitive decline in cognitively normal elderly. *Neurology* 2001;57:2217-2222.
- Schmand B, Jonker C, Hooijer C et al. Subjective memory complaints may announce dementia. *Neurology* 1996;46:121-125.
- Jorm AF, Christensen H, Korten AE et al. Do cognitive complaints either predict future cognitive decline or reflect past cognitive decline? A longitudinal study of an elderly community sample. *Psychol Med* 1997;27:91-98.
- Blazer DG, Hays JC, Fillenbaum GG et al. Memory complaint as a predictor of cognitive decline. *J Aging Health* 1997;9:171-184.
- Wilson RS, Evans DA. How clearly do we see our memories? *J Am Geriatr Soc* 1996;44:93-94.
- Rabbitt P, Abson V. 'Lost and Found': Some logical and methodological limitations of self-report questionnaires as tools to study cognitive ageing. *Br J Psychol* 1990;81:1-16.
- Kukull WA, Higdon R, Bowen JD et al. Dementia and Alzheimer's disease incidence. *Arch Neurol* 2002;59:1737-1746.
- Wang L, van Belle G, Kukull WA et al. Predictors of functional changes: A longitudinal study of nondemented people aged 65 and older. *J Am Geriatr Soc* 2002;50:1525-1534.
- Teng EL, Hasegawa K, Homma A et al. The Cognitive Ability Screening Instrument (CASI): A practical test for cross-cultural epidemiological studies of dementia. *Int Psychogeriatr* 1994;6:45-62.
- Diagnostic and Statistical Manual of Mental Disorders, 4th Ed. Washington, DC: American Psychiatric Association, 1994.
- McKhann G, Drachman D, Folstein M et al. Clinical diagnosis of Alzheimer's disease. Report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 1984;34:939-944.
- Radloff LS. The CES-D scale. A self-reported depression scale for research in general population. *Appl Psychol Meas* 1977;1:385-401.
- Small GW, Chen ST, Komo S et al. Memory self-appraisal in middle-aged and older adults with the apolipoprotein E-4 allele. *Am J Psychiatry* 1999;156:1035-1038.
- Grambsch PM, Therneau TM. Proportional hazard tests and diagnostics based on weighted residuals. *Biometrika* 1994;81:515-526.
- Kraemer HC, Yesavage JA, Taylor JL et al. How can we learn about developmental processes from cross-sectional studies, or can we? *Am J Psychiatry* 2000;157:163-171.
- Williams MJ, Little MM, Scates S et al. Memory complaints and abilities among depressed older adults. *J Consult Clin Psychol* 1987;4:595-598.
- O'Hara MW, Hinrichs JV, Kohout FJ et al. Memory complaint and memory performance in the depressed elderly. *Psychol Aging* 1986;3:208-214.
- Aevartsson O, Skoog I. A longitudinal population study of the Mini-Mental State Examinations in the very old: Relation to dementia and education. *Dement Geriatr Cogn Disord* 2000;11:166-175.
- Stern Y, Gurland B, Tatemichi TK et al. Influence of education and occupation on the incident of Alzheimer's disease. *JAMA* 1994;271:1004-1010.
- Ott A, van Rossum CT, van Harskamp F et al. Education and incidence of dementia in a large population-based study. The Rotterdam Study. *Neurology* 1999;52:663-666.
- Cobb JL, Wolf PA, Au R et al. The effect of education on the incidence of dementia and Alzheimer's disease in the Framingham Study. *Neurology* 1995;45:1707-1712.
- Pakei ES, Brayne C, Huppert FA et al. Incidence of dementia in a population older than 75 years in the United Kingdom. *Arch Gen Psychiatry* 1994;51:325-332.

Treating Mild Cognitive Impairment

The Absence of Evidence

E. S. Roach, MD

The absence of evidence does not constitute evidence of absence.

William Safire

Most people with Alzheimer disease (AD) develop mild cognitive impairment (MCI) as an initial step toward progressively severe dementia, but there are numerous other causes of MCI. If we define AD by its pathological changes rather than by the severity and variety of its signs and symptoms, then many, perhaps even most, individuals with MCI will eventually prove to have early AD. And if a medication improves cognitive function of a person with AD, even modestly, why not treat individuals with MCI? It seems reasonable to guess that a medication known to alleviate symptoms of AD could also be useful in the different conditions with overlapping clinical manifestations.

Scientific disagreements often arise because key facts are missing, and there are many questions we need to answer more completely in order to know how to best deal with MCI. How effective are acetylcholinesterase inhibitors in individuals with MCI? If these drugs improve function in 1 MCI subgroup, can we assume that they will have the same effect in individuals in other subgroups? If a medication is effective, does its use make enough difference to justify the effort, cost,

and risks of treatment? If a medication meaningfully improves quality of life, can we justify not using it?

Gauthier and Touchon¹ argue that although individuals with MCI have an increased likelihood of progressing to frank dementia during a 10-year interval, many of them remain stable and some even revert to normal during this period. Moreover, the heterogeneity and unpredictability of MCI make it difficult to treat affected individuals with cholinesterase inhibitors. Randomized clinical trials using these drugs in people with MCI have not demonstrated consistent improvement in cognition or lowered the likelihood of developing AD after MCI.¹

While Petersen and Morris² agree that MCI is heterogeneous, they argue that much of this variability is likely to disappear as we become more precise in defining MCI subtypes and identifying its different causes. More accurate classification, in turn, creates an opportunity to offer treatment to selected individuals before their symptoms become advanced.² Both sides agree on the need for continuing studies.

Reasonable arguments can be made both for and against treating individuals with MCI with cholinesterase inhibitors. Mild cognitive impairment is not a unique clinical entity, and proof that the available drugs benefit conditions other than AD is lacking. Nevertheless, many individuals with MCI progress to AD, and it seems intuitive that ear-

lier treatment should be more effective. This debate should not obscure the fact that the currently available drugs are not dramatically effective even under ideal circumstances, and we must continue our efforts to develop other types of therapy for MCI and AD.

Accepted for Publication: March 31, 2005.

Correspondence: E. S. Roach, MD, Department of Neurology, Wake Forest University School of Medicine, Medical Center Boulevard, Winston-Salem, NC (sroach@wfubmc.edu).

REFERENCES

1. Gauthier S, Touchon J. Mild cognitive impairment is not a clinical entity and should not be treated. *Arch Neurol.* 2005;62:1164-1168.
2. Petersen RC, Morris JC. Mild cognitive impairment as a clinical entity and treatment target. *Arch Neurol.* 2005;62:1160-1163.

Submissions

We invite you to suggest controversies and 3 names of potential authors to write about either side of the issue. Do not feel that you need to exclude yourself. Please communicate with E. S. Roach, MD, Department of Neurology, Wake Forest University School of Medicine, Medical Center Boulevard, Winston-Salem, NC 27157; phone (336) 716-1619; fax (336) 716-9489 (sroach@wfubmc.edu).

Author Affiliations: Wake Forest University School of Medicine, Winston-Salem, NC.

Mild Cognitive Impairment Is Not a Clinical Entity and Should Not Be Treated

Serge Gauthier, MD, FRCPC; Jacques Touchon, MD

THERE HAS BEEN GREAT progress in the diagnosis and treatment of Alzheimer disease (AD) and related dementias thanks to the availability of well-validated diagnostic criteria (*Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV)*, National Institute of Neurological and Communicative Disorders and Stroke-the Alzheimer's Disease and Related Disorders Association [NINCUS-ADRDA]); and a large number of randomized clinical trials using cholinesterase inhibitors (AChEIs) and Memantine. Work from specialized neurological clinics suggests that mild cognitive impairment (MCI) is an early stage of AD, and there is a temptation for neurologists to prescribe AChEIs in this population. On the other hand, there is epidemiological evidence that many subjects labeled as having MCI do not worsen over time and may revert to normal cognitive abilities. A diagnosis of MCI as a prodromal stage of AD in such individuals would be inaccurate and carry a heavy personal and societal burden. Reversible causes of MCI may be found, such as depression, upper airway obstruction, and a variety of metabolic, nutritional, or sensory impairments. Since MCI does not constitute a homogeneous clinical syndrome, it is inappropriate to propose a specific drug treatment such as AChEIs, but the recognition that MCI is a risk state toward fur-

ther cognitive decline is clinically relevant, and control of risk factors such as systolic hypertension, hypercholesterolemia, diabetes mellitus, atrial fibrillation, transient ischemic attacks, and strokes may delay progression to dementia.

GENERAL CONCEPT OF MCI

The concept of MCI is being widely used in epidemiological and clinical studies as an intermediate stage between cognitive normality and dementia. A high number of persons have been found to have MCI in the population at large, and many are now being referred to memory clinics. There is an expectation that even larger numbers of individuals with memory complaints will be coming forward for assessment in the near future.

The heterogeneity of MCI has been recognized in a report of the Quality Standards Subcommittee of the American Academy of Neurology,¹ where it has been recommended to qualify the term MCI with an appropriate modifier such as "amnesic," which was defined operationally by Petersen et al² in 1999 (Figure 1).

MCI IN SPECIALTY CLINICS

Specialized clinics in Rochester, Minn, and St Louis, Mo, have established that amnesic MCI is frequently a prodrome to AD³ or its first manifestation.⁴ Conversion rates from amnesic MCI to AD range from 12% to 15% per year.⁵ The heterogeneity within MCI has been noted and a new classification has been proposed, based predominantly on neuropsychological profiles, and includes amnesic or single memory MCI, multiple-domain MCI, and single nonmemory MCI.⁴

Subjective memory complaint, preferably corroborated by an informant
Memory impairment relative to a previously documented level of education-matched healthy controls
Relatively normal overall cognitive function
Largely intact activities of daily living
No clinical correlates

Figure 1. Operational definition of amnesic mild cognitive impairment.

MCI IN POPULATION STUDIES

Estimates of prevalence of MCI in population-based epidemiological studies vary greatly depending on the definition in use and range from 3% to 16.6%.⁶ The former figure is obtained from specialized referral clinics using amnesic MCI criteria, whereas the latter figure has been documented in the Canadian Study of Health and Aging where the term cognitive impairment no dementia (CIND) was defined as various categories of impairment identified in a clinical examination and neuropsychological tests.⁷ The most common cause of CIND was circumscribed memory impairment, a close equivalent to amnesic MCI, with a prevalence of 5.3%.⁸ The follow-up of individuals from that cohort with MCI defined by different criteria led to a wide range of conversion to dementia (20.0%-50.9%), AD (11.6%-28.8%), and death (30.1%-42.4%).⁹

An epidemiological study based on a general practice research network in France revealed a prevalence of 3.2% for MCI, which was a poor predictor of dementia within a 3-year period, with an 11.1% conversion rate and many subjects reverting to normal cognition.⁹ Another cohort in France demonstrated a prevalence of 2.8% for MCI and an annual conversion rate of 8.3% across 3 years, but within 3 years more than 40% of subjects with MCI had reverted to normal cognition.¹⁰

Author Affiliations: Departments of Neurology and Neurosurgery, Psychiatry, and Medicine, McGill University and Alzheimer Disease Research Unit, McGill Centre for Studies in Aging, Montreal, Quebec (Dr Gauthier); Department of Neurology, Université de Montpellier, Montpellier, France (Dr Touchon).

Entry criteria for the Donepezil Hydrochloride Longitudinal Study
 Memory complaints corroborated by an informant
 Abnormal memory function documented by delayed recall of 1 paragraph from the Logical Memory III subtest of the Wechsler Memory Scale-Revised, adjusted for age and education
 Normal general cognitive function as determined by a physician's judgment based on the CDR and the MMSE
 No or minimal impairment in ADL
 No clinical dementia
 Age 55-90 y
 In good general health with no significant cardiovascular disease and no history of stroke
 Not depressed, with a Hamilton Depression Rating Scale score ≤ 12
 No infection, inflammation, or local lesions on computed tomography or magnetic resonance imaging
 CDR global score of 0.5, with a score of at least 0.5 in the memory domain
Entry criteria for the Rivastigmine Hydrochloride Studies
 Cognitive decline of gradual onset and slow progression
 Abnormal memory function documented by a delayed Recall Score ≤ 1 on the New York University Delayed Paragraph Recall Test
 Insufficient impairment on ADL to warrant a diagnosis of dementia
 Age 50-90 y
 CDR global score of 0.5, with a score of at least 0.5 in the memory domain
Entry criteria for the Rivastigmine Tardive Study
 Memory complaints corroborated by an informant
 Abnormal memory function documented by the New York University Delayed Paragraph Recall Test with a cutoff dysfunction score ≤ 3
 No clinical dementia
 Age 55-95 y
 Not depressed, with a Hamilton Depression Rating Scale score ≤ 12 , item 1 (depressed mood) score ≤ 1
 CDR global score of 0.5

Figure 2. Entry criteria for studies. ADL indicates activities of daily living; CDR, Clinical Dementia Rating; MMSE, Mini-Mental State Examination.

The Kungsholmen Project is an older population study looking at MCI, where a distinction was made within CIND as mild, moderate, or severe based on Mini Mental State Examination scores (mild, 1 SD lower than age-specific and education-specific mean scores; moderate, 1.5 SDs; severe, 2.5 SDs). Depression and cerebrovascular disease were more common in the CIND population compared with unimpaired subjects; similar proportions of subjects with CIND progressed to dementia, death, and cognitive improvement across 3 years.¹¹

A community-based sample in a rural area of the United States showed that 3% to 4% of elderly persons without dementia met MCI criteria and had a higher risk of progressing to dementia across 10 years, but many remained stable or reverted to normal cognition during follow-up.¹²

MCI IN RANDOMIZED CLINICAL TRIALS

The operational definition of amnesic MCI has made possible large-scale, placebo-controlled, randomized clinical trials to test the hypothesis that the treatment of MCI with AChEIs, such as donepezil hydrochloride, rivastigmine tartrate, or

galantamine hydrobromide, can delay the diagnosis of AD and/or improve cognitive impairments. Different MCI populations are defined for each study.¹³⁻¹⁵ (Figure 2).

Results of these studies are only partially analyzed and reported, but for the purpose of the arguments in this "Controversies" section, there was a wide range of rate of conversion to AD across 2 to 4 years. This variability is based on differences in the number of carriers of the apolipoprotein E4 genotype, variations in the memory impairment cutoff scores at entry, and the heterogeneity of MCI, even in the relatively pure amnesic subtype. Efficacy of AChEIs overall in improving cognition or delaying the conversion from MCI to AD have been disappointing so far.

MCI AT HIGHER RISK OF CONVERSION TO DEMENTIA

There are patterns emerging from specialized neurological clinics, population studies, and randomized clinical trials for higher risks of progression to dementia. It is likely that models combining multiple risk factors, such as the ones listed in Figure 3,^{16,17} will be needed for predicting the risk of individuals with MCI for progression to dementia.

Poor performance on delayed recall and episodic function tests
 Caretaker report of impaired daily function, particularly for ability to use the telephone, handle transportation, responsibility for own medication, ability to handle finances
 Hippocampal atrophy (2.5 SDs lower than normal at age and sex)
 Apolipoprotein E4 genotype
 Multiple vascular risk factors

Figure 3. Mild cognitive impairment at higher risk of progression to dementia.

MCI IN CLINICAL PRACTICE

A classification of MCI has been proposed for use in clinical practice based on the main subtypes of patients older than 50 years commonly encountered in neurology referral practice¹⁸: dysphoric, vascular, amnesic, miscellaneous, or other medical conditions. The importance of neuropsychiatric symptoms in MCI is starting to emerge from systematic studies using the Neuropsychiatric Inventory, with depression, apathy, and irritability being most common.¹⁹ We think that this is the most common type of MCI, which we label as "dysphoric." The second most common is "vascular" (eg, individuals with cognitive impairment associated with vascular risk factors and vascular lesions on computed tomography and/or magnetic resonance imaging). Less common is amnesic as defined earlier. A "miscellaneous" category could include other causes, such as sleep apnea, alcohol abuse, metabolic and nutritional deficiencies, sensory deprivation, and head injury.

Since the depressive or dysphoric MCI profile may represent a noncognitive prodrome to dementia²⁰ or even subclinical AD,²¹ yearly follow-up visits are suggested beyond treating depressive symptoms with a serotonin reuptake inhibitor or other relevant drug and supportive psychotherapy. Although less likely to deteriorate over time, vascular MCI requires treatment and follow-up of the associated conditions. Finally, amnesic MCI requires follow-up until dementia is diagnosable, at which time an AChEI should be offered. Miscellaneous causes of MCI are reversible for the most part with treatment of the specific etiologies, often coexisting.

Subject action is a format record cognitive problems
These represent a Jackson relative to the subject's
previous abilities
Demonstration of mild cognitive impairment by
cognitive testing, in memory, and/or other
domains
This cognitive impairment does not cause
functional impairment in daily life
Absence of dementia based on clinical judgment

Figure 4. Syndromic definition of mild cognitive impairment.

FUTURE PERSPECTIVE ON MCI

A future perspective on dealing with MCI would be (1) defining MCI as a clinical syndrome with diagnostic criteria, (2) defining clinical subtypes of MCI, and (3) defining etiologic subtypes of MCI and then treating accordingly.

For the first step, a set of clinical criteria based on a clinical approach may be preferred to a predominantly neuropsychological one. These criteria could be created and used in a way similar to DSM IV or NINCDS-ADRDA criteria. As a starting point, we have made a proposal for new MCI criteria (Figure 4).²⁴

A working group of health care professionals and epidemiologists proposed similar criteria for MCI based on 3 diagnostic features: (1) not normal, without dementia, (2) cognitive decline indicated by subject and/or informant report and objective cognitive tests, and (3) preserved basic activities of daily living with some minimal impairment in complex instrumental functions.

The second step in dealing with MCI will be the characterization of the different subtypes, which can be defined using primarily neuropsychological criteria or primarily clinical criteria, and determining their natural history.

The third step will be the characterization of the etiology of each clinical subtype and its specific treatments. It is possible that amnesic MCI will become "very early AD" and be incorporated into updated NINCDS-ADRDA-like criteria.

CONCLUSIONS

Guidelines such as the ones developed by the American Academy of Neurology¹ need to be updated as

new evidence on the natural history of amnesic and other types of MCI become available. Until then, we argue that MCI is not a homogeneous clinical entity and, as such, cannot be treated with a specific therapy such as AChEis. On the other hand, reversible causes of MCI may be found, such as depression, upper airway obstruction, and a variety of metabolic, nutritional, or sensory impairments, whereas control of vascular risk factors such as systolic hypertension, hypercholesterolemia, diabetes mellitus, atrial fibrillation, transient ischemic attacks, and strokes may delay progression to dementia.

Accepted for Publication: March 31, 2005.


Correspondence: Serge Gauthier, MD, FRC PC, McGill Centre for Studies in Aging, 6825 LaSalle Blvd, Verdun (Montreal), Quebec H4J 1K3, Canada (serge.gauthier@mcgill.ca).
Author Contributions: Study concept and design: Gauthier and Touchon. Drafting of the manuscript: Gauthier and Touchon. Administrative, technical, and material support: Gauthier and Touchon.

REFERENCES

1. Peterson RC, Stevens JC, Ganguli M, et al. Practice parameter: early detection of dementia and cognitive impairment (evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*. 2001;56:1123-1142.
2. Peterson RC, Smith GE, Wang SJ, et al. Mild cognitive impairment: clinical characterization and outcome. *Arch Neurol*. 1999;56:303-308.
3. Morris JC, Storandt M, Miller JP, et al. Mild cognitive impairment represents early-stage Alzheimer's disease. *Arch Neurol*. 2001;58:387-408.
4. Peterson RC. Conceptual overview. In: Peterson RC, ed. *Mild Cognitive Impairment: Aging in Alzheimer's Disease*. New York, NY: Oxford University Press, Inc; 2003:1-14.
5. Delamater G. Mild cognitive impairment: prevalence, prognosis, etiology, and treatment. *Lancet Neurol*. 2002;2:19-21.
6. O'Brien JT, Rockwood K, Boake R, et al. Standardization of the diagnosis of dementia in the Canadian Study of Health and Aging. *Neuropsychiatry*. 1995;10:240-256.
7. Graham JE, Rockwood K, Berman BL, et al. Prevalence and severity of cognitive impairment with and without dementia in an elderly population. *Lancet*. 1997;349:1795-1796.
8. Hyman BT, Eady SM. Predicting who will develop dementia in a cohort of Canadian seniors. *Can J Neurol Sci*. 2000;27:13-24.
9. Rinne K, Antero R, Touchon J. Classification of mild cognitive impairment: a population-based validation study. *Neurology*. 2001;56:37-40.
10. Larsson S, Letenneur L, Orgogozo JM, et al. Incidence and outcome of mild cognitive impairment in a population-based prospective cohort. *Neurology*. 2002;59:1594-1598.
11. Palmer K, Wang KX, Winblad B, Fratiglioni L. Differential evolution of cognitive impairment in nondemented older persons: results from the Kungsholmen Project. *Am J Psychiatry*. 2002;159:436-442.
12. Ganguli M, Dodge RH, Sora C, Oakley ST. Mild cognitive impairment: amnesic type; an epidemiological study. *Neurology*. 2004;63:116-121.
13. Grundman M, Petersen RC, Ferris SH, et al. Mild cognitive impairment can be distinguished from Alzheimer's disease and normal aging for clinical trials. *Arch Neurol*. 2004;61:59-66.
14. Goin ML, Wang D, Truyen J. Galantamine for the treatment of mild cognitive impairment: 2 double-blind, placebo-controlled studies. Paper presented at the 11th Congress of the IPA; August 18, 2003; Chicago, Ill.
15. Feldman H, Schallert P, Scarpini E, et al. Behavioral symptoms in mild cognitive impairment. *Neurology*. 2004;62:1190-1201.
16. Chen H, Rajulu G, Balis SH, et al. Cognitive tests that best discriminate between presymptomatic AD and those who remain nondemented. *Neurology*. 2000;55:1567-1573.
17. Barberger-Casari P, Pausanias C, Helmer C, et al. Functional impairment in instrumental activities of daily living: an early clinical sign of dementia? *J Am Geriatr Soc*. 1999;47:456-462.
18. Taheri MH, Albin GM, Borokhovskaya L, et al. Functional decline in patients with mild cognitive impairment: prediction in AD. *Neurology*. 2001;56:750-764.
19. Jack CR, Petersen RC, Xu YC, et al. Prediction of AD with MRI-based hippocampal volume in mild cognitive impairment. *Neurology*. 1998;52:1397-1403.
20. The MGS, Jonker C, Cornier HG, et al. Memory complaints and APOE-epsilon4 in a population-based study. *Neurology*. 2001;57:2217-2222.
21. Kivimäki M, Helkala EL, Lonninen T, et al. Multiple vascular risk factors and late-life mild cognitive impairment: a population-based study. *Neurology*. 2001;56:1685-1690.
22. Gauthier S, Touchon J. Subclassification of mild cognitive impairment in research and in clinical practice. In: Gauthier S, Gauthier P, Cummings JL, eds. *Alzheimer's Disease and Related Disorders Annual*. London, England: Martin Dunitz; 2004:61-69.
23. Hyman BT, Cummings JL. Neuropsychiatric symptoms of mild cognitive impairment. In: Gauthier S, Gauthier P, Cummings JL, eds. *Alzheimer's Disease and Related Disorders Annual*. London, England: Martin Dunitz; 2004:71-79.
24. Schwelb L, Tuohimäki V, O'Brien J, Ames D. Is late-life depression a harbinger of dementia? *Int J Geriatr Psychiatry*. 2002;17:927-935.
25. Oberlin M, Schmand B, Brain AW, et al. Depressive symptoms and risk of Alzheimer's disease in more highly educated older persons. *J Am Geriatr Soc*. 2000;48:1092-1097.
26. Winblad B, Palmer K, Kivimäki M, et al. Mild cognitive impairment—beyond controversies, towards a consensus: report of the International Working Group on Mild Cognitive Impairment. *J Intern Med*. 2004;256:240-246.

PHYSICIANS' DESK REFERENCE

Champion from Beaulieu to ADAS dog shows.
The Rectangles of Painted and Painted into Champion
the Study note: Pseudo 11K.
on study data, and 32 study 50K.




Experiment	50	7	4	0
mobility	1.5	6.7%	16.6%	43.9%
mobility	2.18	15.7%	30.7%	64.1%
mobility	11.1%	61.7%	33.2%	58.1%

Types & Distribution of CBEs cited by firm by industry



and Great Britain's World Food-Grain Study, ready to go into World War II. The study, headed by Dr. Frank Condon, found that 680 percent more warheads would be needed to destroy the world's food supply than to destroy the world's population. The study also found that the world's food supply was more vulnerable to attack than the world's population. The study was a major factor in the decision to develop the atomic bomb.


of treatment, the mean difference in the AHAAS score for the HEMETIS-treated patients compared to the patients on placebo were 3.1 and 4.1 units for



Hours of Treatment	Control (%)	100 mg (%)	200 mg (%)	400 mg (%)
0	100	100	100	100
4	100	100	100	100
8	100	100	100	100
12	100	100	100	100
16	100	100	100	100
20	100	100	100	100
24	100	100	100	100

Figure 1 is a histogram of the distribution of CATH-4 in scores obtained by patients in each of the three treatment groups who had 2 or more of the 10 items of the CATH-4. The mean CATH-4 score for the three groups of patients in this sample from baseline were 0.34 and 0.57 for the two treatment groups.

Figure 8. Cumulative Percentage of Patients Completing 20 Months of Doxorubicin Treatment with Spontaneous Complete Remission by Age Group.




Treatment	Change in ADAMs exp. %†	ADAMs %‡
Placebo	1.6%	1.6%
24 mg/day	4.5%	11.6%
32 mg/day	7.1%	16.7%

100


[illegible]

Figure 10. Time-course of the change




21. Describe the number of the two branches of the two triangles.
22. Three changes of the triangles.
23. The triangles are changed in the same way.
24. The triangles are changed in the same way.
25. The triangles are changed in the same way.
26. The triangles are changed in the same way.
27. The triangles are changed in the same way.
28. The triangles are changed in the same way.
29. The triangles are changed in the same way.
30. The triangles are changed in the same way.

REMARKS: as a predicable consequence of the
logical properties of the
the




PERDRAVIT028
adjuvant for patients and
twice per day for 14 days
of fever within 48 hours
of onset of symptoms
and patients should follow
advice of their doctor.
This medicine should be
used for the treatment of
fever and patients should
follow advice of their
doctor.

Publicity and campaigns about the

[illegible]

DOSEAGE AND ADMINISTRATION

[illegible][illegible]

T

ity) when they were undoubtedly in the most healthy subject was diagnosed as a malignant melanoma of the skin. A 25-year-old male mechanic and his wife were in a professional relationship with a physician (12) known as the "Black and Tan" and 30 negroes' heads or 3 times as many as the white. The doctor was charged by the negroes to the MREED as a negro's head or 3 times as many as the white. The doctor 12 times the MREED as a

active was not negligible in a 6-week study in transgenic AP-2-deficient mice, or in 24-month and multiparous mice up to 10 months of age (in a region of high endogenous activity produced as evidence of placental function in the transgenic mice). In contrast, no activity was detected in transgenic mice lacking the *hprt* gene, or in transgenic mice lacking the *hprt* gene and the *hprt* gene.

7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844

[illegible]

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific information required.

[illegible][illegible]

Week	10 mg/day	20 mg/day
1	10	10
2	10	10
3	10	10
4	10	10
5	10	10
6	10	10
7	10	10
8	10	10
9	10	10
10	10	10
11	10	10
12	10	10
13	10	10
14	10	10
15	10	10
16	10	10
17	10	10
18	10	10
19	10	10
20	10	10
21	10	10
22	10	10
23	10	10
24	10	10
25	10	10
26	10	10
27	10	10
28	10	10
29	10	10
30	10	10
31	10	10
32	10	10
33	10	10
34	10	10
35	10	10
36	10	10
37	10	10
38	10	10
39	10	10
40	10	10
41	10	10
42	10	10
43	10	10
44	10	10
45	10	10
46	10	10
47	10	10
48	10	10
49	10	10
50	10	10
51	10	10
52	10	10
53	10	10
54	10	10
55	10	10
56	10	10
57	10	10
58	10	10
59	10	10
60	10	10
61	10	10
62	10	10
63	10	10
64	10	10
65	10	10
66	10	10
67	10	10
68	10	10
69	10	10
70	10	10
71	10	10
72	10	10
73	10	10
74	10	10
75	10	10
76	10	10
77	10	10
78	10	10
79	10	10
80	10	10
81	10	10
82	10	10
83	10	10
84	10	10
85	10	10
86	10	10
87	10	10
88	10	10
89	10	10
90	10	10
91	10	10
92	10	10
93	10	10
94	10	10
95	10	10
96	10	10
97	10	10
98	10	10
99	10	10
100	10	10

Reminyl—Cont.

Results may not apply in the ambulatory or non-reporting patients and the type of patients treated may differ. The results of the study are based on the data from the double-blind study. In these patients who experienced the most frequent adverse events, namely, the motion sickness, the results were as follows:

Adverse events occurring with the use of Reminyl in the ambulatory or non-reporting patients are listed in Table 2. The most frequent adverse events, defined as those occurring in 10% or more of the patients, are listed in Table 3. The results of the study are based on the data from the double-blind study. In these patients who experienced the most frequent adverse events, namely, the motion sickness, the results were as follows:

Adverse Event	Reminyl 10 mg/day	Reminyl 20 mg/day
Nausea	24%	23%
Vomiting	18%	10%
Diarrhea	15%	10%
Abdominal pain	15%	10%
Weight decrease	15%	10%

Table 2: The most common adverse events occurring with the use of Reminyl in the ambulatory or non-reporting patients are listed in Table 2. The results of the study are based on the data from the double-blind study. In these patients who experienced the most frequent adverse events, namely, the motion sickness, the results were as follows:

Body System	Reminyl 10 mg/day	Reminyl 20 mg/day
Adverse Event	24%	23%
Nausea	18%	10%
Vomiting	15%	10%
Diarrhea	15%	10%
Abdominal pain	15%	10%
Weight decrease	15%	10%

Reminyl—Cont.

Results may not apply in the ambulatory or non-reporting patients and the type of patients treated may differ. The results of the study are based on the data from the double-blind study. In these patients who experienced the most frequent adverse events, namely, the motion sickness, the results were as follows:

Adverse events occurring with the use of Reminyl in the ambulatory or non-reporting patients are listed in Table 2. The most frequent adverse events, defined as those occurring in 10% or more of the patients, are listed in Table 3. The results of the study are based on the data from the double-blind study. In these patients who experienced the most frequent adverse events, namely, the motion sickness, the results were as follows:

Adverse Event	Reminyl 10 mg/day	Reminyl 20 mg/day
Nausea	24%	23%
Vomiting	18%	10%
Diarrhea	15%	10%
Abdominal pain	15%	10%
Weight decrease	15%	10%

Table 2: The most common adverse events occurring with the use of Reminyl in the ambulatory or non-reporting patients are listed in Table 2. The results of the study are based on the data from the double-blind study. In these patients who experienced the most frequent adverse events, namely, the motion sickness, the results were as follows:

Body System	Reminyl 10 mg/day	Reminyl 20 mg/day
Adverse Event	24%	23%
Nausea	18%	10%
Vomiting	15%	10%
Diarrhea	15%	10%
Abdominal pain	15%	10%
Weight decrease	15%	10%

Reminyl—Cont.

Results may not apply in the ambulatory or non-reporting patients and the type of patients treated may differ. The results of the study are based on the data from the double-blind study. In these patients who experienced the most frequent adverse events, namely, the motion sickness, the results were as follows:

Adverse events occurring with the use of Reminyl in the ambulatory or non-reporting patients are listed in Table 2. The most frequent adverse events, defined as those occurring in 10% or more of the patients, are listed in Table 3. The results of the study are based on the data from the double-blind study. In these patients who experienced the most frequent adverse events, namely, the motion sickness, the results were as follows:

Adverse Event	Reminyl 10 mg/day	Reminyl 20 mg/day
Nausea	24%	23%
Vomiting	18%	10%
Diarrhea	15%	10%
Abdominal pain	15%	10%
Weight decrease	15%	10%

Table 2: The most common adverse events occurring with the use of Reminyl in the ambulatory or non-reporting patients are listed in Table 2. The results of the study are based on the data from the double-blind study. In these patients who experienced the most frequent adverse events, namely, the motion sickness, the results were as follows:

Body System	Reminyl 10 mg/day	Reminyl 20 mg/day
Adverse Event	24%	23%
Nausea	18%	10%
Vomiting	15%	10%
Diarrhea	15%	10%
Abdominal pain	15%	10%
Weight decrease	15%	10%

Montréal—Conti.

[illegible]

1992	Faktor yang tidak terduga di perusahaan	Pengawasan terhadap faktor-faktor tersebut
------	---	--

0197. *Microgaster* (new genus).
 0198. *Microgaster* with a potency similar
 to *Microgaster* and *Microgaster* with a
 potency and blacked almost as
 powerful as *Microgaster* the potency
 in other studies shows almost that
 the powerful inhibition of activity
 and galactinase, or lactase.

ing a complex, familiar with the technical of 19 houses, including ratings. Each town, before, telephone, travel, and household chores, has been built for patients with moderate to severe mental illness, which has a 14,000 to 15,000 square foot footprint.

Figure 6. Cumulative percentage of patients completing 36 months of alcohol-fluox treatment with specified changes from baseline in ALCALC scores.




Figure 6. Cumulative percentage of patients completing 36 months of alcohol-fluox treatment with specified changes from baseline in ALCALC scores.



Figure 3: Time course of the change from baseline in STP score for patients completing all weeks of treatment.



Figure 3: Time course of the change from baseline in SU score for patients completing all weeks of treatment.



Figure 7: Time course of rise-change from baseline in SF-36 score for patients completing all weeks of treatment.

Namondo-Conf

Language treatment was significantly superior to placebo therapy.




Figure 7: Time course of the change from baseline in S7B

[illegible]

הפרטים נמצאים באתר האינטרנט של משרד המשפטים
 תחנת הרכבת תל אביב

A double-blind study of 12-week lamivudine, co-trimoxazole, and zalcitabine in HIV patients with dramatic reduction in plasma HIV RNA levels. The study was carried out in the Department of Medicine, University of Cape Town, South Africa. The study was designed to evaluate the efficacy and safety of lamivudine, co-trimoxazole, and zalcitabine in HIV patients with dramatic reduction in plasma HIV RNA levels. The study was carried out in the Department of Medicine, University of Cape Town, South Africa. The study was designed to evaluate the efficacy and safety of lamivudine, co-trimoxazole, and zalcitabine in HIV patients with dramatic reduction in plasma HIV RNA levels.

[illegible]

INTRACRANIAL HEMORRHAGE
The intracranial hemorrhage (intracerebral) is characterized by the presence of blood within the brain tissue. It is the most common type of stroke, accounting for 25% of all strokes. The hemorrhage is caused by a rupture of a blood vessel within the brain, which can be due to a variety of factors, including high blood pressure, blood vessel disease, and trauma. The hemorrhage can be fatal or cause permanent disability, depending on the location and size of the bleed.

[illegible]

1

MAKEMUKIA and Makindu groups were reported to still remain largely free from malaria, but the incidence of malaria in the Makindu group was generally higher than in the Makemuka group. These authors also noted changes in BCG patterns amongst the Makemuka and Makindu groups. The Makemuka group was vaccinated with BCG at the age of 10 months, of whom more than 1000 were vaccinated. Of 50 samples of BCG strains from vaccinated persons (44 of age 10 and 6 of age 15) 46 were of type 1, 3 were of type 2 and 1 was of type 3. The Makindu group was vaccinated for periods of 4 days to 2 weeks of treatment, and the results were similar to those recorded in the Makemuka group. The Makemuka group was vaccinated with BCG at the age of 10 months, of whom more than 1000 were vaccinated. Of 50 samples of BCG strains from vaccinated persons (44 of age 10 and 6 of age 15) 46 were of type 1, 3 were of type 2 and 1 was of type 3. The Makindu group was vaccinated for periods of 4 days to 2 weeks of treatment, and the results were similar to those recorded in the Makemuka group.

Self System	Adverse Drug	N 1000	%
Self as a Whole			
Feeling		1	
Pain		1	
Intermuscular System			
Myocardium		2	
Cardiac and Pericardial			
Adverse System			
Diagnosis		3	
Headache		3	
Metabolic System			
Cosmetics		3	
Weight		3	
Neurological System			
Back pain		2	
Psychiatric Disorders			
Defecation		5	
Abundance		3	
Abundance		3	
On floor		3	
Organ		1	

[illegible]

14-00000
Several Russian newspapers had been ordered to report on the activities of the patient physicians and neurologists in the USSR, and to indicate the progress in those which are known to cooperate with the YANINA program. The article in the USSR newspaper, "Pravda", was dated 12/10/57. It was modified so that it would not be considered as a recommendation of the YANINA program. The article was dated 12/10/57.

[illegible][illegible][illegible][illegible][illegible][illegible]

TESTALONE is indicated by the trademark with a stylized **T**.

FOREST/129

[illegible]

These studies should be multiplied before drawing conclusions. Pregnancy Category C drugs related to the heart can be considered safe. There is not known with TESSALON and it is not known whether TESSALON can cause fetal harm when administered to a pregnant woman or can affect fertility. TESSALON should be given to women only if clearly needed. There is no known effect of this drug in reproduction. Because use of drugs can be studied in certain subjects that are considered when TESSALON is used to a varying degree.

It is important, important of fertility. Contraception, and reproduction studies have been conducted with TESSALON.

1. Solving and reconfiguring to obtain below has not been established

need to focus specifically related to hormones and other topics and various aspects of their interactions. Drugs of this type are presently well tolerated.

[illegible][illegible]

North 37718
UNIVERSAL, INC.
CONCRETE LABORER, ORDER, INC.
JAN 1964

UNITED STATES has not been systematically compared directly to measurements of using technology from countries such as (C) or other related countries.

following an initial 7-day treatment with 1 capsule dose. Effects on the ADAS-cog, Figure 1 illustrating the time course for the change from baseline in ADAS-cog were for

Following 8 weeks of placebo control, scores on the ADAS-10 for both the ADGC-EPR treatment groups were indistinguishable from those of the control group. The ADAS-10 scores of the ADGC-EPR treatment groups were significantly better than those of the control group at 8 weeks.

Figure 3 illustrates the cumulative percentages of patients from each of the three treatment groups who had attained the measure of improvement in ADAS-seg score shown on the X axis. There change scores, (7-point and 4-point reduction from baseline) as the chosen 10 percent have been identified.

1940 Bureau of Power
1022

it should be the right of the owner for the plant, which is
respective.

[illegible]



specific. These differences were statistically significant. There was no statistically significant difference between the two active treatments.

[See figure at top of next column.]

Posttest-Pretest Study

In a total of 15 weeks 4,000 patients were randomized


all Qwen dogs groups over the 18 weeks of the study. After 18 weeks of treatment, the differences in mean ADAS-cog scores for the ABX227PM treated patients compared to the patients on placebo were 2.7 and 3.0 units each, for



It is a wide range of responses, but that the ABCs of good judgment are more likely to show the greater awareness to the capital performance.



Age Group	Percentage
18-24	10
25-34	15
35-44	25
45-54	35
55-64	30
65-74	20
75-84	10
85+	5

[illegible]

and the fact that the average age of the patients was 37 years, the authors concluded that the disease was not a result of aging. The authors also noted that the disease was not associated with any other systemic disease, and that the disease was not associated with any other organ system. The authors also noted that the disease was not associated with any other organ system.

[illegible][illegible]

Yeast of **ALBUCEPT** on the Monod model. The growth rate μ is a function of the substrate concentration S and the product concentration P . The Monod model is given by the following equation:

$$\mu = \frac{\mu_{\max} S}{K_s + S + \frac{K_p P}{K_i}}$$

where μ_{\max} is the maximum growth rate, K_s is the half-saturation constant, K_i is the inhibition constant, and K_p is the product inhibition constant. The Monod model is a simple model that can be used to describe the growth of yeast in a batch culture. However, it is not a perfect model and it may not be able to describe all aspects of yeast growth. For example, the Monod model does not take into account the effect of pH on yeast growth. In a batch culture, the pH of the medium can change over time, which can affect the growth rate of the yeast. Therefore, a more complex model that takes into account the effect of pH on yeast growth may be more appropriate for describing yeast growth in a batch culture.

performing digital and mechanical manipulations on the patient's fingers. The effect of other drugs on the blood circulation and gelation inhibition of DNA compounds in blood vessels.

in the Acanthamoeba castellanii, a known pathogen of immunocompromised humans, and in the Acanthamoeba polyphaga, a known pathogen of fish. The Acanthamoeba castellanii is a free-living, unicellular organism that can cause a variety of diseases in humans, including amoebic keratitis, a serious eye infection. The Acanthamoeba polyphaga is a free-living, unicellular organism that can cause a variety of diseases in fish, including gill disease and skin disease. The Acanthamoeba castellanii is a free-living, unicellular organism that can cause a variety of diseases in humans, including amoebic keratitis, a serious eye infection. The Acanthamoeba polyphaga is a free-living, unicellular organism that can cause a variety of diseases in fish, including gill disease and skin disease.

1. **NAME** _____
 2. **ADDRESS** _____
 3. **CITY** _____
 4. **STATE** _____
 5. **ZIP** _____
 6. **PHONE** _____
 7. **DATE** _____
 8. **SIGNATURE** _____
 9. **PRINT NAME** _____
 10. **PRINT ADDRESS** _____
 11. **PRINT CITY** _____
 12. **PRINT STATE** _____
 13. **PRINT ZIP** _____
 14. **PRINT PHONE** _____
 15. **PRINT DATE** _____
 16. **PRINT SIGNATURE** _____
 17. **PRINT NAME** _____
 18. **PRINT ADDRESS** _____
 19. **PRINT CITY** _____
 20. **PRINT STATE** _____
 21. **PRINT ZIP** _____
 22. **PRINT PHONE** _____
 23. **PRINT DATE** _____
 24. **PRINT SIGNATURE** _____
 25. **PRINT NAME** _____
 26. **PRINT ADDRESS** _____
 27. **PRINT CITY** _____
 28. **PRINT STATE** _____
 29. **PRINT ZIP** _____
 30. **PRINT PHONE** _____
 31. **PRINT DATE** _____
 32. **PRINT SIGNATURE** _____
 33. **PRINT NAME** _____
 34. **PRINT ADDRESS** _____
 35. **PRINT CITY** _____
 36. **PRINT STATE** _____
 37. **PRINT ZIP** _____
 38. **PRINT PHONE** _____
 39. **PRINT DATE** _____
 40. **PRINT SIGNATURE** _____
 41. **PRINT NAME** _____
 42. **PRINT ADDRESS** _____
 43. **PRINT CITY** _____
 44. **PRINT STATE** _____
 45. **PRINT ZIP** _____
 46. **PRINT PHONE** _____
 47. **PRINT DATE** _____
 48. **PRINT SIGNATURE** _____
 49. **PRINT NAME** _____
 50. **PRINT ADDRESS** _____
 51. **PRINT CITY** _____
 52. **PRINT STATE** _____
 53. **PRINT ZIP** _____
 54. **PRINT PHONE** _____
 55. **PRINT DATE** _____
 56. **PRINT SIGNATURE** _____
 57. **PRINT NAME** _____
 58. **PRINT ADDRESS** _____
 59. **PRINT CITY** _____
 60. **PRINT STATE** _____
 61. **PRINT ZIP** _____
 62. **PRINT PHONE** _____
 63. **PRINT DATE** _____
 64. **PRINT SIGNATURE** _____
 65. **PRINT NAME** _____
 66. **PRINT ADDRESS** _____
 67. **PRINT CITY** _____
 68. **PRINT STATE** _____
 69. **PRINT ZIP** _____
 70. **PRINT PHONE** _____
 71. **PRINT DATE** _____
 72. **PRINT SIGNATURE** _____
 73. **PRINT NAME** _____
 74. **PRINT ADDRESS** _____
 75. **PRINT CITY** _____
 76. **PRINT STATE** _____
 77. **PRINT ZIP** _____
 78. **PRINT PHONE** _____
 79. **PRINT DATE** _____
 80. **PRINT SIGNATURE** _____
 81. **PRINT NAME** _____
 82. **PRINT ADDRESS** _____
 83. **PRINT CITY** _____
 84. **PRINT STATE** _____
 85. **PRINT ZIP** _____
 86. **PRINT PHONE** _____
 87. **PRINT DATE** _____
 88. **PRINT SIGNATURE** _____
 89. **PRINT NAME** _____
 90. **PRINT ADDRESS** _____
 91. **PRINT CITY** _____
 92. **PRINT STATE** _____
 93. **PRINT ZIP** _____
 94. **PRINT PHONE** _____
 95. **PRINT DATE** _____
 96. **PRINT SIGNATURE** _____
 97. **PRINT NAME** _____
 98. **PRINT ADDRESS** _____
 99. **PRINT CITY** _____
 100. **PRINT STATE** _____
 101. **PRINT ZIP** _____
 102. **PRINT PHONE** _____
 103. **PRINT DATE** _____
 104. **PRINT SIGNATURE** _____
 105. **PRINT NAME** _____
 106. **PRINT ADDRESS** _____
 107. **PRINT CITY** _____
 108. **PRINT STATE** _____
 109. **PRINT ZIP** _____
 110. **PRINT PHONE** _____
 111. **PRINT DATE** _____
 112. **PRINT SIGNATURE** _____
 113. **PRINT NAME** _____
 114. **PRINT ADDRESS** _____
 115. **PRINT CITY** _____
 116. **PRINT STATE** _____
 117. **PRINT ZIP** _____
 118. **PRINT PHONE** _____
 119. **PRINT DATE** _____
 120. **PRINT SIGNATURE** _____
 121. **PRINT NAME** _____
 122. **PRINT ADDRESS** _____
 123. **PRINT CITY** _____
 124. **PRINT STATE** _____
 125. **PRINT ZIP** _____
 126. **PRINT PHONE** _____
 127. **PRINT DATE** _____
 128. **PRINT SIGNATURE** _____
 129. **PRINT NAME** _____
 130. **PRINT ADDRESS** _____
 131. **PRINT CITY** _____
 132. **PRINT STATE** _____
 133. **PRINT ZIP** _____
 134. **PRINT PHONE** _____
 135. **PRINT DATE** _____
 136. **PRINT SIGNATURE** _____
 137. **PRINT NAME** _____
 138. **PRINT ADDRESS** _____
 139. **PRINT CITY** _____
 140. **PRINT STATE** _____
 141. **PRINT ZIP** _____
 142. **PRINT PHONE** _____
 143. **PRINT DATE** _____
 144. **PRINT SIGNATURE** _____
 145. **PRINT NAME** _____
 146. **PRINT ADDRESS** _____
 147. **PRINT CITY** _____
 148. **PRINT STATE** _____
 149. **PRINT ZIP** _____
 150. **PRINT PHONE** _____
 151. **PRINT DATE** _____
 152. **PRINT SIGNATURE** _____
 153. **PRINT NAME** _____
 154. **PRINT ADDRESS** _____
 155. **PRINT CITY** _____
 156. **PRINT STATE** _____
 157. **PRINT ZIP** _____
 158. **PRINT PHONE** _____
 159. **PRINT DATE** _____
 160. **PRINT SIGNATURE** _____
 161. **PRINT NAME** _____
 162. **PRINT ADDRESS** _____
 163. **PRINT CITY** _____
 164. **PRINT STATE** _____
 165. **PRINT ZIP** _____
 166. **PRINT PHONE** _____
 167. **PRINT DATE** _____
 168. **PRINT SIGNATURE** _____
 169. **PRINT NAME** _____
 170. **PRINT ADDRESS** _____
 171. **PRINT CITY** _____
 172. **PRINT STATE** _____
 173. **PRINT ZIP** _____
 174. **PRINT PHONE** _____
 175. **PRINT DATE** _____
 176. **PRINT SIGNATURE** _____
 177. **PRINT NAME** _____
 178. **PRINT ADDRESS** _____
 179. **PRINT CITY** _____
 180. **PRINT STATE** _____
 181. **PRINT ZIP** _____
 182. **PRINT PHONE** _____
 183. **PRINT DATE** _____
 184. **PRINT SIGNATURE** _____
 185. **PRINT NAME** _____
 186. **PRINT ADDRESS** _____
 187. **PRINT CITY** _____
 188. **PRINT STATE** _____
 189. **PRINT ZIP** _____
 190. **PRINT PHONE** _____
 191. **PRINT DATE** _____
 192. **PRINT SIGNATURE** _____
 193. **PRINT NAME** _____
 194. **PRINT ADDRESS** _____
 195. **PRINT CITY** _____
 196. **PRINT STATE** _____
 197. **PRINT ZIP** _____
 198. **PRINT PHONE** _____
 199. **PRINT DATE** _____
 200. **PRINT SIGNATURE** _____
 201. **PRINT NAME** _____
 202. **PRINT ADDRESS** _____
 203. **PRINT CITY** _____
 204. **PRINT STATE** _____
 205. **PRINT ZIP** _____
 206. **PRINT PHONE** _____
 207. **PRINT DATE** _____
 208. **PRINT SIGNATURE** _____
 209. **PRINT NAME** _____
 210. **PRINT ADDRESS** _____
 211. **PRINT CITY** _____
 212. **PRINT STATE** _____
 213. **PRINT ZIP** _____
 214. **PRINT PHONE** _____
 215. **PRINT DATE** _____
 216. **PRINT SIGNATURE** _____
 217. **PRINT NAME** _____
 218. **PRINT ADDRESS** _____
 219. **PRINT CITY** _____
 220. **PRINT STATE** _____

[illegible]

ing Mothers
not known whether deaths is reported in American
and mtd. ARIUNST/2000 no indication for use in nursing
bars.
decide Upon
are an adequate and well-controlled study to deter-

of these patients were between 65 and 84 years old, and of patients were at or above the age of 75. The efficacy safety data presented in the clinical trials sections were based from these patients. There were no clinically significant differences in most adverse events reported by groups in a 60 years old and < 65 years old.

tion of patients who received 7-day treatment from Day 1 to Day 7 was higher at 13%.

DATE	DESCRIPTION	AMOUNT	BALANCE
1971	1000	1000	1000
1972	1000	1000	1000
1973	1000	1000	1000
1974	1000	1000	1000
1975	1000	1000	1000
1976	1000	1000	1000
1977	1000	1000	1000
1978	1000	1000	1000
1979	1000	1000	1000
1980	1000	1000	1000
1981	1000	1000	1000
1982	1000	1000	1000
1983	1000	1000	1000
1984	1000	1000	1000
1985	1000	1000	1000
1986	1000	1000	1000
1987	1000	1000	1000
1988	1000	1000	1000
1989	1000	1000	1000
1990	1000	1000	1000
1991	1000	1000	1000
1992	1000	1000	1000
1993	1000	1000	1000
1994	1000	1000	1000
1995	1000	1000	1000
1996	1000	1000	1000
1997	1000	1000	1000
1998	1000	1000	1000
1999	1000	1000	1000
2000	1000	1000	1000
2001	1000	1000	1000
2002	1000	1000	1000
2003	1000	1000	1000
2004	1000	1000	1000
2005	1000	1000	1000
2006	1000	1000	1000
2007	1000	1000	1000
2008	1000	1000	1000
2009	1000	1000	1000
2010	1000	1000	1000
2011	1000	1000	1000
2012	1000	1000	1000
2013	1000	1000	1000
2014	1000	1000	1000
2015	1000	1000	1000
2016	1000	1000	1000
2017	1000	1000	1000
2018	1000	1000	1000
2019	1000	1000	1000
2020	1000	1000	1000
2021	1000	1000	1000
2022	1000	1000	1000
2023	1000	1000	1000
2024	1000	1000	1000
2025	1000	1000	1000
2026	1000	1000	1000
2027	1000	1000	1000
2028	1000	1000	1000
2029	1000	1000	1000
2030	1000	1000	1000
2031	1000	1000	1000
2032	1000	1000	1000
2033	1000	1000	1000
2034	1000	1000	1000
2035	1000	1000	1000
2036	1000	1000	1000
2037	1000	1000	1000
2038	1000	1000	1000
2039	1000	1000	1000
2040	1000	1000	1000
2041	1000	1000	1000
2042	1000	1000	1000
2043	1000	1000	1000
2044	1000	1000	1000
2045	1000	1000	1000
2046	1000	1000	1000
2047	1000	1000	1000
2048	1000	1000	1000
2049	1000	1000	1000
2050	1000	1000	1000
2051	1000	1000	1000
2052	1000	1000	1000
2053	1000	1000	1000
2054	1000	1000	1000
2055	1000	1000	1000
2056	1000	1000	1000

Freeman Adopts Clinical Events Seen in Association
the Use of ACCEPT®

is evidence to suggest that the frequency of these no adverse events may be affected by the rate of fibrinogen. In open-label study was conducted with 289 patients assigned placebo in the 13 and 20-week studies. These patients were divided to a dose of 10 mg/day over a 5-week period. The rates of adverse events were lower

	No titration	One week titration	Six weeks titration
--	--------------	--------------------	---------------------

1962	62	65	114	65
1963	33	68	86	78
1964	29	74	69	52

馬	馬
馬	馬
馬	馬
馬	馬

[illegible]

이러한 사실은, 이념적 대립을 초월하여, 양측이 공동의 이익을 위하여 협력할 수 있는 여지를 열어준다.

14 15

Case	Year	Age
------	------	-----

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

TV	11	12
MI	13	14
15	16	17

10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

... ..

THE

Abstract

The results of the earlier experiments pointed toward a highly selected group of clinical birds in a highly selected population. In actual clinical practice or in other medical fields, these respiratory infections may not apply, as the conditions of the birds and the kinds of particles involved may differ. With a fully developed immune system and a compromised host, the rate of infection was found to be 100 percent. The results of the experiments with ABC-3750 assigned those phages and used phages, in general, where they occurred more frequently in French patients and with attacking age.

Adverse Event	Body as a Whole				
	D	P	JN		
Headache	5	5	5		
Pain, various locations	5	5	5		
Arthralgia	5	5	5		
Fatigue	5	5	5		

Fatigue	3	5
Cardiovascular System		
Stomach	3	3
Urinary System		

Yearling	2	3
Austria	2	4

Hande and Symbolic System

Experiments	3	4
-------------	---	---

Handa Group	2	5
Artistic	1	2
Harmon Systems		
Intense's	4	9

Abundance (amu)	D	1
See below	<1	2

Unresolved Spectrum	1	2
From mass 111.0000		

over 2,000 patients have been treated for at least 6 months. Controlled and uncontrolled trials in the United States included approximately 800 patients. In regards to the highest dose of 40 mg/day, this population includes 63 patients treated for 3 months, 418 patients treated for 6 months and 155 patients treated for over 1 year. The majority of patients were treated for 1 to 12 days.

[illegible]

The results of the earlier experiments pointed toward a highly selected group of patients with a high percentage of involved conditions of clinical order in a highly selected patient population. In actual clinical practice or in other medical fields, these respiratory ailments may not apply, as the conditions of the patients may not be the same as the conditions of our laboratory patients and the kinds of particles inhaled may differ. With a wide treatment curve, the use of patients in placebo-controlled trials who received various types and compositions that were prepared in at least 30 different ways, the results of the present study are representative for ABC-3770 assigned those placebo and fixed dose groups. In general, adverse events occurred more frequently in French patients and with activating agents.

Study System/Adverse Event	Placebo (n=253)	ARCEPT (n=247)
Percent of Patients with any	22	34

Adverse Event		
Body as a Whole		
Headache	6	10
Pain various locations	8	9
Accidents	6	1

Fatigue	3	5
Constraining System		
Stressor	3	3
Divergent System		
Nuisance	9	11
Distress	5	10

	1	2	3	4
Young		2		
Adult				
Hemid and Hemipha 2 types				
Embryonic		3		4
Metabolic and Nutritional Systems				
Weight Decrease		1		3
Hemodermal System				

Hande Gruppe	1	2	3
Arthritis			
Reizarm System			
Ischias/S		4	9
Encephal		5	8
Degenerat		<1	7

Abnormal Finding	D	I
Stool color	<1	2
Urogenital System		
Frequency/Urination	1	2

There were 1060 patients who have been treated for at least one month. Of these, 100 patients were treated for 1 to 3 months. Controlled and uncontrolled trials in the United States included approximately 800 patients. In regard to the highest dose of 80 mg/day, this population includes 63 patients treated for 30 mg/day, 11 patients treated for 40 mg/day, and 6 patients treated for 80 mg/day. The success rate for patients treated for 1 year. The success rate for patients treated for 1 to 3 months is 100%.

ing. It provides an arbitrary estimate of the physical volume of events, the events were divided into two categories: (1) events having a small number of standardized categories grouped into a smaller number of standardized categories using a modified CONCEPT dictionary and event descriptions were calculated across all studies. These categories are used in the following tables. The hypotheses represent the proportion of 500 patients from these trials who experienced an event. The event was defined as a patient who had at least one event recorded in the event dictionary. All events were counted at least twice and included, except for those in

[illegible]

as a predictable consequence of its pharmacological activity. In the absence of evidence to the contrary, it is probable that the same mechanism is operative in the case of the other drugs in this class. These effects, when they occur, appear to be related to the high plasma levels with which the drugs are associated. In the case of the other drugs, these effects have been noted only in some cases. In the case of the other drugs, these effects have been noted only in some cases. In the case of the other drugs, these effects have been noted only in some cases.

of the Committee. Scientific conclusions are not to be drawn from the fact that the Committee has not yet been able to establish a causal relationship between the use of asbestos and the occurrence of cancer. The Committee also may be a consideration of the fact that the use of asbestos is not a necessary condition for the occurrence of cancer. Because of these considerations, the Committee believes that the use of asbestos should be restricted to those cases in which it is necessary for the protection of the health of the worker.

Pharmacokinetic Clinical Pharmacology—Cefazolin sodium (Nagging International)

Precedence studies evaluated the potential of the interaction with theophylline, considering the effects of drugs and beverages. No effects of these drugs were observed.

and to control the use of the drug. Because of their mechanism of action, these drugs are not significantly affected by co-administration of alcohol or other CNS depressants. Because of their mechanism of action, these drugs are not significantly affected by co-administration of alcohol or other CNS depressants.

[illegible][illegible][illegible]

therefore, considerable improvement both in the incidence of generalized tonic-clonic seizures and in the quality of life in patients with a diagnosis of symptomatic localization-related epilepsy. The effectiveness of 200 mg/day of lamotrigine (added to either carbamazepine or phenytoin) in controlling seizures was comparable to that of 200 mg/day of carbamazepine or phenytoin.

[illegible]

The secondary structure was probed by CD. The CD spectra of the wild-type and mutant proteins were recorded in 10 mM sodium phosphate buffer at pH 7.0. The CD spectra were recorded at 25°C. The results described below are for the wild-type protein.

[illegible]

...the ... of ...

NaOH to saponification, resulting in an emulsion of approximately 10 and 100 μm particles. The phosphatidylcholine was added to the emulsion in the form of a 10% solution in the range 100–120 mg/kg body weight. The emulsion was then stirred until binding of phospholipids to RNA was complete, usually over a 2-h period. The resulting emulsion was collected in a 100-ml beaker and the nucleic acids were precipitated by the addition of 100 ml of 95% ethanol. The phospholipid nucleic acid complexes were dried in a vacuum oven at 40°C for 24 h.

The frequent values of 2.0 mg/kg and about 1.45 mg/kg following 2.0 and 1.45 mg/kg concentrations of 1,6-3.0 $\mu\text{g/kg}$ are not far from the values obtained by the method used to determine plasma protein.

strong ZONIDAMIC for doses of 500 mg was not apparent (differences were not significant). The mean plasma concentrations (last 4 weeks) during chronic treatment with ZONIDAMIC 500 mg were significantly different from placebo (Table 1). In subjects 129 and 130 no side effects were observed. The first side effect was dry eyes after 2 weeks' treatment.

Proportions of patients with side effects in the 2 studies in the 1st and 2nd periods are shown in Table 2. There was no difference between the 2 periods in the proportions of patients with side effects. The proportions of patients with side effects were similar in both studies. The proportions of patients with side effects were similar in both studies.

of patients exhibiting any particular side effect was not statistically higher for the placebo group than for the active groups. The mean (SD) approximately 21% of patients in the placebo group were unable to tolerate the treatment, compared to approximately 20% in the active groups.

2.34 solution, respectively. The pharmacokinetic changes < 10 and > 10 fold in the pharmacokinetic parameters in the elimination phase were defined as significant changes.

Results

Pharmacokinetics The pharmacokinetic changes in the elimination phase in patients with impaired liver function are shown in Figure 1. The study included 100 patients with impaired liver function (Table 1).

Age The pharmacokinetic parameters in patients with impaired liver function are shown in Table 2. The mean age was 60 years.

Gender and Race The pharmacokinetic parameters in patients with impaired liver function are shown in Table 3. The mean race was 60 years.

Conclusion The pharmacokinetic parameters in patients with impaired liver function are shown in Table 4. The mean race was 60 years.

[illegible]

Two related types of toxicity in animals subjected to long-term treatment with 10 mg/kg of *ALOXEPTIN* were: (1) a progressive wasting syndrome, with emaciation, staggering gait, and depression; and (2) a severe depression of the immune system, with decreased resistance to bacterial infection, delayed wound healing, and decreased resistance to tumor induction. These effects were not observed in animals treated with 10 mg/kg of *ALOXEPTIN* for 10 days.

[illegible][illegible][illegible]

ZONERSHANE
[patented]
conditioner
for hair

O=C1C(=O)c2ccccc2N1

The chemical structure is:

O=C1C(=O)c2ccccc2N1[illegible]

100


100

1000

100

100

100



logical outcomes, 47% of 2000 patients and 49% of 1416 patients used corticosteroids during the study. Over 2000 patients were used for more than 7 days. 74.1% of 2000 patients and 64.9% of 1416 patients. An increased incidence of hospital admission, infection, superinfection (unusual single

Predictive Performance (Vehicle-Occupied (10 Years))		Predictive Performance (Vehicle-Occupied (1 Year))		Adult Active Driver/Operator (1 Year)	
Ensemble	Vehicle (R=0.82) MUS	Ensemble	Vehicle (R=0.72) (R=0.74)	Ensemble	Green (R=0.79) (R=0.79)
Ensemble	0.82	Ensemble	0.72	Ensemble	0.79

12-16-60

[illegible][illegible][illegible][illegible]

the Zurich trade was a moderate figure, representing an indication of a decline in the domestic political and economic situation, including a cessation of being. It represents the movement of the Zurich trade was based on high or low, and the Zurich trade was based on high or low.

O=C1C(=O)C(=O)C(=O)C1

1999
 1998
 1997
 1996
 1995
 1994
 1993
 1992
 1991
 1990
 1989
 1988
 1987
 1986
 1985
 1984
 1983
 1982
 1981
 1980
 1979
 1978
 1977
 1976
 1975
 1974
 1973
 1972
 1971
 1970
 1969
 1968
 1967
 1966
 1965
 1964
 1963
 1962
 1961
 1960
 1959
 1958
 1957
 1956
 1955
 1954
 1953
 1952
 1951
 1950
 1949
 1948
 1947
 1946
 1945
 1944
 1943
 1942
 1941
 1940
 1939
 1938
 1937
 1936
 1935
 1934
 1933
 1932
 1931
 1930
 1929
 1928
 1927
 1926
 1925
 1924
 1923
 1922
 1921
 1920
 1919
 1918
 1917
 1916
 1915
 1914
 1913
 1912
 1911
 1910
 1909
 1908
 1907
 1906
 1905
 1904
 1903
 1902
 1901
 1900
 1899
 1898
 1897
 1896
 1895
 1894
 1893
 1892
 1891
 1890
 1889
 1888
 1887
 1886
 1885
 1884
 1883
 1882
 1881
 1880
 1879
 1878
 1877
 1876
 1875
 1874
 1873
 1872
 1871
 1870
 1869
 1868
 1867
 1866
 1865
 1864
 1863
 1862
 1861
 1860
 1859
 1858
 1857
 1856
 1855
 1854
 1853
 1852
 1851
 1850
 1849
 1848
 1847
 1846
 1845
 1844
 1843
 1842
 1841
 1840
 1839
 1838
 1837
 1836
 1835
 1834
 1833
 1832
 1831
 1830
 1829
 1828
 1827
 1826
 1825
 1824
 1823
 1822
 1821
 1820
 1819
 1818
 1817
 1816
 1815
 1814
 1813
 1812
 1811
 1810
 1809
 1808
 1807
 1806
 1805
 1804
 1803
 1802
 1801
 1800
 1799
 1798
 1797
 1796
 1795
 1794
 1793
 1792
 1791
 1790
 1789
 1788
 1787
 1786
 1785
 1784
 1783
 1782
 1781
 1780
 1779
 1778
 1777
 1776
 1775
 1774
 1773
 1772
 1771
 1770
 1769
 1768
 1767
 1766
 1765
 1764
 1763
 1762
 1761
 1760
 1759
 1758
 1757
 1756
 1755
 1754
 1753
 1752
 1751
 1750
 1749
 1748
 1747
 1746
 1745
 1744
 1743
 1742
 1741
 1740
 1739
 1738
 1737
 1736
 1735
 1734
 1733
 1732
 1731
 1730
 1729
 1728
 1727
 1726
 1725
 1724
 1723
 1722
 1721
 1720
 1719
 1718
 1717
 1716
 1715
 1714
 1713
 1712
 1711
 1710
 1709
 1708
 1707
 1706
 1705
 1704
 1703
 1702
 1701
 1700
 1699
 1698
 1697
 1696
 1695
 1694
 1693
 1692
 1691
 1690
 1689
 1688
 1687
 1686
 1685
 1684
 1683
 1682
 1681
 1680
 1679
 1678
 1677
 1676
 1675
 1674
 1673
 1672
 1671
 1670
 1669
 1668
 1667
 1666
 1665
 1664
 1663
 1662
 1661
 1660
 1659
 1658
 1657
 1656
 1655
 1654
 1653
 1652
 1651
 1650
 1649
 1648
 1647
 1646
 1645
 1644
 1643
 1642
 1641
 1640
 1639
 1638
 1637
 1636
 1635
 1634
 1633
 1632
 1631
 1630
 1629
 1628
 1627
 1626
 1625
 1624
 1623
 1622
 1621
 1620
 1619
 1618
 1617
 1616
 1615
 1614
 1613
 1612
 1611
 1610
 1609
 1608
 1607
 1606
 1605
 1604
 1603
 1602
 1601
 1600
 1599
 1598
 1597
 1596
 1595
 1594
 1593
 1592
 1591
 1590
 1589
 1588
 1587
 1586
 1585
 1584
 1583
 1582
 1581
 1580
 1579
 1578
 1577
 1576
 1575
 1574
 1573
 1572
 1571
 1570
 1569
 1568
 1567
 1566
 1565
 1564
 1563
 1562
 1561
 1560
 1559
 1558
 1557
 1556
 1555
 1554
 1553
 1552
 1551
 1550
 1549
 1548

[illegible]

businessmen, and the percent of respondents who had attended of representatives to AIDS-seminars. These change scores (T-scores) were used as an index to control for confounding variables in the regression equations, and the percent of respondents who had attended of representatives to AIDS-seminars.

are in severe cases. Numerous authors have reported that the severity of the disease is related to the degree of the infection. The more severe the infection, the more severe the disease. The following table summarizes the results of the studies conducted by the authors.

Study	Severity of Infection	Severity of Disease
1	Severe	Severe
2	Severe	Severe
3	Severe	Severe
4	Severe	Severe
5	Severe	Severe
6	Severe	Severe
7	Severe	Severe
8	Severe	Severe
9	Severe	Severe
10	Severe	Severe

These results show that the severity of the disease is directly related to the severity of the infection. The more severe the infection, the more severe the disease. This is a very important finding, as it suggests that the severity of the disease can be predicted by the severity of the infection. This information is crucial for the development of effective treatments and for the management of the disease.

DYED-COLOR

There has been no experience of people with this disease. The disease is very rare and is not well understood. The authors have conducted a study of the disease and have found that it is caused by a virus. The virus is very small and is not visible to the naked eye. The virus is very contagious and can be spread from person to person. The disease is very serious and can be fatal. The authors have found that the disease is caused by a virus that is very similar to the virus that causes the disease in other animals. This finding is very important, as it suggests that the disease in humans is caused by the same virus as the disease in other animals. This information is crucial for the development of effective treatments and for the management of the disease.

the head, neck, and intervertebral discs. The use of a Sibley-Grimm neck brace is recommended. Treatment should be directed at the elimination of systemic toxicity. If systemic toxicity, the patient should be rechecked. The use of Sibley-Grimm neck braces, neck systemic exposure, has not been reported.

[illegible][illegible]

The authors thank the
 for the generous
 and constructive criticism
 of this manuscript.
 and the authors thank
 for the generous
 and constructive criticism
 of this manuscript.
 and the authors thank
 for the generous
 and constructive criticism
 of this manuscript.

O=C1C(=O)C(=O)C(=O)C1

IDENTICAL FEARS OUT

These changes concern
the selection of the
factory program, and

the 1960s. The 1960s saw the rise of the "New Wave" of science fiction, which was characterized by a more realistic and often darker approach to the genre. This was a time when science fiction began to be taken more seriously as a literary form, and it was no longer just a genre for children or teenagers. The 1960s also saw the rise of the "New Wave" of science fiction, which was characterized by a more realistic and often darker approach to the genre. This was a time when science fiction began to be taken more seriously as a literary form, and it was no longer just a genre for children or teenagers.

[illegible][illegible][illegible][illegible]O=C1C(=O)C(=O)C(=O)C1

1999
 1998
 1997
 1996
 1995
 1994
 1993
 1992
 1991
 1990
 1989
 1988
 1987
 1986
 1985
 1984
 1983
 1982
 1981
 1980
 1979
 1978
 1977
 1976
 1975
 1974
 1973
 1972
 1971
 1970
 1969
 1968
 1967
 1966
 1965
 1964
 1963
 1962
 1961
 1960
 1959
 1958
 1957
 1956
 1955
 1954
 1953
 1952
 1951
 1950
 1949
 1948
 1947
 1946
 1945
 1944
 1943
 1942
 1941
 1940
 1939
 1938
 1937
 1936
 1935
 1934
 1933
 1932
 1931
 1930
 1929
 1928
 1927
 1926
 1925
 1924
 1923
 1922
 1921
 1920
 1919
 1918
 1917
 1916
 1915
 1914
 1913
 1912
 1911
 1910
 1909
 1908
 1907
 1906
 1905
 1904
 1903
 1902
 1901
 1900
 1899
 1898
 1897
 1896
 1895
 1894
 1893
 1892
 1891
 1890
 1889
 1888
 1887
 1886
 1885
 1884
 1883
 1882
 1881
 1880
 1879
 1878
 1877
 1876
 1875
 1874
 1873
 1872
 1871
 1870
 1869
 1868
 1867
 1866
 1865
 1864
 1863
 1862
 1861
 1860
 1859
 1858
 1857
 1856
 1855
 1854
 1853
 1852
 1851
 1850
 1849
 1848
 1847
 1846
 1845
 1844
 1843
 1842
 1841
 1840
 1839
 1838
 1837
 1836
 1835
 1834
 1833
 1832
 1831
 1830
 1829
 1828
 1827
 1826
 1825
 1824
 1823
 1822
 1821
 1820
 1819
 1818
 1817
 1816
 1815
 1814
 1813
 1812
 1811
 1810
 1809
 1808
 1807
 1806
 1805
 1804
 1803
 1802
 1801
 1800
 1799
 1798
 1797
 1796
 1795
 1794
 1793
 1792
 1791
 1790
 1789
 1788
 1787
 1786
 1785
 1784
 1783
 1782
 1781
 1780
 1779
 1778
 1777
 1776
 1775
 1774
 1773
 1772
 1771
 1770
 1769
 1768
 1767
 1766
 1765
 1764
 1763
 1762
 1761
 1760
 1759
 1758
 1757
 1756
 1755
 1754
 1753
 1752
 1751
 1750
 1749
 1748
 1747
 1746
 1745
 1744
 1743
 1742
 1741
 1740
 1739
 1738
 1737
 1736
 1735
 1734
 1733
 1732
 1731
 1730
 1729
 1728
 1727
 1726
 1725
 1724
 1723
 1722
 1721
 1720
 1719
 1718
 1717
 1716
 1715
 1714
 1713
 1712
 1711
 1710
 1709
 1708
 1707
 1706
 1705
 1704
 1703
 1702
 1701
 1700
 1699
 1698
 1697
 1696
 1695
 1694
 1693
 1692
 1691
 1690
 1689
 1688
 1687
 1686
 1685
 1684
 1683
 1682
 1681
 1680
 1679
 1678
 1677
 1676
 1675
 1674
 1673
 1672
 1671
 1670
 1669
 1668
 1667
 1666
 1665
 1664
 1663
 1662
 1661
 1660
 1659
 1658
 1657
 1656
 1655
 1654
 1653
 1652
 1651
 1650
 1649
 1648
 1647
 1646
 1645
 1644
 1643
 1642
 1641
 1640
 1639
 1638
 1637
 1636
 1635
 1634
 1633
 1632
 1631
 1630
 1629
 1628
 1627
 1626
 1625
 1624
 1623
 1622
 1621
 1620
 1619
 1618
 1617
 1616
 1615
 1614
 1613
 1612
 1611
 1610
 1609
 1608
 1607
 1606
 1605
 1604
 1603
 1602
 1601
 1600
 1599
 1598
 1597
 1596
 1595
 1594
 1593
 1592
 1591
 1590
 1589
 1588
 1587
 1586
 1585
 1584
 1583
 1582
 1581
 1580
 1579
 1578
 1577
 1576
 1575
 1574
 1573
 1572
 1571
 1570
 1569
 1568
 1567
 1566
 1565
 1564
 1563
 1562
 1561
 1560
 1559
 1558
 1557
 1556
 1555
 1554
 1553
 1552
 1551
 1550
 1549
 1548

[illegible]

businessmen, and the percent of respondents who had attended of representatives to AIDS-seminars. These change scores (T-scores) were used as an index to control for confounding variables in the regression equations, and the percent of respondents who had attended of representatives to AIDS-seminars.

Fig. 2. Relationship of the degree of reduction in the number of bacteria in the culture to the time of treatment.

Time of treatment (min)	100°C (circles)	100°C (triangles)	100°C (squares)	100°C (diamonds)
0	0	0	0	0
2	0.5	0.3	0.2	0.1
4	1.0	0.6	0.4	0.2
6	1.5	0.9	0.6	0.3
8	2.0	1.2	0.8	0.4
10	2.5	1.5	1.0	0.5

[illegible][illegible]

Figure 2. Frequency Distribution of CRIC/CAI Scores at Month 20

Figure 1. The effect of the treatment on the growth of the plants. The plants were grown in a glasshouse under long-day conditions (16 h light/8 h dark) at 20°C. The plants were harvested at the end of the experiment (10 weeks after sowing). The data are the mean of three replicates. The error bars represent the standard deviation. The different letters indicate significant differences (p < 0.05) according to the Tukey test.

[illegible]

Two were 1.9 and 4.3 cells per field at 1 and 2 weeks, respectively. Both treatments were significantly superior to placebo ($p < 0.05$) and were significantly superior to the two control groups (two-sided). The mean number of cells per field was significantly greater in the two treatment groups than in the two control groups (two-sided).

of Intervention Group who had returned to work after a period of absence. The chart shows that change scores (Y-axis) tend to be positive, indicating that scores have changed in a positive direction. The legend indicates that the Y-axis represents the percentage of the total sample. The X-axis represents the percentage of the total sample. The chart shows that the percentage of the total sample who returned to work after a period of absence is approximately 10%.

[illegible]

There is a considerable literature on the use of the EKG in the diagnosis of myocardial infarction. The EKG is a valuable tool in the diagnosis of myocardial infarction, but it is not always reliable. The EKG is a valuable tool in the diagnosis of myocardial infarction, but it is not always reliable. The EKG is a valuable tool in the diagnosis of myocardial infarction, but it is not always reliable.

[illegible]

DEVELOPERS
The following are the
names of the developers
of the program, listed
in alphabetical order.
Only those who have
been published in the
literature are listed.

[illegible][illegible][illegible]

Mass Atrial Fibrillation (AF) is a common complication of Coronary Artery Disease (CAD). AF is characterized by a rapid and irregular heart rate, which can lead to a variety of symptoms, including palpitations, dizziness, and shortness of breath. AF is a major risk factor for stroke and heart failure. The purpose of this study was to evaluate the effectiveness of a new treatment for AF, called Catheter Ablation (CA), in patients with CAD. The study included 100 patients who had been diagnosed with AF and CAD. The patients were divided into two groups: a treatment group and a control group. The treatment group received CA, and the control group received a placebo. The primary endpoint of the study was the percentage of patients who were free of AF at the end of the study. The secondary endpoints were the percentage of patients who were free of symptoms, the percentage of patients who were free of side effects, and the percentage of patients who were free of death. The results of the study showed that the treatment group had a significantly higher percentage of patients who were free of AF at the end of the study compared to the control group. The treatment group also had a significantly higher percentage of patients who were free of symptoms, side effects, and death compared to the control group. These findings suggest that CA is an effective treatment for AF in patients with CAD.

FDA rejects memantine for mild Alzheimer's

Increase size of text: A A A

July 26, 2005

The U.S. Food and Drug Administration (FDA) declined to approve memantine (Namenda) to treat mild Alzheimer's disease, Forest Laboratories announced today.

Memantine was approved to treat moderate to severe Alzheimer stages in 2003.

Forest has conducted three studies of memantine as a treatment for mild to moderate Alzheimer's. In one study, participants taking memantine fared better than those receiving a placebo on tests of memory and thinking skills as well as on assessments by their physicians and caregivers. In two other studies, memantine failed to show any statistically significant benefit compared with the placebo. In one of the studies that failed to show benefit, participants were already on a stable dose of a cholinesterase inhibitor at the time they began taking memantine.

For more information, please see:

- Alzheimer's Association [fact sheet](#) on memantine
- Forest Laboratories [press release](#) about FDA's rejection of memantine for mild Alzheimer's disease
- A Medscape [news article](#) on the study in mild Alzheimer's disease that found a statistically significant benefit for memantine
(free registration required to access article)

[Back to top](#)

[Donate](#) | [About this Site](#) | [Security and Privacy Policy](#) | [Disclaimer](#)
Copyright, reprints and permission to use material from this Web site

© 2005 Alzheimer's Association. All rights reserved.

Alzheimer's Association National Office
225 N. Michigan Ave., P.O. 17, Chicago, IL 60601
24/7 Nationwide Contact Center: 1.800.272.3900

alzheimer's 



Seal of Ap

associated with longer-term donepezil use (at least 9 to 12 months of treatment).^{19,20} In a study of patients with early-stage AD, donepezil-treated patients (mean Mini-Mental State Examination [MMSE], 24.1 ± 1.7) showed improvement in cognitive status compared with placebo-treated patients (mean MMSE, 24.3 ± 1.3).²¹ Additional data from clinical studies with physostigmine²² and donepezil²³ have shown enhancement of memory in nondemented individuals, suggesting that benefits may be possible even in individuals with the mildest symptoms of AD or MCI.

Based on the demonstrated treatment benefits of donepezil in mild to moderately severe AD, the present study was conducted to further evaluate the efficacy and safety of donepezil on cognitive impairment in patients with an amnesic form of MCI. One of the goals of this study was to determine appropriate study end points to evaluate efficacy in this patient population. Because there is no established methodology to study MCI, the approach established in AD research was adopted. Both global and cognitive measures were selected based on their potential sensitivity to drug treatment effects in patients with MCI. In addition, a battery of neuropsychologic measures was used to evaluate subjects with MCI on tests of attention and psychomotor speed.

Methods. Patient population. Men and women with MCI between 55 and 90 years of age, generally healthy and living independently, and capable of complying with testing procedures were enrolled. All patients had a documented memory complaint precluding a change from previous functioning, with MMSE²⁴ scores ≥ 24 . In addition, the Global Clinical Dementia Rating (CDR)²⁵ score was required to be 0.5 with memory box scores of 0.5 or 1, no more than one box score other than memory rated as high as 1, and no box score rated greater than 1. ADLs were required to be normal or only slightly impaired, as defined by a summed score ≥ 15 on the three functional CDR domains (house and hobbies, community affairs, and personal care). Patients could not have depressive symptoms and had a 17-item Hamilton Rating Scale for Depression (HAM-D) score ≤ 12 .²⁶ Scores on Paragraph A of the Logical Memory II Delayed Paragraph Recall subtest of the Wechsler Memory Scale-Revised (WMS-R)²⁷ were required to be below predetermined educational level cut-off scores: ≥ 8 for ≥ 16 years of education, ≥ 4 for 8 to 15 years of education, and ≥ 2 for 0 to 7 years of education. A modified Hachinski Ischemia Scale²⁸ score was required to be ≤ 4 .

Participants with a diagnosis of probable or possible AD were excluded. Drug or alcohol abuse or dependence within the previous 5 years was not permitted, as defined by criteria of the *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition Text Revision* (DSM-IV-TR).²⁹ Enrollment was not permitted for patients with poorly controlled diabetes or other medical conditions judged to be incompatible with study participation. Previous treatment with donepezil or other ChEIs (e.g., rivastigmine, tacrine, metrifonate, galantamine, physostigmine) was not permitted.

The study was conducted over a 22-month period from March 1999 to December 2000 and was carried out in compliance with the Declaration of Helsinki and its amendments. Written informed consent was obtained from patients and their legal representatives and informants.

Study design. This 24-week, multicenter, randomized, double-blind, placebo-controlled, parallel-group study was conducted at 23 study centers in the United States. Patients were randomized in a 1:1 ratio to receive either donepezil or placebo, administered orally each evening just prior to bedtime. Patients received donepezil 5 mg daily for 43 days, followed by 10 mg daily for the remainder of the study. Following the forced dose escalation,

patients unable to tolerate 10 mg donepezil were discontinued from the study. Efficacy and safety were assessed at screening, baseline, and at 6-week intervals thereafter through week 24 (final study visit), or at an early termination visit. During the study, flexibility of dosing in the morning or at bedtime was allowed.

Outcome measures. Primary efficacy measures. The primary efficacy measure for cognitive function was the change from baseline to end point on the New York University (NYU) Paragraph Test Delayed Recall.³ Different versions of the NYU Paragraph Test Delayed Recall test were used in each test period when retesting the subjects. The different versions were used in a standard order in both placebo and donepezil groups to eliminate memory and order effects.

The primary efficacy measure for global function was the change from baseline to end point on the AD Cooperative Study's Clinical Global Impression of Change (ADCS-CGIC) instrument.³⁰ The CGIC-MCI, adapted from the ADCS-CGIC, is a clinical global impression of change scale for use with patients with MCI based on semistructured interviews from both the subject and an informant. The CGIC rating is made on a 7-point Likert-type scale, where the clinician's impression of change from baseline is rated as marked improvement (1) to no change (4) to marked worsening (7).

Secondary efficacy measures. Secondary measures of cognitive function included a modified 13-item AD Assessment Scale cognitive subscale (ADAS-cog),³¹ the WMS-R Digit Span Backwards test,²⁷ the Symbol Digit Modalities test,³² and the NYU Paragraph Test Immediate Recall.³ Other secondary measures of the neuropsychologic test battery included the modified Boston Naming Task,³³ the Verbal Fluency test,³⁴ the Maze test,³⁵ and the Number Cancellation test.³⁶ Modifications to the original ADAS-cog test included the use of 12 words in the Word Recall test rather than 10, the addition of Delayed Word Recall, the reduction in number of trials of the Word Recognition Task from 3 to 1, and the addition of a Concentration/Distractibility item.³⁷

In the secondary measure of global function, patients were asked to rate their own perception of change in memory since onset of treatment using the Patient Global Assessment (PGA). The PGA is a 7-point Likert-type scale in which scores range from very much improved (1) to no change (4) to very much worse (7).

Safety measures. The safety and tolerability of treatments were assessed by monitoring adverse events, vital signs, physical examination findings, EKG abnormalities, and evaluation of clinical laboratory measures.

Data analysis. A sample size of 110 patients per treatment group was estimated based on detection of differences between placebo and drug treatment on primary efficacy variables (0.4 on the CGIC-MCI and 1.0 on the NYU Paragraph Delayed Recall) with 80% power at a 0.05 significance level. Sample size was increased by 20% to 130 per group as a result of a greater than expected premature discontinuation rate.

Analyses of efficacy were performed on the intent-to-treat (ITT) population using the last observation carried forward (LOCF) method for missing data points. The ITT population included all patients randomized to treatment who received at least one dose of study medication, had a baseline evaluation, and at least one evaluation for the primary efficacy measure. The ITT-LOCF population included all subjects who received at least one dose of medication and had at least one pretreatment evaluation and one evaluation after the start of treatment. Efficacy of donepezil treatment in the ITT-LOCF population was considered to be the primary outcome measure.

Efficacy of donepezil treatment was also analyzed in the ITT-Observed Cases (OC) and the fully evaluable (FE) population. The ITT-OC population included all subjects who received at least one dose of medication, had a baseline evaluation, and had a double-blind treatment evaluation at week 24 irrespective of compliance and protocol violations. The FE population consisted of all patients who completed 24 weeks of treatment, with $\geq 80\%$ compliance at the week 24 visit and at two other clinic visits, and no significant protocol violations. Because this study also aimed to establish methodology to study MCI, analyses of treatment effect in the OC and FE population were conducted to further evaluate the sensitivity of the selected cognitive and functional measures to donepezil treatment in patients with MCI. In addition, evaluation of the FE population was used to assess treatment effect in the patient population that best complied with the treatment protocol.




Figure 1 Patient enrollment and completion through the study.

Analyses of safety were performed using the safety population, which included all subjects who were randomized and took at least one dose of study medication.

Continuous variables were assessed using analysis of variance (ANOVA) models. Statistical analyses of efficacy measures were performed on the change from baseline scores at weeks 6, 12, 18, and 24 (OC) and at end point (LOCF). The overall treatment effect was assessed by the analysis using type III sums of squares performed at the 5% level to determine significance. Ordinal variables were analyzed using the Cochran-Mantel-Haenszel χ^2 test with MODRIDTS scores. Treatment differences in the incidence rates for adverse events were tested using Fisher's exact test. Statistical tests were two-tailed, and a value of 0.05 or below was accepted as an indicator of significance.

Results. Demographic characteristics. A total of 370 MCI subjects were randomized, of which 133 received donepezil and 137 received placebo (figure 1). Subjects in both the donepezil and placebo groups received study medication for a median of 168 days. Ninety patients from the donepezil group (67.7%) and 114 patients from the placebo group (83.2%) completed the study. Discontinuations were similar in the two study groups between baseline and week 6, but discontinuations increased in the donepezil group following dose escalation after week 6.

Subject characteristics at baseline are summarized in table 1. Subjects randomized to the donepezil and placebo groups were similar with respect to age, sex, race, education, and baseline neuropsychologic and cognitive test scores (e.g., CDR, MMSE, WMS-R, modified ADAS-cog, and NYU Paragraph Delayed Recall). The scores on the HAM-D were low (≤ 2) in both treatment groups. Donepezil-treated patients had slightly more abnormal physical examination findings at screening, but were similar with respect to other common medical comorbidities. Use of concomitant medications was reported in nearly all patients (98.5% of donepezil group and 96.4% of placebo group).

Primary efficacy measures. On the NYU Delayed Paragraph Recall test, no significant treatment effects were observed in the ITT-LOCF (table 2) and ITT-OC analyses. In measures of global function, the least squares mean OGIC-MCI scores were better for donepezil-treated than placebo-treated subjects at end point, but the differences were not significant in the ITT or FE populations. In the ITT OC population, 82.6% of donepezil-treated subjects showed minimal or moderate improvement, compared with

Table 1 Summary of baseline demographic characteristics and neuropsychologic test scores for the safety population

Characteristic	Donepezil, n = 133	Placebo, n = 137
Mean age, y \pm SD (range)	72.1 \pm 8.0 (65–89)	73.7 \pm 8.0 (66–89)
Female patients, n (%)	56 (42.1)	59 (42.8)
Race, white, n (%)	127 (95.2)	127 (92.7)
Education, y, mean \pm SD	14.9 \pm 2.8	15.0 \pm 2.8
Test scores, mean \pm SD		
MMSE	27.5 \pm 2.2	27.4 \pm 2.0
HAM-D	2.0 \pm 2.4	2.0 \pm 2.4
NYU Paragraph Delayed Recall	3.9 \pm 3.0	4.1 \pm 3.0
ADAS-cog total (89 max)	20.0 \pm 6.2	19.5 \pm 6.9
WMS-R Digit Span Backwards	6.7 \pm 2.2	7.0 \pm 2.1
Symbol Digit Modalities (no. correct)	33.8 \pm 11.0	35.1 \pm 11.6

MMSE = Mini-Mental State Examination; HAM-D = Hamilton Rating Scale for Depression; NYU = New York University; ADAS-cog = Alzheimer Disease Assessment Scale—cognitive subscale; WMS-R = Wechsler Memory Scale—Revised.

24.3% of placebo-treated subjects. 51.7% of donepezil-treated and 60.4% of placebo-treated subjects showed no change. These proportions were similar in the FE population.

Secondary efficacy measures. Donepezil treatment was associated with greater improvement on the mean change from baseline on the modified ADAS-cog total score compared with placebo in all populations (ITT-LOCF, ITT-OC, and FE) (figure 2A, see table 2). In the end point analysis, improvements on the ADAS-cog of 7 points or greater were seen in 22.3% of donepezil-treated subjects and in 12.1% of placebo-treated subjects. Improvements of 4 points or greater compared with baseline were seen in 50.0% of donepezil-treated subjects and in 31.8% of placebo-treated subjects (figure 2B). A significant improvement was seen in the donepezil-treated group in the NYU Paragraph Delayed Recall test at week 24 in the FE population. Greater improvements were also observed on the WMS-R Digit Span Backwards test and Symbol Digit Modalities test for subjects receiving donepezil compared with placebo in the FE group (see table 2 and figures E-1A and E-1B, available on the Neurology Web site; go to www.neurology.org). No significant differences were observed on the Boston Naming task, Verbal Fluency test, Mass test, and Number Cancellation test in any of the study populations. There was no difference in the patients' self-rating of their memory on the PGA in the ITT-LOCF population. The mean PGA scores at week 24 significantly favored donepezil in the OC and FE groups.

Safety. The discontinuation rate due to adverse events was 8.0% in the placebo group and 22.0% in the donepezil

Table 3 Changes in primary and secondary outcomes after 24 weeks

Psychometric test: change from baseline \pm SE	Donepezil			Placebo		
	ITT-LOCF, n = 180	ITT-OC, n = 89	FE, n = 83	ITT-LOCF, n = 182	ITT-OC, n = 110	FE, n = 100
Primary efficacy measure						
NYU Paragraph Test Delayed Recall	0.8 \pm 0.3	1.1 \pm 0.4	0.9 \pm 0.4*	0.6 \pm 0.2	0.8 \pm 0.2	0.6 \pm 0.8
Secondary efficacy measures						
ADAS-cog/18 item (total)	-3.1 \pm 0.5†	-2.8 \pm 0.5*	-3.2 \pm 0.5†	-1.2 \pm 0.5	1.0 \pm 0.4	-1.6 \pm 0.6
NYU Paragraph Test Immediate Recall	0.8 \pm 0.3	1.2 \pm 0.3*	1.1 \pm 0.3*	0.3 \pm 0.3	0.2 \pm 0.2	0.3 \pm 0.3
WMS-R Digit Span Backwards	0.6 \pm 0.2	0.6 \pm 0.2	0.6 \pm 0.2*	0.1 \pm 0.2	0.0 \pm 0.2	-0.1 \pm 0.2
Symbol Digit Modalities (no. correct)	4.4 \pm 0.7	4.5 \pm 0.9	4.9 \pm 0.9*	3.7 \pm 0.6	2.4 \pm 0.6	2.2 \pm 0.7

* $p \leq 0.05$ and † $p \leq 0.01$ vs placebo.

ITT = intent-to-treat population; LOCF = last observation carried forward; OC = observed cases analysis; FE = fully evaluable population; NYU = New York University; ADAS-cog = Alzheimer Disease Assessment Scale 18-item cognitive subscale; WMS-R = Wechsler Memory Scale-Revised.

group, with most discontinuations occurring after the forced dose escalation from 5 mg to 10 mg. Overall, adverse events occurred in 57.9% of patients in the donepezil group and 73.0% of patients in the placebo group. The most common adverse events, reported by 5% of donepezil-treated subjects and at a minimum of twice the rate of placebo, are listed in table 3. Study medication-related adverse events significantly more common in donepezil-treated subjects than placebo-treated subjects included diarrhea, abnormal dreams, nausea, leg cramps, and vomiting. Most adverse events were transient and only mild or moderate in severity. Five patients in the donepezil group and six patients in the placebo group experienced serious adverse events; only one serious adverse event (atrial fibrillation) was considered possibly related to study medication. No notable changes were reported from baseline to end point in laboratory examination, EKG, physical examination, or vital signs findings.

Discussion. The identified patient population for this study was characterized as having an amnesic form of MCI.^{1,6} The patients had memory deficits, which distinguish them from normal aging adults, and did not display significant depressive symptoms, as indicated by low HAM-D scores. Functional impairments characteristic of early AD^{1,6} were nearly absent, as indicated by the low scores on the CDR ADL domains.

On the primary outcome measures of cognition and clinician-rated global change, significant improvements were not observed in the donepezil group compared with the placebo group in the ITT populations. On the primary cognitive outcome, the NYU Paragraph Delayed Recall test, there was no significant drug-placebo difference in the ITT population. The NYU Paragraph Delayed Recall test is useful in screening patients who are at increased risk for developing dementia or AD.⁶ This is a difficult test,




Figure 2. (A) Mean (\pm SE) change from baseline in modified AD Assessment Scale-cognitive subscale (ADAS-cog) total test scores for the donepezil- and placebo-treated patients in the intent-to-treat last observation carried forward (ITT-LOCF) and fully evaluable (FE) populations. Filled bars = donepezil; open bars = placebo. (B) Cumulative percentage of patients with specified changes from baseline in the mean change from baseline in ADAS-cog score in donepezil- and placebo-treated patients. ■ = donepezil; ▲ = placebo.

Table 2 Adverse events occurring in $\geq 5\%$ of patients receiving donepezil and at a minimum, of twice the rate of placebo, n (%)

Adverse events	Donepezil, n = 132	Placebo, n = 137
Any	116 (87.9)*	100 (73.0)
Diarrhea	36 (27)*	10 (7)
Nausea	20 (15)*	9 (7)
Vomiting	12 (9)*	0 (0)
Leg cramps	12 (9)*	2 (1)
Abnormal dreams	30 (23)*	5 (4)
Dyspepsia	6 (6)	1 (1)
Insomnia	14 (11)	7 (5)

* $p \leq 0.03$ vs placebo.

even for normal individuals, with points given only for verbatim recall of phrases. Because therapeutic interventions for MCI should be directed at those individuals who are at the highest risk for developing AD, this discriminating measure appears useful. However, possible floor effects due to the relative difficulty of the test appear to limit usefulness for detecting treatment effects within the MCI population.

The CGIC-MCI was chosen as the primary efficacy measure of global function based on its validity and reliability in this population.^{29,34} The ADCS CGIC-MCI scores were better for donepezil-treated than placebo-treated patients but the differences were not significant. Because the defining feature of amnesic MCI is an isolated memory complaint, the absence of a significant effect on the CGIC-MCI is most likely due to minimal global impairment at baseline and limited decline in global function during a 6-month interval. However, this tool may have utility in longer-term studies because additional clinical symptoms (i.e., function, behavior, mood disturbances) may emerge in MCI subjects over longer observation periods. Although clinicians were not able to detect a significantly greater global response to donepezil treatment, the subjects themselves were able to detect treatment effects. In analyses of subjects completing 24 weeks of treatment (both the OC and FE populations), those treated with donepezil subjectively reported greater improvement in memory function than those given placebo. The majority of donepezil treated subjects reported feeling sharper mentally, more organized, and more confident of their memory.

Evidence of the cognitive benefits of donepezil was observed in patient performance on several secondary measures of cognitive function. Significant treatment effects were observed in the NYU Paragraph Delayed Recall test in the FE population. Significant improvements were observed in both the ITT and FE populations on the modified version of the ADAS-cog in the donepezil group. In addition, significant improvements favoring donepezil were observed in the

FE population on the Digit Span Backwards and Symbol Digit Modalities tests. However, the results of secondary analyses should be interpreted with caution given the number of secondary measures employed. In addition, higher dropout rates in the donepezil group may have influenced outcomes in the FE population. Donepezil was safe and reasonably well tolerated in this study population.

Although the treatment effects on the primary end points were not significant, the results in secondary measures suggest directions for further study in this patient population. With no established methodology available, one aim of this study was to establish cognitive and functional measures sensitive to effects of donepezil treatment in patients with MCI. In selecting the primary and secondary outcome measures, the ADAS-cog was chosen as a secondary measure because it was believed to lack sensitivity to detect treatment differences in MCI subjects. However, the study results suggest that the modified ADAS-cog total score is sufficiently sensitive to be useful in studies of MCI patients. A number of reports have found that, relative to normal elderly, patients with MCI have measurable cognitive deficits that extend beyond the memory domain. For example, MCI patients have measurable deficits in language,^{1,35,36} orientation,⁷ and psychomotor/motor function.³⁶⁻³⁸ Since test batteries such as the ADAS-cog tap into multiple domains of neuropsychologic functioning, it should not be surprising that this instrument may be particularly sensitive to a diffuse pattern of cognitive response likely to occur in treatment studies of patients with MCI. As such, consideration of its use as a primary efficacy variable in future studies of MCI appears warranted.

Several additional secondary efficacy measures were employed to evaluate other aspects of cognitive function thought to be impaired in MCI. Donepezil-related benefits in the Digit Span Backwards and Symbol Digit Modalities tests likely reflect improvement in attention, concentration, and psychomotor speed. It should be noted that subjects in both groups demonstrated some improvement on many cognitive tests, suggesting a preserved capacity to learn information, which, to our knowledge, has not been previously reported. Subjects also had a high level of education (mean 14.9 years) and practice effects may have influenced performance. It should be noted that the subjects in this study had predominantly amnesic MCI with a high level of education and a low level of depression, and lacked a medical history or current comorbidities that might account for their cognitive decline. It remains to be seen whether the performance and treatment response of this group is similar to a more heterogeneous MCI cohort.

A higher proportion of patients in the donepezil group reported adverse events compared with the placebo group. Overall, the donepezil-placebo difference in the adverse event discontinuation rate was somewhat higher than in trials of similar design in

patients with AD. Although most adverse events were mild or moderate in severity, increased study withdrawals due to adverse events closely followed the forced dose escalation. As this study did not allow reduction in dose, all subjects unable to tolerate the 10 mg dose were dropped from the study. Perhaps patients earlier in the disease course have higher baseline levels of acetylcholine in the CNS than patients with mild to moderate AD and they may have increased awareness and lower tolerability of adverse events. Consideration should be given to allow more flexible dosing and to relaxing restrictions on dose adjustments to improve tolerability in future MCI trials.

The findings from this study have implications for the design of future efficacy trials for MCI as they provide important insights into outcome measures, sample and effect sizes, dosing, and study duration. In designing this study, investigators expected that there would be a decline in cognitive function of the placebo group over the 24-week observation period (as seen in patients with mild to moderate AD^{13,14} and predicted if MCI is prodromal AD). The lack of decline in placebo-treated subjects over 6 months indicates that loss of cognitive function occurs more slowly in MCI subjects. Thus, efficacy trials of ChEIs may require trials of longer duration to adequately test for symptomatic treatment effects. Finally, the finding that patients with MCI show improvement on some cognitive tests may have implications for developing cognitive rehabilitation strategies for patients with MCI.

Acknowledgment

The authors thank Yvonne Noble, RN, clinical study manager, Eisai Inc.

Appendix

Donepezil 401 MCI Study Group Investigators: Mahesh Cheluvanaiah, DO, Innovative Clinical Research Center, Alexandria, VA; Murali Doraiswamy, MD, PhD, Duke University School of Medicine, Durham, NC; Shelley Dorman, MD, Pivotal Research, Peoria, AZ; David Drachman, MD, University of Massachusetts Medical School, Worcester; James Ferguson, MD, Pharmacology Research Clinic, Salt Lake City, UT; Steven Ferris, PhD, NY University School of Medicine, New York; Barry Gordon, MD, PhD, Johns Hopkins University School of Medicine, Baltimore, MD; Lindy Havell, MD, PhD, University of Alabama at Birmingham; Richard Holub, Neurological Association of Albany, NY; Louis Kirby II, MD, Pivotal Research, Peoria, AZ; John Kheuevich, MD, Neuroscience Options, Inc., Amherst, NH; Gary Leiser, University of Texas, Houston; William McEntee, ICSI-Clinical Studies, Sarasota, FL; Margarita Nunez, MD, ICS Clinical, St. Petersburg, FL; Eric Pfeiffer, USF Simmons Gerontology Center, Tampa, FL; Murray Rosenthal, MD, Behavioral & Medical Research LLC, San Diego, CA; Stephan Salloway, MD, Brown University, Providence, RI; Sarah C. Smith, MD, University of Miami Memory Disorder Center, Ft. Lauderdale, FL; Innovative Medical Research, Inc., Baltimore, MD; Harvey Tilker, MD, Fort Rivers Clinical Research Inc., Paducah, KY; Jeanette Wendt, Neurological Associates of Tucson, AZ; Parvaneh Zolnouni, MD, California Clinical Trials Medical, Beverly Hills.

References

- Morris JC, Storandt M, Miller JP, et al. Mild cognitive impairment represents early-stage Alzheimer's disease. *Arch Neurol* 2001;58:397-405.
- Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Knopman DS. Mild cognitive impairment: clinical characterization and outcome. [published erratum 1999;56:750]. *Arch Neurol* 1999;56:303-308.

- Almkvist O, Swann H, Backman L, et al. Mild cognitive impairment in early stage of Alzheimer's disease? *J Neural Transm Suppl* 1999;54:21-29.
- Shah Y, Tangalos EG, Petersen RC. Mild cognitive impairment. When is it a precursor to Alzheimer's disease? *Geriatrics* 2000;55:62-65-66.
- Collie A, Masliah T. The neuropsychology of preclinical Alzheimer's disease and mild cognitive impairment. *Neurosci Biobehav Rev* 2000;24:389-414.
- Petersen RC, Doody R, Kurz A, et al. Current concepts in mild cognitive impairment. *Arch Neurol* 2001;58:1907-1992.
- Wall H, Grunwald M, Ecks C, et al. The prognosis of mild cognitive impairment in the elderly. *J Neural Transm Suppl* 1998;54:1-61.
- Rubin B, Morris J, Grant L, Vendegua T. Very mild senile dementia of the Alzheimer type. I. Clinical assessment. *Arch Neurol* 1989;46:378-382.
- Kluger A, Morris SH, Golomb J, Mittelman MS, Reisberg B. Neuropsychological prediction of decline to dementia in nondemented elderly. *J Psychiatr Psychiatry* 1999;12:168-170.
- de Leon MJ, Golomb J, George AB, et al. The radiologic prediction of Alzheimer disease: the atrophic hippocampal formation. *ADNR Am J Neuroradiol* 1999;18:897-908.
- Gauthier S. Update on diagnostic methods, natural history and outcome variables in Alzheimer's disease. *Dement Geriatr Cogn Disord* 1999;9(suppl 3):2-7.
- Knapp MJ, Knopman DS, Salmon PR, Pendlebury WW, Davis CS, Gronin SI, for the Tacrine Study Group. A 24-week randomized controlled trial of high-dose tacrine in patients with Alzheimer's disease. *JAMA* 1994;271:986-991.
- Burns A, Rosner M, Barker J, et al. The effects of donepezil in Alzheimer's disease—results from a multinational trial. *Dement Geriatr Cogn Disord* 1999;10:337-344.
- Rogers SL, Farlow MR, Doody RE, Mohr R, Friedhoff LT, for the Donepezil Study Group. A 24-week, double-blind, placebo-controlled trial of donepezil in patients with Alzheimer's disease. *Neurology* 1998;50:136-145.
- Corey-Bloom J, Ahmad R, Veach J. A randomized trial evaluating the efficacy and safety of ENA 710 (rivastigmine tartrate), a new acetylcholinesterase inhibitor, in patients with mild to moderately severe Alzheimer's disease. *Int J Geriatr Psychopharmacol* 1998;1:55-65.
- Tucker PM, Salmon PR, Morris JC, Kershaw P, Lillienfeld S, Ding C, for the Galantamine USA-10 Study Group. A 6-month, randomized, placebo-controlled trial of galantamine in AD. *Neurology* 2000;54:2369-2376.
- Mohr RC, Doody RE, Morris JC, et al. A 1-year, placebo-controlled preservation of function survival study of donepezil in AD patients [published erratum 2001;57:1947]. *Neurology* 2001;57:481-488.
- Winblad B, Englund K, Sjöström H, et al. A 1-year, randomized, placebo-controlled study of donepezil in patients with mild to moderate AD. *Neurology* 2001;57:449-495.
- Geldmacher DS, Provenzano G, McKee T, Mastey V, Ieni JR. Donepezil is associated with delayed nursing home placement in patients with Alzheimer's disease. *J Am Geriatr Soc* 2003;51:987-994.
- Lopez OL, Becker JT, Wisniewski S, Saxton J, Knopman DS, DeKosky ST. Cholinesterase inhibitor treatment alters the natural history of Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 2002;72:310-314.
- Bellack D, Zolnouni P, Nunez M, et al. Donepezil treatment benefits in early-stage Alzheimer's disease. *J Am Geriatr Soc* 2000;48:3100-3101. Abstract 1170.
- Duvyk RJ, Mohr RC, Thimbleberg JR, Pfeifferbaum A, Hollister LE, Kopell BS. Physostigmine: improvement of long-term memory processes in normal humans. *Science* 1978;201:279-284.
- Yasavri JA, Mumenthaler MS, Taylor JL, et al. Donepezil and flight simulator performance: effects on retention of complex skills. *Neurology* 2002;59:133-135.
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the clinician. *J Psychiatr Res* 1975;12:129-138.
- Morris JC. The clinical dementia rating (CDR): current version and scoring rules. *Neurology* 1993;43:2412-2414.
- Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry* 1960;23:56-62.
- Lezak M. *Neuropsychological assessment*. 3rd ed. New York, NY: Oxford University Press, 1995.
- Rapoport WG, Terry RD, Paul PA, Katzman R, Paul A. Pathological verification of dementia score in differentiation of dementia. *Ann Neurol* 1980;7:486-488.
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. Fourth edition, Text revision. Washington, DC: American Psychiatric Association, 2000.
- Schneider LS, Olin JT, Doody RE, et al. Validity and reliability of the Alzheimer's Disease Cooperative Study Clinical Global Impression of Change. The Alzheimer's Disease Cooperative Study. *Alzheimer Dis Assoc Disord* 1997;11(suppl 2):S22-S32.

81. Mills RC, Kraybill D, Freeman RC, et al. Development of cognitive instruments for use in clinical trials of antideementia drugs: additions to the Alzheimer's Disease Assessment Scale that broaden its scope. The Alzheimer's Disease Cooperative Study. *Alzheimer Dis Assoc Disord* 1997;11(suppl 2):S13-S21.
82. Smith A. Symbol digit modalities test. Los Angeles, CA: Western Psychological Service, 1997.
83. Morris JC, Heyman A, Mohs RC, et al. The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part I. Clinical and neuropsychological assessment in Alzheimer's disease. *Neurology* 1989;39:1169-1175.
84. Olin JT, Schneider LS, Dendy RE, et al. Clinical evaluation of global change in Alzheimer's disease: identifying consensus. *J Geriatr Psychiatry Neurol* 1996;9:176-180.
85. Kluger A, Glenshaw J, Colomb J, et al. Patterns of motor impairment in normal aging, mild cognitive decline, and early Alzheimer's disease. *J Gerontol B Psychol Sci Soc Sci* 1997;52P28-P39.
86. Fleiter C, Perrio SM, Reisberg B. Mild cognitive impairment in the elderly: predictors of dementia. *Neurology* 1991;41:1006-1009.
87. Swarand M, Hill R. Very mild senile dementia of the Alzheimer type. II. Psychometric test performance. *Arch Neurol* 1989;46:863-868.
88. Storandt M, Dotwimick J, Danziger W, Berg L, Hughes C. Psychometric differentiation of mild senile dementia of the Alzheimer type. *Arch Neurol* 1984;41:497-499.



WWW.NEUROLOGY.ORG OFFERS IMPORTANT INFORMATION TO PATIENTS AND THEIR FAMILIES

The *Neurology* Patient Page provides:

- a critical review of ground breaking discoveries in neurologic research that are written especially for patients and their families
- up to date patient information about many neurologic diseases
- links to additional information resources for neurologic patients.

All *Neurology* Patient Page articles can be easily downloaded and printed, and may be reproduced to distribute for educational purposes. Click on the Patient Page icon on the home page (www.neurology.org) for a complete index of Patient Pages.

Memantine Treatment in Patients With Moderate to Severe Alzheimer Disease Already Receiving Donepezil

A Randomized Controlled Trial

Pierre N. Tariot, MD

Martin R. Farlow, MD

George T. Grossberg, MD

Stephen M. Graham, PhD

Scott McDonald, PhD

Ivan Gergel, MD

for the Memantine Study Group

ALZHEIMER DISEASE (AD) IS A neurodegenerative disorder characterized by cognitive decline, impaired performance of activities of daily living, and behavioral and psychiatric signs and symptoms. Pathological features of AD include intraneuronal neurofibrillary tangles containing abnormally phosphorylated tau protein, extracellular amyloid plaques containing the peptide β amyloid, neuronal cell death, and anatomic as well as functional impairment of neurotransmitter systems.^{1,2} Alzheimer disease affects approximately 4.5 million people in the United States.³ Treatments approved by the Food and Drug Administration were previously limited to monotherapy with cholinesterase inhibitors in patients with mild to moderate AD.² In October 2003, the Food and Drug Administration approved memantine for the treatment of moderate to severe AD; memantine is now available in more than 40 countries worldwide.

Memantine, a low- to moderate-affinity, uncompetitive N-methyl-D-aspartate (NMDA) receptor antago-

Context Memantine is a low- to moderate-affinity, uncompetitive N-methyl-D-aspartate receptor antagonist. Controlled trials have demonstrated the safety and efficacy of memantine monotherapy for patients with moderate to severe Alzheimer disease (AD) but no controlled trials of memantine in patients receiving a cholinesterase inhibitor have been performed.

Objective To compare the efficacy and safety of memantine vs placebo in patients with moderate to severe AD already receiving stable treatment with donepezil.

Design, Setting, and Participants A randomized, double-blind, placebo-controlled clinical trial of 404 patients with moderate to severe AD and Mini-Mental State Examination scores of 5 to 14, who received stable doses of donepezil, conducted at 37 US sites between June 11, 2001, and June 3, 2002. A total of 322 patients (80%) completed the trial.

Interventions Participants were randomized to receive memantine (starting dose 5 mg/d, increased to 20 mg/d, n=203) or placebo (n=201) for 24 weeks.

Main Outcome Measures Change from baseline on the Severe Impairment Battery (SIB), a measure of cognition, and on a modified 19-item AD Cooperative Study—Activities of Daily Living Inventory (ADCS-ADL₁₉). Secondary outcomes included a Clinician's Interview-Based Impression of Change Plus Caregiver Input (CIBIC-Plus), the Neuropsychiatric Inventory, and the Behavioral Rating Scale for Geriatric Patients (BGR Care Dependency Subscale).

Results The change in total mean (SE) scores favored memantine vs placebo treatment for SIB (possible score range, 0-100), 0.9 (0.67) vs -2.5 (0.69), respectively ($P<.001$); ADCS-ADL₁₉ (possible score range, 0-54), -2.0 (0.50) vs -3.4 (0.51), respectively ($P=.03$); and the CIBIC-Plus (possible score range, 1-7), 4.41 (0.074) vs 4.66 (0.075), respectively ($P=.03$). All other secondary measures showed significant benefits of memantine treatment. Treatment discontinuations because of adverse events for memantine vs placebo were 15 (7.4%) vs 25 (12.4%), respectively.

Conclusions In patients with moderate to severe AD receiving stable doses of donepezil, memantine resulted in significantly better outcomes than placebo on measures of cognition, activities of daily living, global outcome, and behavior and was well tolerated. These results, together with previous studies, suggest that memantine represents a new approach for the treatment of patients with moderate to severe AD.

JAMA. 2004;291:317-324

www.jama.com

nist, represents the first member of a new class of medications showing clinical benefit and good tolerability in AD. Although other NMDA receptor modulators (eg, milacemide and D-cycloserine) have failed in development as

Author Affiliations, Financial Disclosures, and Members of the Memantine Study Group are listed at the end of this article.

Corresponding Author and Reprints: Pierre N. Tariot, MD, Departments of Psychiatry, Medicine, Neurology, and the Center for Aging and Developmental Biology, University of Rochester Medical Center, Monroe Community Hospital, 435 E Henrietta Rd, Rochester, NY 14620 (e-mail: pierre_tariot@urmc.rochester.edu).

potential AD therapeutic agents,^{4,5} memantine has exhibited efficacy and safety in a recent placebo-controlled trial in outpatients with moderate to severe AD and in an earlier study in nursing home patients with dementia.^{6,7} An open-label study suggested that the combination of memantine and various cholinesterase inhibitors was well tolerated.⁸ We hypothesized that administration of memantine to patients with moderate to severe AD receiving stable donepezil therapy would result in clinical benefit and would be safe and well tolerated.

METHODS

Participants

The trial was conducted from June 11, 2001, through June 3, 2002. Participants were recruited from 37 US sites; 404 patients who had a diagnosis of probable AD, according to the National Institute of Neurological and Communicative Disorders and Stroke-Alzheimer Disease and Related Disorders Association criteria, were enrolled. Inclusion criteria were as follows: Mini-Mental State Examination (MMSE) score of 5 to 14 at both screening and baseline; minimum age of 50 years; a recent magnetic resonance imaging or computed tomographic scan (within 12 months) consistent with a diagnosis of probable AD; ongoing cholinesterase inhibitor therapy with donepezil for more than 6 months before entrance into the trial and at a stable dose (5-10 mg/d) for at least 3 months; a knowledgeable and reliable caregiver to accompany the patient to research visits and oversee the administration of the investigational agent during the trial; residence in the community; ambulatory or ambulatory-aided (ie, walker or cane) ability; and stable medical condition. Patients were permitted to continue receiving stable doses of concomitant medications, including antidepressants, antihypertensives, anti-inflammatory drugs, atypical antipsychotics, antiparkinsonian drugs, anticoagulants, laxatives, diuretics, and sedatives/hypnotics.

Patients were excluded for clinically significant B₁₂ or folate deficiency; ac-

tive pulmonary, gastrointestinal, renal, hepatic, endocrine, or cardiovascular disease; other psychiatric or central nervous system disorders; computed tomographic or magnetic resonance imaging evidence of clinically significant central nervous system disorders other than probable AD; dementia complicated by other organic disease; or a modified Hachinski Ischemia Score⁹ of more than 4 at screening. Written informed consent was obtained from the caregiver and either the patient (if possible) or a legally acceptable representative (if different from the caregiver) before the initiation of any study-specific procedures. The study was reviewed and approved by the institutional review board at each site.

Interventions

This study was a prospective randomized, placebo-controlled, parallel-group, fixed-dose trial in which participants were assigned to double-blind treatment for 24 weeks, with a 1- to 2-week single-blind placebo lead-in period before randomization solely to assess compliance. Patients were randomly allocated to 1 of the 2 treatment groups in permuted blocks of 4 in accordance with the randomization list generated and retained by the Department of Biostatistics at Forest Laboratories. At the baseline visit, each investigator sequentially assigned a randomization number to each patient. No individual patient randomization code was revealed during the trial.

Patients assigned to double-blind memantine treatment were titrated in 5-mg weekly increments from a starting dose of 5 mg/d to 20 mg/d (administered as two 5-mg tablets twice daily) at the beginning of week 4. Masked study medication was supplied to each study site for dispensation in blister packs at each visit. Drug and placebo tablets were visually identical and all patients received 4 tablets of study medication daily (in combinations of memantine [5 mg] and matching placebo tablets). All patients were to maintain stable donepezil therapy at entry dose as prescribed by the patient's phy-

sician for the duration of the study; adherence to this protocol was monitored by routine assessment of concomitant medication use. Any change in the dosing regimen or discontinuation of donepezil was recorded, and patients were discontinued from the study if the inclusion criterion of concomitant donepezil therapy was no longer met. From week 3 to the end of week 8 of double-blind treatment, transient dosage adjustments for memantine treatment were permitted for patients experiencing dose-limiting adverse events. All patients receiving memantine were required to receive the target dose of 20 mg/d by the end of week 8. Patients not tolerating the target dose by week 8 were disenrolled. Adherence with study medication was assessed by returned tablets and more than 95% of both treatment groups had more than 75% compliance (95% for the placebo-treatment group and 96.5% for the memantine-treatment group). Most patients who completed the double-blind phase entered the currently ongoing open-label extension.

Outcome Measures

Cognitive, functional, and global outcome measures were obtained at baseline and at the end of weeks 4, 8, 12, 18, and 24, unless otherwise specified. Patients who discontinued prematurely were evaluated during the final visit. The primary efficacy parameters were the change from baseline on the Severe Impairment Battery (SIB) and on a modified 19-item AD Cooperative Study-Activities of Daily Living Inventory (ADCS-ADL₁₉) at week 24.

The SIB is a 40-item test developed for the evaluation of cognitive dysfunction in patients with more severe AD. Six primary subscales assess memory, orientation, language, attention, visuospatial ability, and construction. In addition, the scale assesses praxis, social interaction, and orienting to name.^{10,11} Validity, reliability, and sensitivity to longitudinal change have been established.^{10,11} The SIB scores range from 0 to 100, with higher scores reflecting

higher levels of cognitive ability. The SIB was assessed at baseline and all subsequent visits.

The ADCS-ADL₁₉ was the second primary efficacy instrument.¹² This 19-item subset of the original 42-item inventory focuses on items appropriate for the assessment of later stages of dementia (ie, the level of independence in performing everyday tasks including eating, walking, grooming, telephone use, hobbies, complex tasks, and communications). The sensitivity and reliability of this modification have been established.¹³ The ADCS-ADL₁₉ was administered as an interview to the patient's caregiver and focused on the performance of each activity of daily living during the previous 4 weeks. Possible scores range from 0 to 54. Higher scores reflect higher levels of functioning. The ADCS-ADL₁₉ was assessed at baseline and all subsequent visits.

The secondary outcomes included a Clinician's Interview-Based Impression of Change Plus Caregiver Input (CIBIC-Plus),¹⁴ the Neuropsychiatric Inventory (NPI), and the Behavioral Rating Scale for Geriatric Patients (BGP). The CIBIC-Plus was administered according to the format of the Alzheimer Disease Cooperative Study—Clinician's Global Impression of Change. The CIBIC-Plus is used to assess the effect of medication on overall clinical status in patients with dementia, incorporating caregiver observations as well as patient interviews. Change is rated on a scale from 1 (marked improvement) to 7 (marked worsening). A global assessment of severity of illness was made at baseline; the CIBIC-Plus was assessed at all post-baseline visits.

The NPI was designed to assess the frequency and severity of behavioral symptoms in patients with dementia, based on an interview of the caregiver.¹⁵ The 12-item version of the instrument was used with a total score ranging from 0 to 144. Higher scores reflect greater symptoms. The NPI was assessed at baseline, at the end of week 12, and at the final visit.

The BGP consists of 35 items (scored 0, 1, or 2 by the rater) assessing observable aspects of cognition, function, and behavior.¹⁶ A higher score reflects worse function. The BGP care dependency subscale reflects cognitive and functional characteristics associated with increased need for care. The BGP was administered at baseline and the final visit.

The Functional Assessment Staging (FAST) was administered as an index of staging and not as a secondary outcome.¹⁷ The FAST evaluates a patient's ability to perform daily and necessary life activities and is divided into 7 major stages, from normality (FAST stage 1) to severe dementia (FAST stage 7). Stages 6 and 7 are further divided into 11 substages (6a to 6e and 7a to 7f), each of which is based on specific functional deficits. The FAST was administered at baseline and the final visit.

Concomitant medications and vital signs were recorded at every visit; adverse events were recorded at baseline and all subsequent visits; and laboratory tests, electrocardiograms, and physical examinations were performed at the screening and final visits.

Sample Size

Assuming a hypothetical effect size of 0.35, a sample size of at least 170 patients in each treatment group provided a 90% power at a 2-sided α level of .05, based on a 2-sample *t* test for change from baseline to week 24 in both SIB and ADCS-ADL₁₉ scores.

Statistical Analyses

Three populations were considered in the statistical analyses. The randomized population consisted of all patients randomized into the study ($n=404$); the safety population consisted of all randomized patients who received at least 1 dose of double-blind study medication ($n=403$); the modified intent-to-treat population specified by the protocol consisted of patients in the safety population who completed at least 1 postbaseline SIB or ADCS-ADL₁₉ assessment ($n=395$). The statistical analysis plan for this study

stipulated that only postbaseline data could be carried forward. Particularly for the CIBIC-Plus, it is not possible to carry forward baseline data because by definition this is a change score and is not applicable to baseline. All efficacy analyses were based on the modified intent-to-treat population. Primary efficacy analyses were conducted by using the last observation carried forward (LOCF) approach for missing data imputation. Supportive analyses were performed by using the observed case approach. Change from baseline was compared between memantine and placebo groups using a 2-way analysis of covariance, with treatment group and center as main effects and baseline total score as the covariate. The study was to be declared positive if memantine was statistically significantly better than placebo ($P<.05$) on both the SIB and ADCS-ADL₁₉. For categorical measures, the Cochran-Mantel-Haenszel statistic using modified Ridit scores (Van Elteren test) controlling for study center was used to compare distributions between memantine and placebo groups. No interim analyses were performed. SAS version 6.12 (SAS Institute, Cary, NC) was used for all analyses.


RESULTS

Participants

The trial profile is summarized in FIGURE 1. Of the 404 patients who entered the study, 201 were randomized to placebo and 203 were randomized to memantine (1 in the memantine group withdrew consent before receiving treatment). No patients were excluded during the placebo lead-in period for lack of compliance. Significantly more participants in the memantine group ($n=172$, 85.1%) completed the study than in the placebo group ($n=150$, 74.6%, $P=.01$). No patients discontinued because of changes in administration of donepezil.

The demographic and clinical characteristics of the 2 groups at baseline are summarized in TABLE 1. Patients in the memantine group were slightly heavier ($F=.003$) than those in the pla-

Figure 1. Study Flow



cebo group; retrospectively adding this variable to the analysis of covariance did not affect the primary outcomes. No clinically relevant group differences were observed for the duration of donepezil use before baseline or for any other characteristic at baseline. Most patients (87%) had a FAST rating between 4 and 6c. The most frequent medical conditions were not recorded, but the following body systems were noted to be affected at screening for placebo and memantine groups, respectively: eyes-ears-nose-throat (43% and 43%), neurological (34% and 38%), appearance/skin (40% and 33%), musculoskeletal (29% and 29%), cardiovascular (20% and 23%), abdomen (12% and 17%), head/neck (6% and 9%), other (10% and 9%), and pulmonary (3% and 5%). The most frequent medication classes (>20%) used during treatment with placebo and memantine, respectively, were vitamins (74% and 77%), analgesics (48% and 48%), antidepressants (36% and 36%), mineral supplements (22% and 27%), lipid-reducing agents (23% and 25%), anxiolytics/neuroleptics (26% and 22%), and anti-inflammatory agents (21% and

24%). There were no statistically significant differences between groups in the number or type of medical disorders experienced previously or at the time of enrollment, or in the number or type of concomitant medications used during the study.

Efficacy

Statistically significant benefits of treatment with memantine vs treatment with placebo were observed on all primary and secondary outcome measures as presented. TABLE 2 summarizes primary and secondary efficacy outcomes at week 24 and at end point, using both the observed case and LOCF analytical approaches.

Primary Outcomes

Analyses using the LOCF approach showed a statistically significant benefit of memantine treatment vs treatment with placebo on the SIB ($P<.001$) and the ADCS-ADL₁₉ ($P=.03$), as did analyses using the observed case approach ($P<.001$ for SIB; $P=.02$ for ADCS-ADL₁₉). Post hoc analyses including all randomized patients also showed statistically significant benefits consistent with analyses using the modified intent-to-treat population (for SIB, $P<.001$ and for ADCS-ADL₁₉, $P=.03$).

FIGURE 2 depicts the mean change from baseline by visit and at end point on the SIB by using observed case and LOCF, showing statistically significant differences between the memantine and placebo groups at all visits beginning at week 8; the mean SIB values for the patients receiving memantine remained above baseline throughout the trial. Figure 2 also depicts the mean change in total ADCS-ADL₁₉ from baseline by visit and at end point by using observed case and LOCF, respectively, showing a statistically significant difference ($P<.05$) from placebo beginning at week 4.

Secondary Outcomes

A CIBIC-Plus score was used as a measure of overall clinical response to therapy. The mean CIBIC-Plus score was

statistically significantly better for the memantine group vs the placebo group using both observed case and LOCF (Table 2). Furthermore, 55% of the memantine group was rated as improved or unchanged vs 45% of the placebo group at end point. FIGURE 3 provides the distribution of CIBIC-Plus ratings at end point using LOCF analysis.

The total NPI score was significantly lower for the memantine group compared with the placebo group at week 24 ($P=.01$ with observed case analysis and $P=.002$ with LOCF), representing fewer behavioral disturbances and psychiatric symptoms for patients in the memantine group. The BGP care dependency subscale was also statistically significantly improved for the memantine group compared with the placebo group ($P=.001$ using observed case and $P=.001$ using LOCF; Table 2).

Safety and Tolerability

Overall treatment-emergent adverse events are summarized in TABLE 3. More participants ($n=25$, 12.4%) in the placebo-treated group discontinued prematurely because of adverse events than in the memantine group ($n=15$, 7.4%; Figure 1). The adverse event most often associated with discontinuation was confusion, resulting in discontinuation in 1.5% of patients in the placebo group and 2% in the memantine group.

Adverse events occurred in 72% of the placebo and 78% of the memantine groups. Most adverse events were rated as mild or moderate in severity and were judged to be not related to study drug for participants in both treatment groups. The only adverse events that occurred in at least 5% of the memantine group and with an incidence of at least twice that of the placebo group were confusion (7.9% vs 2.0%, respectively; $P=.01$) and headache (6.4% vs 2.5%, respectively; $P=.09$). By similar criteria, lower incidences of diarrhea (4.5% vs 8.5%) and fecal incontinence (2.0% vs 5.0%) were observed in the memantine group compared with the placebo group, respectively. Other gastrointes-

tinal effects of interest for patients receiving cholinesterase inhibitors included nausea, which was reported by 3.5% of the placebo group and 0.5% of the memantine group, and constipation, which was reported by 1.5% of the placebo group and 3.0% of the memantine group.

Of the patients who experienced confusion, 4 (25%) of 16 patients receiving memantine discontinued treatment because of this adverse event, whereas 3 (75%) of 4 patients receiving placebo did so. In most of the patients receiving memantine, confusion was rated as mild, occurred at a median of 32 days, and remitted within 2 weeks. In patients receiving placebo, confusion was more likely to be rated as severe, occurred at a median of 55 days, and did not remit. No patients discontinued because of headache, which usually lasted 1 day.

No clinically significant differences were detected between treatment groups in the mean change from baseline to end point or in the incidence of potentially clinically significant values for laboratory tests, vital sign measurements, or electrocardiogram parameters.

COMMENT

To our knowledge, this is the first published, prospective, double-blind, placebo-controlled study examining the benefits of an NMDA receptor antagonist in patients with AD receiving a

stable dose of donepezil. Efficacy of memantine was significantly better than placebo for treatment of moderate to severe AD in community-dwelling patients. Specifically, measures of cognitive function, activities of daily living,

Table 1. Baseline Demographic and Clinical Characteristics*

Characteristics	Placebo (n = 201)	Memantine (n = 202)
Men	67 (33)	74 (37)
Women	134 (67)	128 (63)
Age, mean (SD), y	75.5 (8.73)	75.5 (8.45)
Weight, mean (SD), kg	68.4 (14.12)	70.7 (14.31)†
White race	186 (92.5)	182 (90.1)
MMSE score, mean (SD)	10.2 (2.98)	9.9 (3.13)
Duration of donepezil treatment, mean (SD), wk	129 (70.3)	126 (64.9)
Donepezil dose, mean (SD), mg	9.49 (1.88)	9.25 (1.79)
Any concurrent medical condition	149 (74.1)	149 (73.8)
Any concomitant medication during treatment	197 (98.0)	197 (97.5)
Tocopherol	120 (59.7)	131 (64.9)
Multivitamins	78 (38.8)	80 (39.6)
Acetylsalicylic acid	76 (37.8)	73 (36.1)
Ascorbic acid	35 (17.4)	43 (21.3)
Paracetamol	25 (12.4)	32 (15.8)
Ginkgo biloba	24 (11.9)	31 (15.3)
Calcium	21 (10.4)	26 (12.4)

Abbreviation: MMSE, Mini-Mental State Examination.

*Data are No. (%) unless otherwise specified. One randomized patient discontinued the study prior to receiving any treatment and was not included in the analyses.

†P = .003.

Table 2. Efficacy Outcomes at Week 24 (Observed Case) and at End Point (LOCF)*

Outcome Measure	Least Squares Mean Score (SE)							
	Baseline		Change From Baseline			Week 24 Observed Case		
	Placebo	Memantine	End Point LOCF†		P Value	Placebo	Memantine	P Value
			Placebo	Memantine				
SIB	80.0 (1.13)	78.0 (1.11)	-2.5 (0.69)	0.9 (0.67)	<.001	-2.4 (0.74)	1.0 (0.70)	<.001
No. of patients	197	198	196	198		153	171	
ADCS-ADL ₁₀	35.8 (0.74)	35.5 (0.73)	-3.4 (0.51)	-2.0 (0.50)	.03	-3.3 (0.55)	-1.7 (0.51)	.02
No. of patients	197	198	197	198		152	172	
CIBIC-Plus‡	NA	NA	4.66 (0.075)	4.41 (0.074)	.03	4.64 (0.087)	4.38 (0.081)	.03
No. of patients	197	198	196	198		162	172	
NPI	13.4 (1.08)	13.4 (1.07)	3.7 (0.89)	-0.1 (0.98)	.002	2.9 (1.06)	-0.5 (0.99)	.01
No. of patients	197	198	189	193		152	171	
BGP Care Dependency Subscale	9.8 (0.46)	9.5 (0.45)	2.3 (0.38)	0.6 (0.37)	.001	2.2 (0.40)	0.6 (0.37)	.001
No. of patients	196§	198	179	185		151	172	


Abbreviations: ADCS-ADL₁₀, 10-item Alzheimer Disease Cooperative Study-Activities of Daily Living Inventory; BGP, Behavioral Rating Scale for Geriatric Patients; CIBIC-Plus, Clinician's Interview-Based Impression of Change Plus Caregiver Input; LOCF, last observation carried forward; NA, not applicable; NPI, Neuropsychiatric Inventory; SIB, Severe Impairment Battery.

*SIB range of possible scores, 0 to 100; higher score indicates better function. ADCS-ADL₁₀ range of possible scores, 0 to 54; higher score indicates better function. CIBIC-Plus was defined as a change score, therefore baseline values are not applicable; range of possible scores, 1 (marked improvement) to 7 (marked worsening). NPI range of possible scores, 0 to 144; higher scores indicate worse symptoms. BGP range of possible scores, 0 to 70; higher scores indicate worse function.


†For the end point LOCF approach, only postbaseline assessments were carried forward.

‡Arithmetic mean.

§One patient had an incomplete BGP baseline assessment and was not included.

Figure 2. SIB and ADCS-ADL₁₉ by Visit (Observed Case) and at End Point (LOCF)

SIB indicates Severe Impairment Battery; ADCS-ADL₁₉, 19-item Alzheimer Disease Cooperative Study-Activities of Daily Living Inventory; LOCF, last observation carried forward. For the Severe Impairment Battery, the mean (SD) score at baseline was 79.8 (14.18) for the placebo group and 77.8 (15.46) for the memantine group. For the Alzheimer Disease Cooperative Study-Activities of Daily Living Inventory, the mean (SD) score at baseline was 36.2 (9.32) for the placebo group and 35.9 (9.75) for the memantine group. Only patients with at least 1 postbaseline assessment were included in the LOCF analysis. The end point is the last nonmissing post-baseline assessment carried forward to end of study. Error bars indicate SEM.

Figure 3. Distribution of CIBIC-Plus Ratings at End Point (LOCF)

CIBIC-Plus indicates Clinician's Interview-Based Impression of Change Plus Caregiver Input; LOCF, last observation carried forward. $P=.03$ for the comparison between the distribution of values for the memantine and placebo groups, determined by the Cochran Mantel-Haenszel statistic using modified Ridit scores (Van Elteren test) controlling for study center.

behavior, and clinical global status were significantly improved with memantine compared with placebo. Treatment with memantine during the 6-month trial in patients with MMSE scores of 5 to 14 resulted in the maintenance of cognitive function (0.9 increase in SIB score compared with baseline), whereas treatment with pla-

cebo was associated with cognitive decline (2.5 decrease in SIB score compared with baseline). In comparison, the AD Cooperative Study group reported that for untreated patients with AD with MMSE scores of 5 to 9, the mean deterioration rate on the SIB was roughly 3.19 per month and for untreated patients with AD with MMSE scores of 10

to 15, the rate of change was 2.08 per month.¹⁰ Treatment with memantine was associated with less decline on the CIBIC-Plus.

These efficacy findings confirm and extend results from previous placebo-controlled trials of memantine in dementia. A 12-week multicenter European trial⁶ of memantine 10 mg/d was conducted in 166 nursing home residents with severe dementia, including both Alzheimer type and vascular dementias, diagnosed by *Diagnostic and Statistical Manual of Mental Disorders, Revised Third Edition* criteria¹⁸ (mean baseline MMSE of 6.3). Significant benefit of memantine vs placebo was observed on the Clinician's Global Impression of Change and the BGP care dependency subscale, and there were no clinically relevant differences in adverse events between memantine (21%) and placebo (22%) groups. A more recent 28-week multicenter US trial of memantine 20 mg/d monotherapy was conducted in 252 patients with moderate to severe probable AD by the National Institute of Neurological and Communicative Disorders and Stroke-

Alzheimer Disease and Related Disorders Association criteria and who were not permitted to receive a cholinesterase inhibitor.⁷ Significant benefit of memantine treatment was observed on the ADCS-ADL₁₀, an assessment of function, and on the SIB, an assessment of cognition using both observed case and LOCF approaches, and on the CIBIC-Plus in the observed case but not the LOCF analysis. Adverse events were similar between the memantine (84%) and placebo (87%) groups.

Patients in the memantine monotherapy outpatient study⁷ were more cognitively impaired (mean baseline MMSE of 7.9), more functionally impaired (mean baseline ADCS-ADL₁₀ score of approximately 27), and experienced more psychopathology (mean baseline NPI of approximately 20; rates of agitation as an adverse event in 32% and 18% of patients treated with placebo and memantine, respectively) than patients included in this trial. In addition, the magnitude of the memantine-placebo differences in outcomes common to both studies, as well as the magnitude of decline in most measures over time, was greater in the memantine monotherapy study than observed in this trial. This finding may be related to the higher severity of dementia in patients enrolled in the memantine monotherapy trial or because the present trial required donepezil therapy and permitted use of most psychotropics, factors which may have contributed to slower rates of decline in both the memantine and placebo groups. However, this type of inference is speculative given the absence of patients who were not treated with donepezil. Similar to the finding in the present trial, discontinuation rates because of adverse events in the monotherapy study were lower in patients receiving memantine than in those receiving placebo (10% vs 17%, respectively).⁷ These trials support the efficacy of memantine for patients with moderate to severe AD.

Memantine administered at a dosage of 20 mg/d to patients receiving stable doses of donepezil was safe and well tolerated. Significantly more pa-

tients receiving placebo discontinued the trial than patients receiving memantine and the rate of discontinuation because of adverse events was lower in the memantine-treated group than in the placebo-treated group. The incidence of individual adverse events was generally similar in the 2 groups. Confusion, although occurring at a low frequency, was more common in patients receiving memantine than in those patients receiving placebo. However, it did not lead to a greater proportion of discontinuations and was mild in intensity and duration. The gastrointestinal adverse effects associated with cholinergic compounds were more commonly reported by patients receiving placebo, which was suggestive of a possible amelioration of these adverse events by the addition of memantine treatment to patients receiving a stable regimen of donepezil therapy. There were no clinically significant memantine-related mean changes in laboratory test results, vital signs, or electrocardiogram parameters.

There are limitations to the generalizability of our results. The trial did not address different doses or titration rates, the use of other cholinesterase inhibitors besides donepezil, or the impact of commencing memantine therapy before donepezil. Although there is no a priori reason to expect different results with other cholinesterase inhibitors, studies of memantine in combination with other cholinesterase inhibitors are being conducted to address this issue. Furthermore, results from an open-label European trial indicated that tolerability was not affected when donepezil or other cholinesterase inhibitors were administered to patients already receiving memantine or vice versa.⁸ Preclinical studies show that memantine does not affect the inhibition of acetylcholinesterase by donepezil, nor does it bind to muscarinic receptors.¹⁹⁻²¹ Furthermore, in healthy volunteers, no pharmacokinetic or pharmacodynamic interactions were observed between memantine and donepezil.²² Although memantine has demonstrated positive

Table 3. Adverse Events Reported in at Least 5% of Patients in Either Treatment Group*

	Adverse Event, No. (%)	
	Placebo (n = 201)	Memantine (n = 202)
Agitation	24 (11.9)	19 (9.4)
Confusion	4 (2.0)	16 (7.9)
Fall	14 (7.0)	15 (7.4)
Influenza-like symptoms	13 (6.5)	15 (7.4)
Dizziness	18 (8.0)	14 (6.9)
Headache	5 (2.5)	13 (6.4)
Urinary tract infection	10 (5.0)	12 (5.9)
Urinary incontinence	6 (3.0)	11 (5.4)
Accidental injury	16 (8.0)	10 (5.0)
Upper respiratory tract infection	13 (6.5)	10 (5.0)
Peripheral edema	8 (4.0)	10 (5.0)
Diarrhea	17 (8.5)	9 (4.5)
Fecal incontinence	10 (5.0)	4 (2.0)

*Patients may have reported more than 1 adverse event.

cognitive effects in patients with mild to moderate vascular dementia, the efficacy of memantine administered alone or along with any cholinesterase inhibitor in other forms of dementia was not systematically evaluated in this trial.^{23,24}

The long-term effects of memantine and cholinesterase inhibitor treatment were not addressed in this double-blind trial but are the focus of the open-label extension and other ongoing trials. Considering that patients in this study had been receiving stable long-term donepezil therapy before enrollment, it is possible that participants were more likely to experience good tolerability and efficacy in the trial, perhaps because of having fewer medical problems or experiencing a slower rate of decline than patients without any prior AD treatment. However, the use of concomitant medications was typical for this elderly patient population and was similar between the groups. In addition, the dropout rate was approximately 15% in the memantine group vs approximately 25% in the placebo group, a phenomenon that perhaps led to an underestimation of the effect of memantine.

Drugs that target the glutamatergic system appear to have a therapeutic role in AD.^{25,26} Memantine may block pathological activation of NMDA receptors while dissociating from the NMDA re-

ceptor channel during normal physiological conditions,²⁰ in theory improving cognition in states of glutamatergic excess. It is plausible that combining donepezil and memantine, which affect separate neurotransmitter systems, may confer independent clinical benefits. However, given the complex interconnection of different neurotransmitter systems, a synergistic mechanism is also plausible. Although the specific mechanisms and interactions between these therapies have not yet been defined, this and other studies demonstrate that memantine alone or together with a cholinesterase inhibitor results in significantly better outcomes than placebo in patients with moderate to severe AD.

Author Affiliations: Departments of Psychiatry, Medicine, Neurology, and the Center for Aging and Developmental Biology, University of Rochester Medical Center, Monroe Community Hospital, Rochester, NY (Dr Tariot); Department of Neurology, Indiana University School of Medicine, Indianapolis (Dr Farlow); Department of Geriatric Psychiatry, St Louis University School of Medicine, St Louis, Mo (Dr Grossberg); Forest Laboratories Inc, Jersey City, NJ (Drs Graham, McDonald, and Gergel).

Financial Disclosures: Dr Tariot has received research support, consultation fees, and honoraria from Abbott, AstraZeneca, Bristol-Myers Squibb, Eisai, Eli Lilly, Forest, Janssen, Novartis, Pfizer, and Schwabe. Dr Farlow receives research support and consultation fees from Axonyx, Bristol-Myers Squibb, Eisai, Eli Lilly, EUNOE, Forest, Janssen, Novartis, Pfizer, and Schwabe. Dr Grossberg received consultation fees from AstraZeneca, Forest, Janssen, and Novartis and honoraria from AstraZeneca, Forest, Janssen, and Novartis. Drs Graham, McDonald, and Gergel are employees with stock options at Forest Laboratories.

Author Contributions: Dr Tariot had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Tariot, Gergel.

Acquisition of data: Tariot, Farlow, McDonald.

Analysis and interpretation of data: Tariot, Farlow, Grossberg, Graham, McDonald, Gergel.

Drafting of the manuscript: Tariot, Farlow, Grossberg, Graham.

Critical revision of the manuscript for important intellectual content: Tariot, Farlow, Grossberg, Graham, McDonald, Gergel.

Statistical expertise: Gergel.

Obtained funding: Gergel.

Administrative, technical, or material support: Tariot, Farlow, Graham, McDonald, Gergel.

Study supervision: Tariot, Farlow, Graham, McDonald, Gergel.

Members of the Memantine Study Group: Paul S. Aisen, MD, Washington, DC; Peter Aupperle, MD, Piscataway, NJ; Barry Baumeister, MD, Fort Lauderdale, Fla; William Burke, MD, Omaha, Neb; Jody Corey-Bloom, MD, PhD, San Diego, Calif; P. Murali Doraiswamy, MD, Durham, NC; Larry Eisner, MD, Fort Lauderdale, Fla; Martin R. Farlow, MD, Indianapolis, Ind; M. Fred Farmer, MD, St Petersburg, Fla; George T. Grossberg, MD, St

Louis, Mo; Daniel Grosz, MD, Northridge, Calif; Howard A. Hassman, DO, Clementon, NJ; Jon Heiser, MD, Newport Beach, Calif; Richard F. Holub, MD, Albany, NY; Ari Kiev, MD, New York, NY; Louis Kirby, MD, Peoria, Ariz; Steven Kobetz, MD, Miami, Fla; Anne M. Lipton, MD, PhD, Dallas, Tex; Scott N. Losk, PhD, Portland, Ore; David I. Margolin, MD, PhD, Fresno, Calif; Jeffrey A. Mattes, MD, Princeton, NJ; Craig McCarthy, MD, Mesa, Ariz; Barry Meyers, MD, White Plains, NY; Bruce L. Miller, MD, San Francisco, Calif; Jacobo Mintzer, MD, Charleston, SC; Eric Pfeiffer, MD, Tampa, Fla; Ralph Richter, MD, FACP, Tulsa, Okla; Carl H. Sadowsky, MD, FAAN, West Palm Beach, Fla; Beth Safirstein, MD, Fort Lauderdale, Fla; Mary Sano, MD, New York, NY; Benjamin Seltzer, MD, New Orleans, La; Joshua Shua-Haim, MD, Lakehurst, NJ; Pierre N. Tariot, MD, Rochester, NY; Harvey A. Tilker, PhD, Paducah, Ky; Larry Tune, MD, Atlanta, Ga; Mahmood A. Usman, MD, Pittsburgh, Pa; Chris H. van Dyck, MD, New Haven, Conn. **Funding/Support:** This study was supported by funding from Forest Research Institute, a division of Forest Laboratories Inc.

Role of the Sponsor: Forest Laboratories provided all financial and material support for the research, consulted with the authors and the members of the Memantine Study Group on the study design, monitored the conduct of the study as well as the collection of the data, analyzed and interpreted the data, and assisted the authors in the preparation, review, and approval of the manuscript.

Independent Analysis: Weichung J. Shih, PhD, from the University of Medicine and Dentistry of New Jersey provided an independent statistical analysis confirming the primary efficacy results.

Acknowledgment: We thank James Jin, PhD, from Forest Laboratories, who provided statistical support for the analysis of the data, and Grace Lee, BA, from Forest Laboratories, who provided editorial support for the manuscript.

REFERENCES

1. Selkoe DJ. The molecular pathology of Alzheimer's disease. *Neuron*. 1991;6:487-498.
2. Cummings JL, Cole G. Alzheimer disease. *JAMA*. 2002;287:2335-2338.
3. Hebert LE, Scherr PA, Bienias JL, Bennett BD, Evans DA. Alzheimer's disease in the US population. *Arch Neurol*. 2003;60:1119-1122.
4. Dysken MW, Mendels J, LeWitt P, et al. Milamceme: a placebo-controlled study in senile dementia of the Alzheimer type. *J Am Geriatr Soc*. 1992;40:503-506.
5. Laake K, Oeksengaard AR. D-cycloserine for Alzheimer's disease. *Cochrane Database Syst Rev*. 2002;(2):CD003153.
6. Winblad B, Poritis N. Memantine in severe dementia: results of the M-BEST study (benefit and efficacy in severely demented patients during treatment with memantine). *Int J Geriatr Psychiatry*. 1999;14:135-146.
7. Reisberg B, Doody R, Stöffler A, Schmitt F, Ferris S, Möbius HJ. Memantine in moderate-to-severe Alzheimer's disease. *N Engl J Med*. 2003;348:1333-1341.
8. Hartmann S, Möbius HJ. Tolerability of memantine in combination with cholinesterase inhibitors in dementia therapy. *Int Clin Psychopharmacol*. 2003;18:81-85.
9. Rosen WC, Terry RD, Fuld PA, Katzman R, Peck A. Pathological verification of ischemic score in differentiation of dementias. *Ann Neurol*. 1980;7:486-488.
10. Schmitt FA, Ashford W, Ernesto C, et al. The severe impairment battery: concurrent validity and the assessment of longitudinal change in Alzheimer's disease. *Alzheimer Dis Assoc Disord*. 1997;11(suppl 2):S51-S56.
11. Panisset M, Roudier M, Saxton J, Boller F. Severe impairment battery: a neuropsychological test for severely demented patients. *Arch Neurol*. 1994;51:41-45.
12. Galasko D, Bennett D, Sano M, et al. An inventory to assess activities of daily living for clinical trials in Alzheimer's disease. *Alzheimer Dis Assoc Disord*. 1997;11(suppl 2):S33-S39.
13. Galasko DR, Schmitt FA, Jin S, et al. Detailed assessment of cognition and activities of daily living in moderate to severe Alzheimer's disease. *Neurobiol Aging*. 2000;21(suppl 1):S168.
14. Schneider LS, Olin JT, Doody RS, et al. Validity and reliability of the Alzheimer's Disease Cooperative Study—Clinical Global Impression of Change. *Alzheimer Dis Assoc Disord*. 1997;11(suppl 2):S22-S32.
15. Cummings JL, Mega M, Gray K, Rosenberg-Thompson S, Carusi DA, Gombeln J. The Neuropsychiatric Inventory: comprehensive assessment of psychopathology in dementia. *Neurology*. 1994;44:2308-2314.
16. van der Kam P, Mol F, Wimmers M. *Beoordelingsschaal voor Oudere Patienten (BOP)*. Deventer, the Netherlands: Van Loghum Slaterus; 1971.
17. Scam SC, Reisberg B. Functional assessment staging (FAST) in Alzheimer's disease: reliability, validity, and ordinality. *Int Psychogeriatr*. 1992;4(suppl 1):55-69.
18. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Revised Third Edition*. Washington, DC: American Psychiatric Association; 1987.
19. Wenk GL, Quack G, Möbius H-J, Danysz W. No interaction of memantine with acetylcholinesterase inhibitors approved for clinical use. *Life Sci*. 2000;66:1079-1083.
20. Parsons CG, Danysz W, Quack G. Memantine is a clinically well tolerated N-methyl-D-aspartate (NMDA) receptor antagonist: a review of preclinical data. *Neuropharmacology*. 1999;38:735-767.
21. Danysz W, Parsons CG, Kornhuber J, Schmidt WJ, Quack G. Amino acid antagonists as NMDA receptor antagonists and antiparkinsonian agents—preclinical studies. *Neurosci Biobehav Rev*. 1997;21:455-468.
22. Periclou A, Ventura D, Sherman T, Rao N, Abramowitz W. A pharmacokinetic study of the NMDA receptor antagonist memantine and donepezil in healthy young subjects (abstract). *J Am Geriatr Soc*. 2003;51:S225.
23. Wilcock G, Möbius HJ, Stöffler A. A double-blind, placebo-controlled multicentre study of memantine in mild to moderate vascular dementia (MMM500). *Int Clin Psychopharmacol*. 2002;17:297-305.
24. Orgogozo JM, Rigaud AS, Stöffler A, Möbius HJ, Forette F. Efficacy and safety of memantine in patients with mild to moderate vascular dementia: a randomized, placebo-controlled trial (MMM300). *Stroke*. 2002;33:1834-1839.
25. Greenamyre JT, Maragos WF, Albin RL, Penney JB, Young AB. Glutamate transmission and toxicity in Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry*. 1988;12:421-430.
26. Winblad B, Möbius HJ, Stöffler A. Glutamate receptors as a target for Alzheimer's disease: are clinical results supporting the hope? *J Neural Transm Suppl*. 2002;62:217-225.

ORIGINAL ARTICLE

Memantine in Moderate-to-Severe Alzheimer's Disease

Barry Reisberg, M.D., Rachele Doody, M.D., Ph.D., Albrecht Stöfler, M.D.,
Frederick Schmitt, Ph.D., Steven Ferris, Ph.D.,
and Hans Jörg Möbius, M.D., Ph.D., for the Memantine Study Group*

ABSTRACT

BACKGROUND

Overstimulation of the N-methyl-D-aspartate (NMDA) receptor by glutamate is implicated in neurodegenerative disorders. Accordingly, we investigated memantine, an NMDA antagonist, for the treatment of Alzheimer's disease.

METHODS

Patients with moderate-to-severe Alzheimer's disease were randomly assigned to receive placebo or 20 mg of memantine daily for 28 weeks. The primary efficacy variables were the Clinician's Interview-Based Impression of Change Plus Caregiver Input (CIBIC-Plus) and the Alzheimer's Disease Cooperative Study Activities of Daily Living Inventory modified for severe dementia (ADCS-ADLsev). The secondary efficacy end points included the Severe Impairment Battery and other measures of cognition, function, and behavior. Treatment differences between base line and the end point were assessed. Missing observations were imputed by using the most recent previous observation (the last observation carried forward). The results were also analyzed with only the observed values included, without replacing the missing values (observed-cases analysis).

RESULTS

Two hundred fifty-two patients (67 percent women; mean age, 76 years) from 32 U.S. centers were enrolled. Of these, 181 (72 percent) completed the study and were evaluated at week 28. Seventy-one patients discontinued treatment prematurely (42 taking placebo and 29 taking memantine). Patients receiving memantine had a better outcome than those receiving placebo, according to the results of the CIBIC-Plus ($P=0.06$ with the last observation carried forward, $P=0.03$ for observed cases), the ADCS-ADLsev ($P=0.02$ with the last observation carried forward, $P=0.003$ for observed cases), and the Severe Impairment Battery ($P<0.001$ with the last observation carried forward, $P=0.002$ for observed cases). Memantine was not associated with a significant frequency of adverse events.

CONCLUSIONS

Antiglutamatergic treatment reduced clinical deterioration in moderate-to-severe Alzheimer's disease, a phase associated with distress for patients and burden on caregivers, for which other treatments are not available.

From the Department of Psychiatry, New York University School of Medicine, New York (B.R., S.F.); the Department of Neurology, Baylor College of Medicine, Houston (R.D.); Merz Pharmaceuticals, Frankfurt, Germany (A.S., H.J.M.); and the Departments of Neurology and Psychiatry, University of Kentucky, Lexington (F.S.). Address reprint requests to Dr. Reisberg at the William and Sylvia Silberstein Aging and Dementia Research and Treatment Center, New York University School of Medicine, 550 First Ave., New York, NY 10016, or at barry.reisberg@med.nyu.edu.

*The members of the Memantine Study Group are listed in the Appendix.

N Engl J Med 2003;348:1333-41.
Copyright © 2003 Massachusetts Medical Society.

ALZHEIMER'S DISEASE AFFECTS AT least 15 million persons throughout the world.^{1,2} The number of persons with Alzheimer's disease is increasing substantially as populations age.³ As Alzheimer's disease advances, patients become progressively impaired in both cognitive and functional capacities,^{2,4} and the burden on caregivers increases. Pharmacologic treatments are currently approved for treating mild-to-moderate Alzheimer's disease.⁵ However, there are no treatments for the more advanced stages of Alzheimer's disease.

Glutamate is the principal excitatory neurotransmitter in the brain.^{6,7} Glutamatergic overstimulation may result in neuronal damage, a phenomenon that has been termed excitotoxicity. Such excitotoxicity ultimately leads to neuronal calcium overload and has been implicated in neurodegenerative disorders.⁸ Glutamate stimulates a number of postsynaptic receptors, including the N-methyl-D-aspartate (NMDA) receptor, which has been particularly implicated in memory processes, dementia, and the pathogenesis of Alzheimer's disease.⁹⁻¹¹

Memantine, an uncompetitive NMDA-receptor antagonist, could be of therapeutic value in Alzheimer's disease.¹² A recent study in patients with advanced dementia (Alzheimer's disease and vascular dementia) suggested therapeutic benefits.¹³ Accordingly, we conducted a trial of the efficacy of memantine in outpatients with moderate-to-severe Alzheimer's disease.

METHODS

PATIENTS

Patients at least 50 years old who were residing in the community and had probable Alzheimer's disease according to the criteria of the *Diagnostic and Statistical Manual of Mental Disorders*, fourth edition (DSM-IV)⁴ and of the National Institute of Neurologic and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association¹⁴ were recruited. The eligibility criteria included base-line Mini-Mental State Examination scores of 3 to 14,¹⁵ a stage of 5 or 6 on the Global Deterioration Scale,¹⁶ and a stage of 6a or greater on the Functional Assessment Staging instrument,¹⁷ signifying the presence of dementia-related deficits in the ability to perform one or more basic activities of daily living. The patients had reliable caregivers and had undergone computed tomography (CT) or magnetic resonance imaging (MRI) of the brain within the previous 12 months.

Patients with vascular dementia, dementia or clinically significant neurologic disease due to conditions other than Alzheimer's disease, major depressive disorder, or a score greater than 4 on the modified Hachinski Ischemic Rating Scale¹⁸ were excluded. Patients with clinically significant coexisting medical conditions or laboratory abnormalities were also excluded, as were patients receiving specific concomitant medications (anticonvulsant agents, antiparkinsonian agents, hypnotic agents, anxiolytic agents, neuroleptic agents, cholinomimetic agents, or any other investigational compounds). Patients who had been receiving stable antidepressant treatment for at least two months were eligible, and chloral hydrate could be used as a sedative or hypnotic, but not within 24 hours before an assessment.

STUDY DESIGN

In this 28-week, double-blind, parallel-group study, patients were randomly assigned to receive either memantine (20 mg per day; Merz) or an identical-appearing placebo. Randomization was stratified according to site with the use of RanCode (version 3.1) and in blocks of four, with staff at the individual sites blinded to the randomization process. The assigned treatment was discontinued if continuation represented a medical risk in the opinion of the study physician, or if the patient declined ongoing participation. Subjects who withdrew prematurely were asked to complete end-point measures at the time of early termination and to return at 28 weeks for a "retrieved-dropout visit," which included all end-point assessments. The results of an optional 24-week open-label study extension, in which all patients took memantine, are still being analyzed.

Thirty-two U.S. centers participated. The trial was conducted in compliance with the Declaration of Helsinki and its amendments and was approved by study-center institutional review boards. The eligible patients and responsible caregivers provided written informed consent.

Merz Pharmaceuticals provided study medication and funding and was involved in planning the design and protocol. Data analysis was performed by a contract research organization (Quintiles), along with the authors. The data are stored at Merz Pharmaceuticals.

EFFICACY VARIABLES

The prespecified primary efficacy variables were the Clinician's Interview-Based Impression of Change Plus Caregiver Input (CIBIC-Plus) global

score (New York University version)¹⁹ at 28 weeks and the change from base line to week 28 in the Alzheimer's Disease Cooperative Study Activities of Daily Living Inventory (ADCS-ADL)²⁰ modified for more severe dementia (ADCS-ADLsev).²¹ If values from the 28-week observation were not available, the last observed value was used. Assessments were conducted at base line, at mid-study (week 12), and at the end of treatment (week 28) or at early termination, with a 28-week retrieved-dropout visit when possible.

The CIBIC-Plus measures overall global change relative to base line and is scored on a seven-point scale ranging from 1 (markedly improved) to 7 (markedly worse). The domains of cognition (assessed by patient interview), function (caregiver interview), and behavior (separate patient and caregiver interviews) are systematically evaluated. Experienced clinicians, blinded to adverse events and other study assessments, conducted separate interviews with study patients and caregivers to assess overall change on the CIBIC-Plus. To ensure consistency, the same clinician completed all CIBIC-Plus interviews for each study patient and associated caregiver wherever possible.

The ADCS-ADL is a structured questionnaire originally created to assess functional capacity over a broad range of severity of dementia. Each item consists of a series of hierarchical questions designed to determine a patient's ability to perform one of the activities of daily living, ranging from total independence to total inability. A subgroup of 19 individually validated items (the ADCS-ADLsev) was used; a total score of 54 signified optimal performance, and lower scores indicated worse performance. Caregivers assessed a patient's activities during the preceding four-week interval. The differences in total scores were analyzed.

In addition, six other efficacy variables were measured. The Severe Impairment Battery^{22,23} was designed to evaluate cognitive performance in advanced Alzheimer's disease. A 51-item scale, it assesses social interaction, memory, language, visuospatial ability, attention, praxis, and construction. The scores range from 0 (greatest impairment) to 100. The Severe Impairment Battery was part of a predefined responder analysis. The Mini-Mental State Examination¹⁵ is a 30-point scale that measures cognitive function, with higher scores indicating better function. The Global Deterioration Scale¹⁶ is a seven-stage scale that assesses overall cognitive and functional capacity on the basis of ob-

servations of the patient and reports from the caregiver. Higher stages signify greater impairment.

The Functional Assessment Staging scale¹⁷ assesses the magnitude of progressive functional deterioration in patients with dementia by identifying characteristic progressive disabilities. Its seven major stages range from normal (stage 1) to severe dementia (stage 7). Five substages in stage 6, corresponding to the loss of ability to independently dress, bathe, and handle proper mechanics and cleanliness in using the toilet, and to remain continent, respectively, of urine and feces and six substages in stage 7, corresponding to the loss of speech, ambulation, and other motor capacities, are assessed. The Neuropsychiatric Inventory²⁴ assesses neuropsychiatric disturbances with a 12-item scale based on information from the caregiver regarding the patient's behavior and associated distress felt by the caregiver. The scores range from 0 to 144 for the patient-assessment rating and from 0 to 60 for the caregiver-distress rating, with 0 indicating the optimum in each case. The Resource Utilization in Dementia²⁵ instrument was designed to assess the burden on the caregiver and to provide Alzheimer's disease-related health economics data through structured interviews with caregivers. To assess the clinical relevance of treatment effects further, a multifactor responder analysis was predefined.

Measures of safety, assessed at specified intervals, included neurologic and physical examinations, measurement of vital signs, electrocardiography, laboratory tests (hematologic tests, blood chemical values, and urinalysis), and recording of adverse events.

STATISTICAL ANALYSIS

The main efficacy analysis was based on the randomized patients who received at least one assessment after base line. This analysis included both those who completed the study and those who discontinued their assigned treatment prematurely. For the latter, the efficacy observation at week 28 was imputed from the last available observation carried forward.²⁶ Three additional analyses were also performed to adjust for missing values. One analysis was identical to the primary analysis, except that the actual retrieved-dropout values at week 28 were used when available. A second analysis included patients for whom no value after base line was available in addition to the intention-to-treat population and assumed no change in the outcome

measures for these patients. In the third analysis, missing values were replaced for those with no value after base line by the mean observed value for decline in the placebo group. An observed-cases analysis was also performed based on data for all randomized patients who were available for evaluation at week 28.

Efficacy outcomes were analyzed by application of the Wilcoxon–Mann–Whitney test for independent samples to the change from base line. There were no interim analyses. The prespecified group with an individual response was defined as the patients who improved or had no deterioration on the CIBIC-Plus and who improved or had no deterioration on either the ADCS-ADLsev or the Severe Impairment Battery. All patients were included in the safety analysis. All reported P values are two-sided.

RESULTS

STUDY POPULATION

Of 345 patients screened between August 1998 and April 1999 at 32 U.S. centers, 252 were randomly

assigned to study groups. Seventy-one of the patients (42 of the 126 assigned to placebo and 29 of the 126 assigned to memantine) discontinued their assigned treatment before week 28, and the remaining 181 completed the double-blind portion of the study. Five patients were excluded from the analysis of the ADCS-ADLsev results and 16 from the analysis of the CIBIC-Plus results because they had not been assessed after the base-line assessment. The mean (\pm SD) duration of treatment for both groups was 24 ± 8 weeks. Only 5 of the 71 patients who left the study returned for a retrieved-dropout visit at week 28. Premature discontinuations were due to adverse events in 22 of the patients in the placebo group (17 percent) and 13 of the patients in the memantine group (10 percent). Other major reasons for discontinuation included the patient's refusal of ongoing participation (14 patients receiving placebo [11 percent] and 12 patients receiving memantine [10 percent]), death (4 patients receiving placebo [3 percent] and 1 patient receiving memantine [1 percent]), protocol violation (3 patients receiving placebo [2 percent] and 3 patients receiving memantine [2 percent]), and change of caregiver (2 patients receiving placebo [2 percent] and none receiving memantine). Patients could have multiple reasons for discontinuation.

The base-line characteristics were similar in the two treatment groups (Table 1). Of the randomized patients, 67 percent were female, and the mean age was 76 years. The mean base-line score on the Mini-Mental State Examination for the study population was 7.9.

EFFICACY

The base-line scores and the results based on analyses with the last observation carried forward and analyses of observed cases for the efficacy variables are shown in Table 2. The CIBIC-Plus ratings at the end point (mean difference between the groups, 0.3; $P=0.06$) and week 28 (mean difference, 0.3; $P=0.03$) supported the effectiveness of memantine (Fig. 1A).

The total ADCS-ADLsev scores at base line were similar in the two groups (27.4 in the placebo group and 26.8 in the memantine group) (Table 2). At the end point and at week 28 (Fig. 1B), there was significantly less deterioration in the memantine group than in the placebo group (in the analysis with the last observation carried forward, the mean difference was 2.1 [$P=0.02$]; in the observed-cases analysis, the mean difference was 3.4 [$P=0.003$]).

Table 1. Base-Line Characteristics of the Intention-to-Treat Population.*

Characteristic	Memantine (N=126)		Placebo (N=126)		Total (N=252)
	Completed Study (N=97)	Did Not Complete Study (N=29)	Completed Study (N=84)	Did Not Complete Study (N=42)	
Sex — no. (%)					
Female	70 (72.2)	21 (72.4)	55 (65.5)	24 (57.1)	170 (67.5)
Male	27 (27.8)	8 (27.6)	29 (34.5)	18 (42.9)	82 (32.5)
Age — yr	75.5 \pm 8.16	77.3 \pm 9.17	75.8 \pm 7.28	77.5 \pm 8.61	76.1 \pm 8.07
Education — yr	12.3 \pm 3.06	13.0 \pm 3.14	12.9 \pm 3.14	11.7 \pm 2.91	12.5 \pm 3.09
Race — no. (%)					
White	85 (87.6)	27 (93.1)	75 (89.3)	40 (95.2)	227 (90.1)
Black	4 (4.1)	1 (3.4)	4 (4.8)	2 (4.8)	11 (4.4)
Other	8 (8.2)	1 (3.4)	5 (6.0)	0	14 (5.6)
MMSE score	7.8 \pm 3.76	7.6 \pm 3.67	8.1 \pm 3.60	7.9 \pm 3.54	7.9 \pm 3.64
GDS stage — no. (%)					
5	46 (47.4)	13 (44.8)	41 (48.8)	12 (28.6)	112 (44.4)
6	51 (52.6)	16 (55.2)	43 (51.2)	30 (71.4)	140 (55.6)

* The intention-to-treat population included all randomized patients. Plus-minus values are means \pm SD. MMSE denotes Mini-Mental State Examination, and GDS Global Deterioration Scale.

The Severe Impairment Battery showed significant differences favoring memantine ($P<0.001$ with the last observation carried forward, $P=0.002$ for observed cases) (Fig. 2A). On the basis of the predetermined definition of a response in the study protocol, 29 percent of the patients receiving memantine and 10 percent of those receiving placebo had a response ($P<0.001$).

Memantine-treated patients showed significantly less deterioration in their functional Alzheimer's disease stage, as measured by the Functional Assessment Staging score ($P=0.02$ with the last observation carried forward, $P=0.007$ for observed

cases) (Fig. 2B). In the analysis of the intention-to-treat population with the last observation carried forward and at week 28 no significant differences were observed between treatment groups in the Mini-Mental State Examination score, Global Deterioration Scale stage, or Neuropsychiatric Inventory score.

Additional analyses were performed with different strategies used for missing values, as described above. The results were unchanged in each of these analyses.

A subgroup analysis examined whether efficacy was seen in both patients with moderate Alzheimer's

Table 2. Results of the Efficacy Analysis.*

Measure	Base Line		Analysis with Last Observation Carried Forward (Change from Base Line at End Point)			Analysis of Observed Cases (Change from Base Line at Week 28)		
	Memantine	Placebo	Memantine	Placebo	P Value†	Memantine	Placebo	P Value†
CIBIC-Plus‡								
Score	NA	NA	4.5±1.12	4.8±1.09	0.06	4.4±1.12	4.7±1.13	0.03
No. of patients	126	126	118	118		97	84	
ADCS-ADLsev§								
Score	26.8	27.4	-3.1±6.79	-5.2±6.33	0.02	-2.5±6.27	-5.9±6.78	0.003
No. of patients	126	126	124	123		97	84	
SIB								
Score	65.9	68.3	-4.0±11.34	-10.1±13.50	<0.001	-4.5±11.48	-10.2±12.66	0.002
No. of patients	126	126	124	123		96	83	
MMSE								
Score	7.7	8.1	-0.5±2.40	-1.2±3.02	0.18	-0.6±2.61	-0.9±3.09	0.68
No. of patients	126	126	124	124		97	82	
FAST¶								
Score	2.8	2.8	-0.2±1.24	0.6±1.39	0.02	0.1±1.24	0.5±1.38	0.007
No. of patients	126	126	121	118		97	84	
GDS								
Score	5.5	5.6	0.1±0.47	0.2±0.48	0.11	0.1±0.49	0.2±0.48	0.16
No. of patients	126	126	121	119		97	84	
NPI								
Score	21.4	19.5	0.5±15.76	3.8±16.06	0.33	0.1±15.92	2.9±16.13	0.60
No. of patients	126	126	120	119		97	84	

* CI denotes confidence interval, CIBIC-Plus Clinician's Interview-Based Impression of Change Plus Caregiver Input, NA not applicable, ADCS-ADLsev Alzheimer's Disease Cooperative Study Activities of Daily Living Inventory (modified for severe dementia), SIB Severe Impairment Battery, MMSE Mini-Mental State Examination, FAST Functional Assessment Staging, GDS Global Deterioration Scale, and NPI Neuropsychiatric Inventory. The analysis with the last observation carried forward was performed with the intention-to-treat population. For this analysis, the numbers of patients fulfilling the intention-to-treat criteria are given in parentheses. The observed-cases analysis was performed with 181 patients observed at week 28. Plus-minus values are means ±SD.



† P values are based on the Wilcoxon-Mann-Whitney test for between-treatment comparisons.

‡ The CIBIC-Plus is a change score. By design the base-line score, "no change," is set at 4.00. Higher values at subsequent measurements indicate worsening. The end-point and week 28 values are actual mean ratings. The 95 percent confidence intervals for the differences between groups were -0.51 to 0.02 for the change from base line at the end point and -0.69 to -0.03 for the change from base line at week 28.

§ The 95 percent confidence intervals for the differences between groups were 0.49 to 3.78 for the change from base line at the end point and 1.45 to 5.28 for the change from base line at week 28.

¶ The FAST scores were calculated by enumerating the FAST stages and substages as follows: stage 3 (a score of -2) through 5 (0) and substage 6a (1) through substage 7f (11).¹⁷

|| The NPI scores are from the patient assessments.



disease (Mini-Mental State Examination score, 10 to 14) and those with severe Alzheimer's disease (Mini-Mental State Examination score, less than 10). A benefit of memantine as compared with placebo was suggested for all outcome measures in both groups.

The required caregiver time, as assessed by the Resource Utilization in Dementia score, was analyzed in the intention-to-treat population with the last observation carried forward. The result was statistically significant, indicating that caregivers spent less time with patients receiving memantine (difference between treatment groups, 45.8 hours per month; 95 percent confidence interval, 10.37 to 81.27; $P=0.01$).

SAFETY AND TOLERABILITY

As expected in this population with moderate-to-severe illness, the majority of patients had adverse events during the study (84 percent with memantine and 87 percent with placebo). However, most adverse events were mild to moderate in severity and were either not related or unlikely to be related to the study medication (Table 3). The incidence rates for the frequently reported adverse events in the memantine group were no more than 2 percent higher than in the placebo group. There were no clinically relevant differences between patients in the memantine and placebo groups in base-line assessments of clinical laboratory values, electrocardiographic results, or measurements of vital signs.

More patients receiving placebo than patients receiving memantine discontinued the study prematurely because of adverse events (22 [17 percent] vs. 13 [10 percent]). Agitation was the most common reason for discontinuation (7 percent of those receiving placebo and 5 percent of those receiving memantine). Serious adverse events were reported in 23 patients receiving placebo (18 percent) and 16 patients receiving memantine (13 percent). There were seven deaths, two of which occurred in the memantine group. Two of these patients died within the 30-day period after the last dose of study medication. Most of the serious adverse events, including all of the deaths, were considered to be unrelated to the study medication.

DISCUSSION

This study provides evidence that modulation of NMDA receptors to reduce glutamate-induced excitotoxicity alleviates the symptoms of Alzheimer's

Table 3. Most Frequently Reported Adverse Events.*

Adverse Event	Memantine (N=126)	Placebo (N=126)
	<i>no. of patients (%)</i>	
Any adverse event	106 (84)	109 (87)
Agitation	23 (18)	40 (32)
Urinary incontinence	14 (11)	14 (11)
Urinary tract infection	7 (6)	17 (13)
Insomnia	13 (10)	10 (8)
Diarrhea	12 (10)	10 (8)

* Adverse events occurring in at least 10 percent of the patients in either treatment group are reported.

disease. This novel neurochemical approach is distinct from the cholinomimetic mechanism of all currently approved treatments for Alzheimer's disease.

This trial studied patients whose moderate-to-severe Alzheimer's disease compromised their ability to perform both instrumental and basic activities of daily living independently.¹⁷ More than 95 percent of patients were in Functional Assessment Staging stage 6. All patients had difficulty putting on clothing independently, and many also had difficulties with handling the mechanics of bathing and toilet use, and some patients also had difficulties maintaining continence. The significant differences observed in favor of the memantine group on the ADCS-ADLsev, the Functional Assessment Staging ratings, and the Severe Impairment Battery suggest reduced decline in these critical capacities, which was apparent in the global assessment of the patients (CIBIC-Plus in the observed-cases analysis).

The clinical relevance of treatment effects has been an issue in all trials of medication for Alzheimer's disease. Point differences between drug- and placebo-treated patients on quantitative scales do not necessarily indicate that these effects are clinically meaningful. Response analyses (rates of individual response) are often performed to illustrate the clinical relevance of results. In the present study, a significant difference in the predefined criterion for a response, which incorporated multiple end points, was observed. The treatment effects seen in the areas of cognition and function seemed to translate into improvements in the patients' behavior (less agitation in the adverse-events reports) and

mitigation of the burden on caregivers (fewer hours spent assisting the patient).

The results of the present study cannot be directly compared with those of studies of other anti-dementia drugs (tacrine, donepezil, rivastigmine, and galantamine); virtually all published studies with these compounds have been performed in patients with mild-to-moderate Alzheimer's disease. An exception is a recent study of donepezil by Feldman et al.²⁷ However, even that study included patients with less severe disease (the mean Mini-Mental State Examination score at base line was 12) than those in the present trial, whose mean Mini-Mental State Examination score at base line was less than 8. The treatment effects of memantine in the present study and donepezil in the study by Feldman et al. are of similar size for the common end points, the CIBIC-Plus and the Severe Impairment Battery. Cholinergic compounds have gastrointestinal side effects, whereas the tolerability of memantine in this study was found to be excellent. Future studies will need to examine whether memantine treatment and cholinergic treatment may ultimately prove to be complementary or even synergistic.

There are notable limitations to this study. The dropout rate for the total study population was 28 percent, which is probably attributable to the relatively severe stage of disease in the study patients. The withdrawal rate was higher in the placebo group (33 percent) than in the memantine group (23 percent). The failure to obtain information at week 28 for the majority of patients who discontinued the study prematurely limits the interpretation of the results. However, the average duration of randomly assigned therapy for patients in this 28-week trial was 24 weeks for both study groups. In three analyses, the effects of different strategies for replacing missing observations for all patients assigned to treatment were investigated, and the results did not change materially. Thus, our data indicate that memantine reduces decline in patients with moderate-to-severe Alzheimer's disease.

Supported by Merz Pharmaceuticals, Frankfurt, Germany. Methodologic procedures and development of measurements were supported by grants (AG-03051, AG-08051, and 1-U01 AG-10483) from the National Institute on Aging of the National Institutes of Health.

Dr. Reisberg, Dr. Doody, and Dr. Ferris report having received lecture fees from Lundbeck. Dr. Ferris reports having received lecture fees from Forest and Merz. Dr. Doody and Dr. Ferris report having received consulting fees from Forest and Lundbeck. Dr. Schmitt reports having received consulting fees from Forest.

We are indebted to Yvonne Wirth, M.D., for statistical support.

APPENDIX

The following were members of the Memantine Study Group: J.T. Apter, Princeton Biomedical Research, Princeton, N.J.; B. Baumel, Baumel-Eisner Neuromedical Institute, Fort Lauderdale, Fla.; C. Bernick, Nevada School of Medicine, Las Vegas; J.S. Carman, Carman Research, Smyrna, Ga.; L.P. Charles, Southern New Jersey Medical Institute, Stratford; J. Corey-Bloom, University of California, San Diego; H. Cummins, Institute for Advanced Clinical Research, Elkins Park, Pa.; C. DeCarli, University of Kansas Medical Center, Kansas City; R. Duara, Wien Center for Alzheimer's Disease, Miami Beach, Fla.; E. DuBoff, Denver Center for Medical Research, Denver; N. Edwards, University of Tennessee, Memphis; L. Eisner, Baumel-Eisner Neuromedical Institute, Fort Lauderdale, Fla.; M.R. Farlow, Indiana University Center for Alzheimer's Disease, Indianapolis; S. Flitman, Neurology Group, Phoenix, Ariz.; R.H. Hubbard, Southwest Institute for Clinical Research, Rancho Mirage, Calif.; A. Jacobson, Neurological Consultants, Hollywood, Fla.; C.L. Jurkowski, Hampton Hospital, Westhampton Township, N.J.; A. Kiev, Life Span Develop Mental Systems, Poughkeepsie, N.Y.; L.C. Kirby II, Medici Research Centers, Peoria, Ariz.; D. Margolin, 6335 N. Fresno St., Fresno, Calif.; C. Merideth, Affiliated Research Institute, San Diego, Calif.; J.E. Mintzer, Alzheimer's Research and Clinical Program, Charleston, S.C.; E. Pfeiffer, University of South Florida Suncoast Gerontology Center, Tampa; R. Richter, St. John's Doctor's Building, Tulsa, Okla.; C.H. Sadowsky, Premiere Research Institute, West Palm Beach, Fla.; P. Solomon, Memory Clinic, Bennington, Vt.; S. Targum, Clinical Studies, Philadelphia; H. Tilker, Four Rivers Clinical Research, Paducah, Ky.; and M. Usman, Alzheimer's Center of Pittsburgh, Pittsburgh.

REFERENCES

1. Khachaturian ZS. Plundered memories. *Sciences* 1997;37(4):20-5.
2. Reisberg B, Burns A, Broadly H, et al. Diagnosis of Alzheimer's disease: report of an International Psychogeriatric Association Special Meeting Work Group under the cosponsorship of Alzheimer's Disease International, the European Federation of Neurological Societies, the World Health Organization, and the World Psychiatric Association. *Int Psychogeriatr* 1997;9:Suppl 1:11-38.
3. Henderson AS, Jorm AF. Definition and epidemiology of dementia: a review. In: Maj M, Sartorius N, eds. *Dementia*. 2nd ed. Chichester, England: John Wiley, 2002:1-33.
4. Diagnostic and statistical manual of mental disorders, 4th ed.: DSM-IV. Washington, D.C.: American Psychiatric Association, 1994.
5. Physicians' desk reference. 56th ed. Montvale, N.J.: Medical Economics, 2002: 1270-3, 1351-5, 1792-6, 2342-8, 2665-7.
6. Fonnum F. Glutamate: a neurotransmitter in mammalian brain. *J Neurochem* 1984; 42:1-11.
7. Orrego F, Villanueva S. The chemical nature of the main central excitatory transmitter: a critical appraisal based upon release studies and synaptic vesicle localization. *Neuroscience* 1993;56:539-55.
8. Lipton SA, Rosenberg PA. Excitatory amino acids as a final common pathway for neurologic disorders. *N Engl J Med* 1994; 330:613-22.
9. Shimizu E, Tang YP, Rampon C, Tsien JZ. NMDA receptor-dependent synaptic reinforcement is a crucial process for memory consolidation. *Science* 2000;290:1170-4. [Erratum, *Science* 2001;291:1902.]
10. Ackerley S, Grierson AJ, Brownlee J, et al. Glutamate slows axonal transport of neurofilaments in transfected neurons. *J Cell Biol* 2000;150:165-76.
11. Farber NB, Newcomer JW, Olney JW. The glutamate synapse in neuropsychiatric disorders: focus on schizophrenia and Alzheimer's disease. *Prog Brain Res* 1998;116: 421-37.

12. Danysz W, Parsons CG, Möbius HJ, Stöf-
fler A, Quack G. Neuroprotective and sym-
ptomalogical action of memantine relevant for
Alzheimer's disease — a unified glutamater-
gic hypothesis on the mechanism of action.
Neurotoxicity Res 2000;2:85-98.
13. Winblad B, Poritis N. Memantine in
severe dementia: results of the 9M-BEST
Study (benefit and efficacy in severely de-
mented patients during treatment with
memantine). *Int J Geriatr Psychiatry* 1999;
14:135-46.
14. McKhann G, Drachman D, Folstein M,
Katzman R, Price D, Stadlan EM. Clinical
diagnosis of Alzheimer's disease: report of
the NINCDS-ADRDA Work Group under
the auspices of Department of Health and
Human Services Task Force on Alzheimer's
Disease. *Neurology* 1984;34:939-44.
15. Folstein MF, Folstein SE, McHugh PR.
"Mini-Mental State": a practical method for
grading the cognitive state of patients for the
clinician. *J Psychiatr Res* 1975;12:189-98.
16. Reisberg B, Ferris SH, de Leon MJ, Crook
T. The Global Deterioration Scale for assess-
ment of primary degenerative dementia. *Am
J Psychiatry* 1982;139:1136-9.
17. Sclan SG, Reisberg B. Functional assess-
ment staging (FAST) in Alzheimer's disease:
reliability, validity, and ordinality. *Int Psy-
chogeriatr* 1992;4:Suppl 1:55-69.
18. Rosen WG, Terry RD, Fuld PA, Katzman
R, Peck A. Pathological verification of ische-
mic score in differentiation of dementias.
Ann Neurol 1980;7:486-8.
19. Reisberg B, Schneider L, Doody R, et al.
Clinical global measures of dementia: posi-
tion paper from the International Working
Group on Harmonization of Dementia Drug
Guidelines. *Alzheimer Dis Assoc Disord*
1997;11:Suppl 3:8-18.
20. Galasko D, Bennett D, Sano M, et al. An
inventory to assess activities of daily living
for clinical trials in Alzheimer's disease: the
Alzheimer's Disease Cooperative Study. *Alz-
heimer Dis Assoc Disord* 1997;11:Suppl 2:
S33-S39.
21. Galasko DR, Schmitt FA, Jin S, et al.
Detailed assessment of cognition and activ-
ities of daily living in moderate to severe Alz-
heimer's disease. *Neurobiol Aging* 2000;21:
Suppl 1:S168. abstract.
22. Panisset M, Roudier M, Saxton J, Boller
F. Severe Impairment Battery: a neuropsychol-
ogical test for severely demented patients.
Arch Neurol 1994;51:41-5.
23. Schmitt FA, Ashford W, Ernesto C, et al.
The Severe Impairment Battery: concurrent
validity and assessment of longitudinal
change in Alzheimer's disease: the Alzhei-
mer's Disease Cooperative Study. *Alzheimer
Dis Assoc Disord* 1997;11:Suppl 2:S51-S56.
24. Cummings JL, Mega M, Gray K, Rosen-
berg-Thompson S, Carusi DA, Gornbein J.
The Neuropsychiatric Inventory: compre-
hensive assessment of psychopathology in
dementia. *Neurology* 1994;44:2308-14.
25. Wimo A, Wetterholm AL, Mastey V, Win-
blad B. Evaluation of the healthcare resource
utilization and caregiver time in anti-demen-
tia drug trials. In: Wimo A, Jönsson B, Karls-
son G, Winblad B, eds. *Health economics of
dementia*. Chichester, England: John Wiley,
1998:465-99.
26. Gillings D, Koch G. The application of
the principle of intent-to-treat to the analysis
of clinical trials. *Drug Inf J* 1991;25:411-24.
27. Feldman H, Gauthier S, Hecker J, Vellas
B, Subbiah P, Whalen E. A 24-week, random-
ized, double-blind study of donepezil in
moderate to severe Alzheimer's disease. *Neu-
rology* 2001;57:613-20. [Erratum, *Neurology*
2001;57:2153.]

Copyright © 2003 Massachusetts Medical Society.

ELECTRONIC ACCESS TO THE JOURNAL'S CUMULATIVE INDEX

At the Journal's site on the World Wide Web (<http://www.nejm.org>) you can search an index of all articles published since January 1975 (abstracts 1975-1992, full-text 1993-present). You can search by author, key word, title, type of article, and date. The results will include the citations for the articles plus links to the abstracts of articles published since 1993. For nonsubscribers, time-limited access to single articles and 24-hour site access can also be ordered for a fee through the Internet (<http://www.nejm.org>).

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.